

# Package ‘actuar’

January 6, 2021

**Type** Package

**Title** Actuarial Functions and Heavy Tailed Distributions

**Version** 3.1-0

**Date** 2021-01-05

**Description** Functions and data sets for actuarial science:  
modeling of loss distributions; risk theory and ruin theory;  
simulation of compound models, discrete mixtures and compound  
hierarchical models; credibility theory. Support for many additional  
probability distributions to model insurance loss size and  
frequency: 23 continuous heavy tailed distributions; the  
Poisson-inverse Gaussian discrete distribution; zero-truncated and  
zero-modified extensions of the standard discrete distributions.  
Support for phase-type distributions commonly used to compute ruin  
probabilities.

**Depends** R (>= 3.3.0)

**Imports** stats, graphics, expint

**LinkingTo** expint

**Suggests** MASS

**License** GPL (>= 2)

**URL** <https://gitlab.com/vigou3/actuar>

**BugReports** <https://gitlab.com/vigou3/actuar/-/issues>

**Encoding** UTF-8

**LazyData** yes

**Classification/MSC-2010** 62P05, 91B30, 62G32

**NeedsCompilation** yes

**Author** Vincent Goulet [cre, aut],  
Sébastien Auclair [ctb],  
Christophe Dutang [aut],  
Nicholas Langevin [ctb],  
Xavier Milhaud [ctb],

Tommy Ouellet [ctb],  
 Alexandre Parent [ctb],  
 Mathieu Pigeon [aut],  
 Louis-Philippe Pouliot [ctb],  
 Jeffrey A. Ryan [aut] (Package API),  
 Robert Gentleman [aut] (Parts of the R to C interface),  
 Ross Ihaka [aut] (Parts of the R to C interface),  
 R Core Team [aut] (Parts of the R to C interface),  
 R Foundation [aut] (Parts of the R to C interface)

**Maintainer** Vincent Goulet <vincent.goulet@act.ulaval.ca>

**Repository** CRAN

**Date/Publication** 2021-01-06 16:01:15 UTC

## R topics documented:

adjCoef	4
aggregateDist	6
betaint	11
BetaMoments	13
Burr	14
ChisqSupp	16
cm	18
coverage	24
CTE	26
dental	27
discretize	28
elev	30
emm	32
ExponentialSupp	33
Extract.grouped.data	35
FellerPareto	36
GammaSupp	39
gdental	40
GeneralizedBeta	40
GeneralizedPareto	42
grouped.data	45
Gumbel	47
hachemeister	49
hist.grouped.data	49
InverseBurr	51
InverseExponential	53
InverseGamma	54
InverseGaussian	56
InverseParalogistic	59
InversePareto	60
InverseTransformedGamma	62
InverseWeibull	64

Logarithmic . . . . .	66
Loggamma . . . . .	68
Loglogistic . . . . .	69
LognormalMoments . . . . .	71
mde . . . . .	72
mean.grouped.data . . . . .	74
NormalSupp . . . . .	75
ogive . . . . .	76
Paralogistic . . . . .	78
Pareto . . . . .	80
Pareto2 . . . . .	82
Pareto3 . . . . .	84
Pareto4 . . . . .	86
PhaseType . . . . .	88
PoissonInverseGaussian . . . . .	90
quantile.aggregateDist . . . . .	93
quantile.grouped.data . . . . .	94
rcompound . . . . .	95
rmixture . . . . .	97
ruin . . . . .	98
severity . . . . .	100
simul . . . . .	101
simul.summaries . . . . .	104
SingleParameterPareto . . . . .	107
TransformedBeta . . . . .	108
TransformedGamma . . . . .	111
UniformSupp . . . . .	113
unroll . . . . .	114
VaR . . . . .	115
WeibullMoments . . . . .	116
ZeroModifiedBinomial . . . . .	117
ZeroModifiedGeometric . . . . .	119
ZeroModifiedLogarithmic . . . . .	121
ZeroModifiedNegativeBinomial . . . . .	123
ZeroModifiedPoisson . . . . .	125
ZeroTruncatedBinomial . . . . .	127
ZeroTruncatedGeometric . . . . .	129
ZeroTruncatedNegativeBinomial . . . . .	131
ZeroTruncatedPoisson . . . . .	133



## Details

In the typical case reinsurance = "none", the coefficient of determination is the smallest (strictly) positive root of the Lundberg equation

$$h(x) = E[e^{xB-xcW}] = 1$$

on  $[0, m)$ , where  $m = \text{upper.bound}$ ,  $B$  is the claim severity random variable,  $W$  is the claim interarrival (or wait) time random variable and  $c = \text{premium.rate}$ . The premium rate must satisfy the positive safety loading constraint  $E[B - cW] < 0$ .

With reinsurance = "proportional", the equation becomes

$$h(x, y) = E[e^{xyB-xc(y)W}] = 1,$$

where  $y$  is the retention rate and  $c(y)$  is the premium rate function.

With reinsurance = "excess-of-loss", the equation becomes

$$h(x, y) = E[e^{x \min(B, y) - xc(y)W}] = 1,$$

where  $y$  is the retention limit and  $c(y)$  is the premium rate function.

One can use argument  $h$  as an alternative way to provide function  $h(x)$  or  $h(x, y)$ . This is necessary in cases where random variables  $B$  and  $W$  are not independent.

The root of  $h(x) = 1$  is found by minimizing  $(h(x) - 1)^2$ .

## Value

If reinsurance = "none", a numeric vector of length one. Otherwise, a function of class "adjCoef" inheriting from the "function" class.

## Author(s)

Christophe Dutang, Vincent Goulet <vincent.goulet@act.ulaval.ca>

## References

Bowers, N. J. J., Gerber, H. U., Hickman, J., Jones, D. and Nesbitt, C. (1986), *Actuarial Mathematics*, Society of Actuaries.

Centeno, M. d. L. (2002), Measuring the effects of reinsurance by the adjustment coefficient in the Sparre-Anderson model, *Insurance: Mathematics and Economics* **30**, 37–49.

Gerber, H. U. (1979), *An Introduction to Mathematical Risk Theory*, Huebner Foundation.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2008), *Loss Models, From Data to Decisions, Third Edition*, Wiley.

## Examples

```
## Basic example: no reinsurance, exponential claim severity and wait
## times, premium rate computed with expected value principle and
## safety loading of 20%.
adjCoef(mgfexp, premium = 1.2, upper = 1)
```

```

## Same thing, giving function h.
h <- function(x) 1/((1 - x) * (1 + 1.2 * x))
adjCoef(h = h, upper = 1)

## Example 11.4 of Klugman et al. (2008)
mgfx <- function(x) 0.6 * exp(x) + 0.4 * exp(2 * x)
adjCoef(mgfx(x), mgfexp(x, 4), prem = 7, upper = 0.3182)

## Proportional reinsurance, same assumptions as above, reinsurer's
## safety loading of 30%.
mgfx <- function(x, y) mgfexp(x * y)
p <- function(x) 1.3 * x - 0.1
h <- function(x, a) 1/((1 - a * x) * (1 + x * p(a)))
R1 <- adjCoef(mgfx, premium = p, upper = 1, reins = "proportional",
             from = 0, to = 1, n = 11)
R2 <- adjCoef(h = h, upper = 1, reins = "p",
             from = 0, to = 1, n = 101)
R1(seq(0, 1, length = 10)) # evaluation for various retention rates
R2(seq(0, 1, length = 10)) # same
plot(R1) # graphical representation
plot(R2, col = "green", add = TRUE) # smoother function

## Excess-of-loss reinsurance
p <- function(x) 1.3 * levgamma(x, 2, 2) - 0.1
mgfx <- function(x, l)
  mgfgamma(x, 2, 2) * pgamma(1, 2, 2 - x) +
  exp(x * l) * pgamma(1, 2, 2, lower = FALSE)
h <- function(x, l) mgfx(x, l) * mgfexp(-x * p(l))
R1 <- adjCoef(mgfx, upper = 1, premium = p, reins = "excess-of-loss",
             from = 0, to = 10, n = 11)
R2 <- adjCoef(h = h, upper = 1, reins = "e",
             from = 0, to = 10, n = 101)

plot(R1)
plot(R2, col = "green", add = TRUE)

```

---

 aggregateDist

*Aggregate Claim Amount Distribution*


---

### Description

Compute the aggregate claim amount cumulative distribution function of a portfolio over a period using one of five methods.

### Usage

```

aggregateDist(method = c("recursive", "convolution", "normal",
                        "npower", "simulation"),
             model.freq = NULL, model.sev = NULL, p0 = NULL,
             x.scale = 1, convolve = 0, moments, nb.simul, ...)

```

```

        tol = 1e-06, maxit = 500, echo = FALSE)

## S3 method for class 'aggregateDist'
print(x, ...)

## S3 method for class 'aggregateDist'
plot(x, xlim, ylab = expression(F[S](x)),
     main = "Aggregate Claim Amount Distribution",
     sub = comment(x), ...)

## S3 method for class 'aggregateDist'
summary(object, ...)

## S3 method for class 'aggregateDist'
mean(x, ...)

## S3 method for class 'aggregateDist'
diff(x, ...)

```

### Arguments

method	method to be used
model.freq	for "recursive" method: a character string giving the name of a distribution in the $(a, b, 0)$ or $(a, b, 1)$ families of distributions. For "convolution" method: a vector of claim number probabilities. For "simulation" method: a frequency simulation model (see <a href="#">rcomphierarc</a> for details) or NULL. Ignored with normal and npower methods.
model.sev	for "recursive" and "convolution" methods: a vector of claim amount probabilities. For "simulation" method: a severity simulation model (see <a href="#">rcomphierarc</a> for details) or NULL. Ignored with normal and npower methods.
p0	arbitrary probability at zero for the frequency distribution. Creates a zero-modified or zero-truncated distribution if not NULL. Used only with "recursive" method.
x.scale	value of an amount of 1 in the severity model (monetary unit). Used only with "recursive" and "convolution" methods.
convolve	number of times to convolve the resulting distribution with itself. Used only with "recursive" method.
moments	vector of the true moments of the aggregate claim amount distribution; required only by the "normal" or "npower" methods.
nb.simul	number of simulations for the "simulation" method.
...	parameters of the frequency distribution for the "recursive" method; further arguments to be passed to or from other methods otherwise.
tol	the resulting cumulative distribution in the "recursive" method will get less than tol away from 1.
maxit	maximum number of recursions in the "recursive" method.
echo	logical; echo the recursions to screen in the "recursive" method.

x, object	an object of class "aggregateDist".
xlim	numeric of length 2; the $x$ limits of the plot.
ylab	label of the y axis.
main	main title.
sub	subtitle, defaulting to the calculation method.

## Details

aggregateDist returns a function to compute the cumulative distribution function (cdf) of the aggregate claim amount distribution in any point.

The "recursive" method computes the cdf using the Panjer algorithm; the "convolution" method using convolutions; the "normal" method using a normal approximation; the "npower" method using the Normal Power 2 approximation; the "simulation" method using simulations. More details follow.

## Value

A function of class "aggregateDist", inheriting from the "function" class when using normal and Normal Power approximations and additionally inheriting from the "ecdf" and "stepfun" classes when other methods are used.

There are methods available to summarize (summary), represent (print), plot (plot), compute quantiles (quantile) and compute the mean (mean) of "aggregateDist" objects.

For the diff method: a numeric vector of probabilities corresponding to the probability mass function evaluated at the knots of the distribution.

## Recursive method

The frequency distribution must be a member of the  $(a, b, 0)$  or  $(a, b, 1)$  families of discrete distributions.

To use a distribution from the  $(a, b, 0)$  family, model.freq must be one of "binomial", "geometric", "negative binomial" or "poisson", and  $p_0$  must be NULL.

To use a zero-truncated distribution from the  $(a, b, 1)$  family, model.freq may be one of the strings above together with  $p_0 = 0$ . As a shortcut, model.freq may also be one of "zero-truncated binomial", "zero-truncated geometric", "zero-truncated negative binomial", "zero-truncated poisson" or "logarithmic", and  $p_0$  is then ignored (with a warning if non NULL).

(Note: since the logarithmic distribution is always zero-truncated. model.freq = "logarithmic" may be used with either  $p_0 = \text{NULL}$  or  $p_0 = 0$ .)

To use a zero-modified distribution from the  $(a, b, 1)$  family, model.freq may be one of standard frequency distributions mentioned above with  $p_0$  set to some probability that the distribution takes the value 0. It is equivalent, but more explicit, to set model.freq to one of "zero-modified binomial", "zero-modified geometric", "zero-modified negative binomial", "zero-modified poisson" or "zero-modified logarithmic".

The parameters of the frequency distribution must be specified using names identical to the arguments of the appropriate function `dbinom`, `dgeom`, `dnbinom`, `dpois` or `dlogarithmic`. In the latter case, do take note that the parametrization of `dlogarithmic` is different from Appendix B of Klugman et al. (2012).



If the length of `p0` is greater than one, only the first element is used, with a warning.

`model.sev` is a vector of the (discretized) claim amount distribution  $X$ ; the first element **must** be  $f_X(0) = \Pr[X = 0]$ .

The recursion will fail to start if the expected number of claims is too large. One may divide the appropriate parameter of the frequency distribution by  $2^n$  and convolve the resulting distribution  $n = \text{convolve times}$ .

Failure to obtain a cumulative distribution function less than `tol` away from 1 within `maxit` iterations is often due to too coarse a discretization of the severity distribution.

### Convolution method

The cumulative distribution function (cdf)  $F_S(x)$  of the aggregate claim amount of a portfolio in the collective risk model is

$$F_S(x) = \sum_{n=0}^{\infty} F_X^{*n}(x) p_n,$$

for  $x = 0, 1, \dots$ ;  $p_n = \Pr[N = n]$  is the frequency probability mass function and  $F_X^{*n}(x)$  is the cdf of the  $n$ th convolution of the (discrete) claim amount random variable.

`model.freq` is vector  $p_n$  of the number of claims probabilities; the first element **must** be  $\Pr[N = 0]$ .

`model.sev` is vector  $f_X(x)$  of the (discretized) claim amount distribution; the first element **must** be  $f_X(0)$ .

### Normal and Normal Power 2 methods

The Normal approximation of a cumulative distribution function (cdf)  $F(x)$  with mean  $\mu$  and standard deviation  $\sigma$  is

$$F(x) \approx \Phi\left(\frac{x - \mu}{\sigma}\right).$$

The Normal Power 2 approximation of a cumulative distribution function (cdf)  $F(x)$  with mean  $\mu$ , standard deviation  $\sigma$  and skewness  $\gamma$  is

$$F(x) \approx \Phi\left(-\frac{3}{\gamma} + \sqrt{\frac{9}{\gamma^2} + 1 + \frac{6}{\gamma} \frac{x - \mu}{\sigma}}\right).$$

This formula is valid only for the right-hand tail of the distribution and skewness should not exceed unity.

### Simulation method

This methods returns the empirical distribution function of a sample of size `nb.simul` of the aggregate claim amount distribution specified by `model.freq` and `model.sev`. `rcomphierarc` is used for the simulation of claim amounts, hence both the frequency and severity models can be mixtures of distributions.

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Louis-Philippe Pouliot

## References

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

Daykin, C.D., Pentikäinen, T. and Pesonen, M. (1994), *Practical Risk Theory for Actuaries*, Chapman & Hall.

## See Also

[discretize](#) to discretize a severity distribution; [mean.aggregateDist](#) to compute the mean of the distribution; [quantile.aggregateDist](#) to compute the quantiles or the Value-at-Risk; [CTE.aggregateDist](#) to compute the Conditional Tail Expectation (or Tail Value-at-Risk); [rcomphierarc](#).

## Examples

```
## Convolution method (example 9.5 of Klugman et al. (2012))
fx <- c(0, 0.15, 0.2, 0.25, 0.125, 0.075,
        0.05, 0.05, 0.05, 0.025, 0.025)
pn <- c(0.05, 0.1, 0.15, 0.2, 0.25, 0.15, 0.06, 0.03, 0.01)
Fs <- aggregateDist("convolution", model.freq = pn,
                   model.sev = fx, x.scale = 25)

summary(Fs)
c(Fs(0), diff(Fs(25 * 0:21))) # probability mass function
plot(Fs)

## Recursive method (example 9.10 of Klugman et al. (2012))
fx <- c(0, crossprod(c(2, 1)/3,
                    matrix(c(0.6, 0.7, 0.4, 0, 0, 0.3), 2, 3)))
Fs <- aggregateDist("recursive", model.freq = "poisson",
                   model.sev = fx, lambda = 3)

plot(Fs)
Fs(knots(Fs))      # cdf evaluated at its knots
diff(Fs)           # probability mass function

## Recursive method (high frequency)
fx <- c(0, 0.15, 0.2, 0.25, 0.125, 0.075,
        0.05, 0.05, 0.05, 0.025, 0.025)
## Not run: Fs <- aggregateDist("recursive", model.freq = "poisson",
                              model.sev = fx, lambda = 1000)

## End(Not run)
Fs <- aggregateDist("recursive", model.freq = "poisson",
                   model.sev = fx, lambda = 250, convolve = 2, maxit = 1500)

plot(Fs)

## Recursive method (zero-modified distribution; example 9.11 of
## Klugman et al. (2012))
Fn <- aggregateDist("recursive", model.freq = "binomial",
                   model.sev = c(0.3, 0.5, 0.2), x.scale = 50,
                   p0 = 0.4, size = 3, prob = 0.3)

diff(Fn)

## Equivalent but more explicit call
```

```

aggregateDist("recursive", model.freq = "zero-modified binomial",
              model.sev = c(0.3, 0.5, 0.2), x.scale = 50,
              p0 = 0.4, size = 3, prob = 0.3)

## Recursive method (zero-truncated distribution). Using 'fx' above
## would mean that both Pr[N = 0] = 0 and Pr[X = 0] = 0, therefore
## Pr[S = 0] = 0 and recursions would not start.
fx <- discretize(pexp(x, 1), from = 0, to = 100, method = "upper")
fx[1L] # non zero
aggregateDist("recursive", model.freq = "zero-truncated poisson",
              model.sev = fx, lambda = 3, x.scale = 25, echo=TRUE)

## Normal Power approximation
Fs <- aggregateDist("npower", moments = c(200, 200, 0.5))
Fs(210)

## Simulation method
model.freq <- expression(data = rpois(3))
model.sev <- expression(data = rgamma(100, 2))
Fs <- aggregateDist("simulation", nb.simul = 1000,
                   model.freq, model.sev)

mean(Fs)
plot(Fs)

## Evaluation of ruin probabilities using Beekman's formula with
## Exponential(1) claim severity, Poisson(1) frequency and premium rate
## c = 1.2.
fx <- discretize(pexp(x, 1), from = 0, to = 100, method = "lower")
phi0 <- 0.2/1.2
Fs <- aggregateDist(method = "recursive", model.freq = "geometric",
                   model.sev = fx, prob = phi0)
1 - Fs(400) # approximate ruin probability
u <- 0:100
plot(u, 1 - Fs(u), type = "l", main = "Ruin probability")

```

---

betaint

*The "Beta Integral"*


---

### Description

The "beta integral" which is just a multiple of the non regularized incomplete beta function. This function merely provides an R interface to the C level routine. It is not exported by the package.

### Usage

```
betaint(x, a, b)
```

### Arguments

x	vector of quantiles.
a, b	parameters. See Details for admissible values.

**Details**

Function `betaint` computes the “beta integral”

$$B(a, b; x) = \Gamma(a + b) \int_0^x t^{a-1} (1-t)^{b-1} dt$$

for  $a > 0$ ,  $b \neq -1, -2, \dots$  and  $0 < x < 1$ . (Here  $\Gamma(\alpha)$  is the function implemented by R’s `gamma()` and defined in its help.) When  $b > 0$ ,

$$B(a, b; x) = \Gamma(a)\Gamma(b)I_x(a, b),$$

where  $I_x(a, b)$  is `pbeta(x, a, b)`. When  $b < 0$ ,  $b \neq -1, -2, \dots$ , and  $a > 1 + [-b]$ ,

$$\begin{aligned} B(a, b; x) = & -\Gamma(a + b) \left[ \frac{x^{a-1}(1-x)^b}{b} + \frac{(a-1)x^{a-2}(1-x)^{b+1}}{b(b+1)} \right. \\ & + \dots + \left. \frac{(a-1) \cdots (a-r)x^{a-r-1}(1-x)^{b+r}}{b(b+1) \cdots (b+r)} \right] \\ & + \frac{(a-1) \cdots (a-r-1)}{b(b+1) \cdots (b+r)} \Gamma(a-r-1) \\ & \times \Gamma(b+r+1) I_x(a-r-1, b+r+1), \end{aligned}$$

where  $r = [-b]$ .

This function is used (at the C level) to compute the limited expected value for distributions of the transformed beta family; see, for example, [levtrbeta](#).

**Value**

The value of the integral.

Invalid arguments will result in return value NaN, with a warning.

**Note**

The need for this function in the package is well explained in the introduction of Appendix A of Klugman et al. (2012). See also chapter 6 and 15 of Abramowitz and Stegun (1972) for definitions and relations to the hypergeometric series.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>

**References**

- Abramowitz, M. and Stegun, I. A. (1972), *Handbook of Mathematical Functions*, Dover.
- Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**Examples**

```

x <- 0.3
a <- 7

## case with b > 0
b <- 2
actuar:::betaint(x, a, b)
gamma(a) * gamma(b) * pbeta(x, a, b) # same

## case with b < 0
b <- -2.2
r <- floor(-b) # r = 2
actuar:::betaint(x, a, b)

## "manual" calculation
s <- (x^(a-1) * (1-x)^b)/b +
  ((a-1) * x^(a-2) * (1-x)^(b+1))/(b * (b+1)) +
  ((a-1) * (a-2) * x^(a-3) * (1-x)^(b+2))/(b * (b+1) * (b+2))
-gamma(a+b) * s +
  (a-1)*(a-2)*(a-3) * gamma(a-r-1)/(b*(b+1)*(b+2)) *
  gamma(b+r+1)*pbeta(x, a-r-1, b+r+1)

```

BetaMoments

*Raw and Limited Moments of the Beta Distribution***Description**

Raw moments and limited moments for the (central) Beta distribution with parameters shape1 and shape2.

**Usage**

```

mbeta(order, shape1, shape2)
levbeta(limit, shape1, shape2, order = 1)

```

**Arguments**

order            order of the moment.  
limit            limit of the loss variable.  
shape1, shape2   positive parameters of the Beta distribution.

**Details**

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$  and the  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ ,  $k > -\alpha$ .

The noncentral beta distribution is not supported.

**Value**

`mbeta` gives the  $k$ th raw moment and `levbeta` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**See Also**

[Beta](#) for details on the beta distribution and functions `[dppr]beta`.

**Examples**

```
mbeta(2, 3, 4) - mbeta(1, 3, 4)^2
levbeta(10, 3, 4, order = 2)
```

---

Burr

*The Burr Distribution*

---

**Description**

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Burr distribution with parameters `shape1`, `shape2` and `scale`.

**Usage**

```
dburr(x, shape1, shape2, rate = 1, scale = 1/rate,
      log = FALSE)
pburr(q, shape1, shape2, rate = 1, scale = 1/rate,
      lower.tail = TRUE, log.p = FALSE)
qburr(p, shape1, shape2, rate = 1, scale = 1/rate,
      lower.tail = TRUE, log.p = FALSE)
rburr(n, shape1, shape2, rate = 1, scale = 1/rate)
mburr(order, shape1, shape2, rate = 1, scale = 1/rate)
levburr(limit, shape1, shape2, rate = 1, scale = 1/rate,
        order = 1)
```

**Arguments**

<code>x, q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>shape1, shape2, scale</code>	parameters. Must be strictly positive.
<code>rate</code>	an alternative way to specify the scale.
<code>log, log.p</code>	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
<code>order</code>	order of the moment.
<code>limit</code>	limit of the loss variable.

**Details**

The Burr distribution with parameters  $\text{shape1} = \alpha$ ,  $\text{shape2} = \gamma$  and  $\text{scale} = \theta$  has density:

$$f(x) = \frac{\alpha\gamma(x/\theta)^\gamma}{\theta[1 + (x/\theta)^\gamma]^{\alpha+1}}$$

for  $x > 0$ ,  $\alpha > 0$ ,  $\gamma > 0$  and  $\theta > 0$ .

The Burr is the distribution of the random variable

$$\theta \left( \frac{X}{1-X} \right)^{1/\gamma},$$

where  $X$  has a beta distribution with parameters 1 and  $\alpha$ .

The Burr distribution has the following special cases:

- A [Loglogistic](#) distribution when `shape1 == 1`;
- A [Paralogistic](#) distribution when `shape2 == shape1`;
- A [Pareto](#) distribution when `shape2 == 1`.

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $-\gamma < k < \alpha\gamma$ .

The  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ ,  $k > -\gamma$  and  $\alpha - k/\gamma$  not a negative integer.

**Value**

`dburr` gives the density, `pburr` gives the distribution function, `qburr` gives the quantile function, `rburr` generates random deviates, `mburr` gives the  $k$ th raw moment, and `levburr` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

**Note**

levburr computes the limited expected value using [betaint](#).

Distribution also known as the Burr Type XII or Singh-Maddala distribution. See also Kleiber and Kotz (2003) for alternative names and parametrizations.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**See Also**

[dpareto4](#) for an equivalent distribution with a location parameter.

**Examples**

```
exp(dburr(1, 2, 3, log = TRUE))
p <- (1:10)/10
pburr(qburr(p, 2, 3, 2), 2, 3, 2)

## variance
mburr(2, 2, 3, 1) - mburr(1, 2, 3, 1) ^ 2

## case with shape1 - order/shape2 > 0
levburr(10, 2, 3, 1, order = 2)

## case with shape1 - order/shape2 < 0
levburr(10, 1.5, 0.5, 1, order = 2)
```

---

ChisqSupp

*Moments and Moment Generating Function of the (non-central) Chi-Squared Distribution*

---

**Description**

Raw moments, limited moments and moment generating function for the chi-squared ( $\chi^2$ ) distribution with df degrees of freedom and optional non-centrality parameter ncp.



**Usage**

```
mchisq(order, df, ncp = 0)
levchisq(limit, df, ncp = 0, order = 1)
mgfchisq(t, df, ncp = 0, log= FALSE)
```

**Arguments**

order	order of the moment.
limit	limit of the loss variable.
df	degrees of freedom (non-negative, but can be non-integer).
ncp	non-centrality parameter (non-negative).
t	numeric vector.
log	logical; if TRUE, the cumulant generating function is returned.

**Details**

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ , the  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$  and the moment generating function is  $E[e^{tX}]$ .

Only integer moments are supported for the non central Chi-square distribution ( $ncp > 0$ ).

The limited expected value is supported for the centered Chi-square distribution ( $ncp = 0$ ).

**Value**

`mchisq` gives the  $k$ th raw moment, `levchisq` gives the  $k$ th moment of the limited loss variable, and `mgfchisq` gives the moment generating function in `t`.

Invalid arguments will result in return value NaN, with a warning.

**Author(s)**

Christophe Dutang, Vincent Goulet <vincent.goulet@act.ulaval.ca>

**References**

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

Johnson, N. L. and Kotz, S. (1970), *Continuous Univariate Distributions, Volume 1*, Wiley.

**See Also**

[Chisquare](#)

**Examples**

```
mchisq(2, 3, 4)
levchisq(10, 3, order = 2)
mgfchisq(0.25, 3, 2)
```

**Description**

Fit the following credibility models: Bühlmann, Bühlmann-Straub, hierarchical, regression (Hachemeister) or linear Bayes.

**Usage**

```
cm(formula, data, ratios, weights, subset,
   regformula = NULL, regdata, adj.intercept = FALSE,
   method = c("Buhlmann-Gisler", "Ohlsson", "iterative"),
   likelihood, ...,
   tol = sqrt(.Machine$double.eps), maxit = 100, echo = FALSE)

## S3 method for class 'cm'
print(x, ...)

## S3 method for class 'cm'
predict(object, levels = NULL, newdata, ...)

## S3 method for class 'cm'
summary(object, levels = NULL, newdata, ...)

## S3 method for class 'summary.cm'
print(x, ...)
```

**Arguments**

formula	character string "bayes" or an object of class " <a href="#">formula</a> ": a symbolic description of the model to be fit. The details of model specification are given below.
data	a matrix or a data frame containing the portfolio structure, the ratios or claim amounts and their associated weights, if any.
ratios	expression indicating the columns of data containing the ratios or claim amounts.
weights	expression indicating the columns of data containing the weights associated with ratios.
subset	an optional logical expression indicating a subset of observations to be used in the modeling process. All observations are included by default.
regformula	an object of class " <a href="#">formula</a> ": symbolic description of the regression component (see <a href="#">lm</a> for details). No left hand side is needed in the formula; if present it is ignored. If NULL, no regression is done on the data.
regdata	an optional data frame, list or environment (or object coercible by <a href="#">as.data.frame</a> to a data frame) containing the variables in the regression model.

<code>adj.intercept</code>	if TRUE, the intercept of the regression model is located at the barycenter of the regressor instead of the origin.
<code>method</code>	estimation method for the variance components of the model; see details below.
<code>likelihood</code>	a character string giving the name of the likelihood function in one of the supported linear Bayes cases; see details below.
<code>tol</code>	tolerance level for the stopping criteria for iterative estimation method.
<code>maxit</code>	maximum number of iterations in iterative estimation method.
<code>echo</code>	logical; whether to echo the iterative procedure or not.
<code>x, object</code>	an object of class "cm".
<code>levels</code>	character vector indicating the levels to predict or to include in the summary; if NULL all levels are included.
<code>newdata</code>	data frame containing the variables used to predict credibility regression models.
<code>...</code>	parameters of the prior distribution for cm; additional attributes to attach to the result for the <code>predict</code> and <code>summary</code> methods; further arguments to <code>format</code> for the <code>print.summary</code> method; unused for the <code>print</code> method.

## Details

`cm` is the unified front end for credibility models fitting. The function supports hierarchical models with any number of levels (with Bühlmann and Bühlmann-Straub models as special cases) and the regression model of Hachemeister. Usage of `cm` is similar to `lm` for these cases. `cm` can also fit linear Bayes models, in which case usage is much simplified; see the section on linear Bayes below.

When not "bayes", the `formula` argument symbolically describes the structure of the portfolio in the form *terms*. Each term is an interaction between risk factors contributing to the total variance of the portfolio data. Terms are separated by + operators and interactions within each term by :. For a portfolio divided first into sectors, then units and finally contracts, `formula` would be `~ sector + sector:unit + sector:unit:contract`, where `sector`, `unit` and `contract` are column names in `data`. In general, the formula should be of the form `~ a + a:b + a:b:c + a:b:c:d + ...`

If argument `regformula` is not NULL, the regression model of Hachemeister is fit to the data. The response is usually time. By default, the intercept of the model is located at time origin. If argument `adj.intercept` is TRUE, the intercept is moved to the (collective) barycenter of time, by orthogonalization of the design matrix. Note that the regression coefficients may be difficult to interpret in this case.

Arguments `ratios`, `weights` and `subset` are used like arguments `select`, `select` and `subset`, respectively, of function `subset`.

Data does not have to be sorted by level. Nodes with no data (complete lines of NA except for the portfolio structure) are allowed, with the restriction mentioned above.

## Value

Function `cm` computes the structure parameters estimators of the model specified in `formula`. The value returned is an object of class `cm`.

An object of class "cm" is a list with at least the following components:

<code>means</code>	a list containing, for each level, the vector of linearly sufficient statistics.
--------------------	--

weights	a list containing, for each level, the vector of total weights.
unbiased	a vector containing the unbiased variance components estimators, or NULL.
iterative	a vector containing the iterative variance components estimators, or NULL.
cred	for multi-level hierarchical models: a list containing, the vector of credibility factors for each level. For one-level models: an array or vector of credibility factors.
nodes	a list containing, for each level, the vector of the number of nodes in the level.
classification	the columns of data containing the portfolio classification structure.
ordering	a list containing, for each level, the affiliation of a node to the node of the level above.

Regression fits have in addition the following components:

adj.models	a list containing, for each node, the credibility adjusted regression model as obtained with <code>lm.fit</code> or <code>lm.wfit</code> .
transition	if <code>adj.intercept</code> is TRUE, a transition matrix from the basis of the orthogonal design matrix to the basis of the original design matrix.
terms	the <code>terms</code> object used.

The method of `predict` for objects of class "cm" computes the credibility premiums for the nodes of every level included in argument `levels` (all by default). Result is a list the same length as `levels` or the number of levels in `formula`, or an atomic vector for one-level models.

### Hierarchical models

The credibility premium at one level is a convex combination between the linearly sufficient statistic of a node and the credibility premium of the level above. (For the first level, the complement of credibility is given to the collective premium.) The linearly sufficient statistic of a node is the credibility weighted average of the data of the node, except at the last level, where natural weights are used. The credibility factor of node  $i$  is equal to

$$\frac{w_i}{w_i + a/b},$$

where  $w_i$  is the weight of the node used in the linearly sufficient statistic,  $a$  is the average within node variance and  $b$  is the average between node variance.

### Regression models

The credibility premium of node  $i$  is equal to

$$y' b_i^a,$$

where  $y$  is a matrix created from `newdata` and  $b_i^a$  is the vector of credibility adjusted regression coefficients of node  $i$ . The latter is given by

$$b_i^a = Z_i b_i + (I - Z_i) m,$$

where  $b_i$  is the vector of regression coefficients based on data of node  $i$  only,  $m$  is the vector of collective regression coefficients,  $Z_i$  is the credibility matrix and  $I$  is the identity matrix. The credibility matrix of node  $i$  is equal to

$$A^{-1}(A + s^2 S_i),$$

where  $S_i$  is the unscaled regression covariance matrix of the node,  $s^2$  is the average within node variance and  $A$  is the within node covariance matrix.

If the intercept is positioned at the barycenter of time, matrices  $S_i$  and  $A$  (and hence  $Z_i$ ) are diagonal. This amounts to use Bühlmann-Straub models for each regression coefficient.

Argument `newdata` provides the “future” value of the regressors for prediction purposes. It should be given as specified in [predict.lm](#).

### Variance components estimation

For hierarchical models, two sets of estimators of the variance components (other than the within node variance) are available: unbiased estimators and iterative estimators.

Unbiased estimators are based on sums of squares of the form

$$B_i = \sum_j w_{ij} (X_{ij} - \bar{X}_i)^2 - (J - 1)a$$

and constants of the form

$$c_i = w_i - \sum_j \frac{w_{ij}^2}{w_i},$$

where  $X_{ij}$  is the linearly sufficient statistic of level  $(ij)$ ;  $\bar{X}_i$  is the weighted average of the latter using weights  $w_{ij}$ ;  $w_i = \sum_j w_{ij}$ ;  $J$  is the effective number of nodes at level  $(ij)$ ;  $a$  is the within variance of this level. Weights  $w_{ij}$  are the natural weights at the lowest level, the sum of the natural weights the next level and the sum of the credibility factors for all upper levels.

The Bühlmann-Gisler estimators (method = "Buhlmann-Gisler") are given by

$$b = \frac{1}{I} \sum_i \max\left(\frac{B_i}{c_i}, 0\right),$$

that is the average of the per node variance estimators truncated at 0.

The Ohlsson estimators (method = "Ohlsson") are given by

$$b = \frac{\sum_i B_i}{\sum_i c_i},$$

that is the weighted average of the per node variance estimators without any truncation. Note that negative estimates will be truncated to zero for credibility factor calculations.

In the Bühlmann-Straub model, these estimators are equivalent.

Iterative estimators method = "iterative" are pseudo-estimators of the form

$$b = \frac{1}{d} \sum_i w_i (X_i - \bar{X})^2,$$

where  $X_i$  is the linearly sufficient statistic of one level,  $\bar{X}$  is the linearly sufficient statistic of the level above and  $d$  is the effective number of nodes at one level minus the effective number of nodes of the level above. The Ohlsson estimators are used as starting values.

For regression models, with the intercept at time origin, only iterative estimators are available. If method is different from "iterative", a warning is issued. With the intercept at the barycenter of time, the choice of estimators is the same as in the Bühlmann-Straub model.

### Linear Bayes

When formula is "bayes", the function computes pure Bayesian premiums for the following combinations of distributions where they are linear credibility premiums:

- $X|\Theta = \theta \sim \text{Poisson}(\theta)$  and  $\Theta \sim \text{Gamma}(\alpha, \lambda)$ ;
- $X|\Theta = \theta \sim \text{Exponential}(\theta)$  and  $\Theta \sim \text{Gamma}(\alpha, \lambda)$ ;
- $X|\Theta = \theta \sim \text{Gamma}(\tau, \theta)$  and  $\Theta \sim \text{Gamma}(\alpha, \lambda)$ ;
- $X|\Theta = \theta \sim \text{Normal}(\theta, \sigma_2^2)$  and  $\Theta \sim \text{Normal}(\mu, \sigma_1^2)$ ;
- $X|\Theta = \theta \sim \text{Bernoulli}(\theta)$  and  $\Theta \sim \text{Beta}(a, b)$ ;
- $X|\Theta = \theta \sim \text{Binomial}(\nu, \theta)$  and  $\Theta \sim \text{Beta}(a, b)$ ;
- $X|\Theta = \theta \sim \text{Geometric}(\theta)$  and  $\Theta \sim \text{Beta}(a, b)$ .
- $X|\Theta = \theta \sim \text{Negative Binomial}(r, \theta)$  and  $\Theta \sim \text{Beta}(a, b)$ .

The following combination is also supported:  $X|\Theta = \theta \sim \text{Single Parameter Pareto}(\theta)$  and  $\Theta \sim \text{Gamma}(\alpha, \lambda)$ . In this case, the Bayesian estimator not of the risk premium, but rather of parameter  $\theta$  is linear with a "credibility" factor that is not restricted to  $(0, 1)$ .

Argument `likelihood` identifies the distribution of  $X|\Theta = \theta$  as one of "poisson", "exponential", "gamma", "normal", "bernoulli", "binomial", "geometric", "negative binomial" or "pareto".

The parameters of the distributions of  $X|\Theta = \theta$  (when needed) and  $\Theta$  are set in ... using the argument names (and default values) of `dgamma`, `dnorm`, `dbeta`, `dbinom`, `dnbinom` or `dpareto1`, as appropriate. For the Gamma/Gamma case, use `shape.lik` for the shape parameter  $\tau$  of the Gamma likelihood. For the Normal/Normal case, use `sd.lik` for the standard error  $\sigma_2$  of the Normal likelihood.

Data for the linear Bayes case may be a matrix or data frame as usual; an atomic vector to fit the model to a single contract; missing or NULL to fit the prior model. Arguments `ratios`, `weights` and `subset` are ignored.

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>, Xavier Milhaud, Tommy Ouellet, Louis-Philippe Pouliot

### References

Bühlmann, H. and Gisler, A. (2005), *A Course in Credibility Theory and its Applications*, Springer.  
 Belhadj, H., Goulet, V. and Ouellet, T. (2009), On parameter estimation in hierarchical credibility, *Astin Bulletin* **39**.

Goulet, V. (1998), Principles and application of credibility theory, *Journal of Actuarial Practice* **6**, ISSN 1064-6647.

Goovaerts, M. J. and Hoogstad, W. J. (1987), *Credibility Theory*, Surveys of Actuarial Studies, No. 4, Nationale-Nederlanden N.V.

### See Also

[subset](#), [formula](#), [lm](#), [predict.lm](#).

### Examples

```
data(hachemeister)

## Buhlmann-Straub model
fit <- cm(~state, hachemeister,
         ratios = ratio.1:ratio.12, weights = weight.1:weight.12)
fit # print method
predict(fit) # credibility premiums
summary(fit) # more details

## Two-level hierarchical model. Notice that data does not have
## to be sorted by level
X <- data.frame(unit = c("A", "B", "A", "B", "B"), hachemeister)
fit <- cm(~unit + unit:state, X, ratio.1:ratio.12, weight.1:weight.12)
predict(fit)
predict(fit, levels = "unit") # unit credibility premiums only
summary(fit)
summary(fit, levels = "unit") # unit summaries only

## Regression model with intercept at time origin
fit <- cm(~state, hachemeister,
         regformula = ~time, regdata = data.frame(time = 12:1),
         ratios = ratio.1:ratio.12, weights = weight.1:weight.12)
fit
predict(fit, newdata = data.frame(time = 0))
summary(fit, newdata = data.frame(time = 0))

## Same regression model, with intercept at barycenter of time
fit <- cm(~state, hachemeister, adj.intercept = TRUE,
         regformula = ~time, regdata = data.frame(time = 12:1),
         ratios = ratio.1:ratio.12, weights = weight.1:weight.12)
fit
predict(fit, newdata = data.frame(time = 0))
summary(fit, newdata = data.frame(time = 0))

## Poisson/Gamma pure Bayesian model
fit <- cm("bayes", data = c(5, 3, 0, 1, 1),
         likelihood = "poisson", shape = 3, rate = 3)
fit
predict(fit)
summary(fit)
```

```
## Normal/Normal pure Bayesian model
cm("bayer", data = c(5, 3, 0, 1, 1),
    likelihood = "normal", sd.lik = 2,
    mean = 2, sd = 1)
```

---

 coverage

*Density and Cumulative Distribution Function for Modified Data*


---

### Description

Compute probability density function or cumulative distribution function of the payment per payment or payment per loss random variable under any combination of the following coverage modifications: deductible, limit, coinsurance, inflation.

### Usage

```
coverage(pdf, cdf, deductible = 0, franchise = FALSE,
          limit = Inf, coinsurance = 1, inflation = 0,
          per.loss = FALSE)
```

### Arguments

pdf, cdf	function object or character string naming a function to compute, respectively, the probability density function and cumulative distribution function of a probability law.
deductible	a unique positive numeric value.
franchise	logical; TRUE for a franchise deductible, FALSE (default) for an ordinary deductible.
limit	a unique positive numeric value larger than deductible.
coinsurance	a unique value between 0 and 1; the proportion of coinsurance.
inflation	a unique value between 0 and 1; the rate of inflation.
per.loss	logical; TRUE for the per loss distribution, FALSE (default) for the per payment distribution.

### Details

coverage returns a function to compute the probability density function (pdf) or the cumulative distribution function (cdf) of the distribution of losses under coverage modifications. The pdf and cdf of unmodified losses are pdf and cdf, respectively.

If pdf is specified, the pdf is returned; if pdf is missing or NULL, the cdf is returned. Note that cdf is needed if there is a deductible or a limit.

### Value

An object of mode "function" with the same arguments as pdf or cdf, except "lower.tail", "log.p" and "log", which are not supported.



**Note**

Setting arguments of the function returned by coverage using [formals](#) may very well not work as expected.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**See Also**

vignette("coverage") for the exact definitions of the per payment and per loss random variables under an ordinary or franchise deductible.

**Examples**

```
## Default case: pdf of the per payment random variable with
## an ordinary deductible
coverage(dgamma, pgamma, deductible = 1)

## Add a limit
f <- coverage(dgamma, pgamma, deductible = 1, limit = 7)
f <- coverage("dgamma", "pgamma", deductible = 1, limit = 7) # same
f(0, shape = 3, rate = 1)
f(2, shape = 3, rate = 1)
f(6, shape = 3, rate = 1)
f(8, shape = 3, rate = 1)
curve(dgamma(x, 3, 1), xlim = c(0, 10), ylim = c(0, 0.3)) # original
curve(f(x, 3, 1), xlim = c(0.01, 5.99), col = 4, add = TRUE) # modified
points(6, f(6, 3, 1), pch = 21, bg = 4)

## Cumulative distribution function
F <- coverage(cdf = pgamma, deductible = 1, limit = 7)
F(0, shape = 3, rate = 1)
F(2, shape = 3, rate = 1)
F(6, shape = 3, rate = 1)
F(8, shape = 3, rate = 1)
curve(pgamma(x, 3, 1), xlim = c(0, 10), ylim = c(0, 1)) # original
curve(F(x, 3, 1), xlim = c(0, 5.99), col = 4, add = TRUE) # modified
curve(F(x, 3, 1), xlim = c(6, 10), col = 4, add = TRUE) # modified

## With no deductible, all distributions below are identical
coverage(dweibull, pweibull, limit = 5)
coverage(dweibull, pweibull, per.loss = TRUE, limit = 5)
coverage(dweibull, pweibull, franchise = TRUE, limit = 5)
coverage(dweibull, pweibull, per.loss = TRUE, franchise = TRUE,
        limit = 5)
```

```
## Coinsurance alone; only case that does not require the cdf
coverage(dgamma, coinsurance = 0.8)
```

---

CTE *Conditional Tail Expectation*

---

### Description

Conditional Tail Expectation, also called Tail Value-at-Risk.  
TVaR is an alias for CTE.

### Usage

```
CTE(x, ...)

## S3 method for class 'aggregateDist'
CTE(x, conf.level = c(0.9, 0.95, 0.99),
     names = TRUE, ...)

TVaR(x, ...)
```

### Arguments

<code>x</code>	an R object.
<code>conf.level</code>	numeric vector of probabilities with values in $[0, 1)$ .
<code>names</code>	logical; if true, the result has a <code>names</code> attribute. Set to FALSE for speedup with many probs.
<code>...</code>	further arguments passed to or from other methods.

### Details

The Conditional Tail Expectation (or Tail Value-at-Risk) measures the average of losses above the Value at Risk for some given confidence level, that is  $E[X|X > \text{VaR}(X)]$  where  $X$  is the loss random variable.

CTE is a generic function with, currently, only a method for objects of class "aggregateDist".

For the recursive, convolution and simulation methods of [aggregateDist](#), the CTE is computed from the definition using the empirical cdf.

For the normal approximation method, an explicit formula exists:

$$\mu + \frac{\sigma}{(1 - \alpha)\sqrt{2\pi}} e^{-\text{VaR}(X)^2/2},$$

where  $\mu$  is the mean,  $\sigma$  the standard deviation and  $\alpha$  the confidence level.

For the Normal Power approximation, the explicit formula given in Castañer et al. (2013) is

$$\mu + \frac{\sigma}{(1 - \alpha)\sqrt{2\pi}} e^{-\text{VaR}(X)^2/2} \left(1 + \frac{\gamma}{6} \text{VaR}(X)\right),$$

where, as above,  $\mu$  is the mean,  $\sigma$  the standard deviation,  $\alpha$  the confidence level and  $\gamma$  is the skewness.

### Value

A numeric vector, named if names is TRUE.

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Tommy Ouellet

### References

Castañer, A. and Claramunt, M.M. and Mármol, M. (2013), Tail value at risk. An analysis with the Normal-Power approximation. In *Statistical and Soft Computing Approaches in Insurance Problems*, pp. 87-112. Nova Science Publishers, 2013. ISBN 978-1-62618-506-7.

### See Also

[aggregateDist](#); [VaR](#)

### Examples

```
model.freq <- expression(data = rpois(7))
model.sev <- expression(data = rnorm(9, 2))
Fs <- aggregateDist("simulation", model.freq, model.sev, nb.simul = 1000)
CTE(Fs)
```

---

dental

*Individual Dental Claims Data Set*

---

### Description

Basic dental claims on a policy with a deductible of 50.

### Usage

```
dental
```

### Format

A vector containing 10 observations

### Source

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (1998), *Loss Models, From Data to Decisions*, Wiley.

---

 discretize

*Discretization of a Continuous Distribution*


---

### Description

Compute a discrete probability mass function from a continuous cumulative distribution function (cdf) with various methods.

discretise is an alias for discretize.

### Usage

```
discretize(cdf, from, to, step = 1,
           method = c("upper", "lower", "rounding", "unbiased"),
           lev, by = step, xlim = NULL)
```

```
discretise(cdf, from, to, step = 1,
            method = c("upper", "lower", "rounding", "unbiased"),
            lev, by = step, xlim = NULL)
```

### Arguments

cdf	an expression written as a function of $x$ , or alternatively the name of a function, giving the cdf to discretize.
from, to	the range over which the function will be discretized.
step	numeric; the discretization step (or span, or lag).
method	discretization method to use.
lev	an expression written as a function of $x$ , or alternatively the name of a function, to compute the limited expected value of the distribution corresponding to cdf. Used only with the "unbiased" method.
by	an alias for step.
xlim	numeric of length 2; if specified, it serves as default for <code>c(from, to)</code> .

### Details

Usage is similar to [curve](#).

discretize returns the probability mass function (pmf) of the random variable obtained by discretization of the cdf specified in cdf.

Let  $F(x)$  denote the cdf,  $E[\min(X, x)]$  the limited expected value at  $x$ ,  $h$  the step,  $p_x$  the probability mass at  $x$  in the discretized distribution and set  $a = \text{from}$  and  $b = \text{to}$ .

Method "upper" is the forward difference of the cdf  $F$ :

$$p_x = F(x + h) - F(x)$$

for  $x = a, a + h, \dots, b - \text{step}$ .

Method "lower" is the backward difference of the cdf  $F$ :

$$p_x = F(x) - F(x - h)$$

for  $x = a + h, \dots, b$  and  $p_a = F(a)$ .

Method "rounding" has the true cdf pass through the midpoints of the intervals  $[x - h/2, x + h/2)$ :

$$p_x = F(x + h/2) - F(x - h/2)$$

for  $x = a + h, \dots, b - step$  and  $p_a = F(a + h/2)$ . The function assumes the cdf is continuous. Any adjustment necessary for discrete distributions can be done via cdf.

Method "unbiased" matches the first moment of the discretized and the true distributions. The probabilities are as follows:

$$p_a = \frac{E[\min(X, a)] - E[\min(X, a + h)]}{h} + 1 - F(a)$$

$$p_x = \frac{2E[\min(X, x)] - E[\min(X, x - h)] - E[\min(X, x + h)]}{h}, \quad a < x < b$$

$$p_b = \frac{E[\min(X, b)] - E[\min(X, b - h)]}{h} - 1 + F(b),$$

### Value

A numeric vector of probabilities suitable for use in [aggregateDist](#).

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>

### References

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

### See Also

[aggregateDist](#)

### Examples

```
x <- seq(0, 5, 0.5)

op <- par(mfrow = c(1, 1), col = "black")

## Upper and lower discretization
fu <- discretize(pgamma(x, 1), method = "upper",
                from = 0, to = 5, step = 0.5)
fl <- discretize(pgamma(x, 1), method = "lower",
                from = 0, to = 5, step = 0.5)
curve(pgamma(x, 1), xlim = c(0, 5))
```

```

par(col = "blue")
plot(stepfun(head(x, -1), diffinv(fu)), pch = 19, add = TRUE)
par(col = "green")
plot(stepfun(x, diffinv(fl)), pch = 19, add = TRUE)
par(col = "black")

## Rounding (or midpoint) discretization
fr <- discretize(pgamma(x, 1), method = "rounding",
                from = 0, to = 5, step = 0.5)
curve(pgamma(x, 1), xlim = c(0, 5))
par(col = "blue")
plot(stepfun(head(x, -1), diffinv(fr)), pch = 19, add = TRUE)
par(col = "black")

## First moment matching
fb <- discretize(pgamma(x, 1), method = "unbiased",
                lev = levgamma(x, 1), from = 0, to = 5, step = 0.5)
curve(pgamma(x, 1), xlim = c(0, 5))
par(col = "blue")
plot(stepfun(x, diffinv(fb)), pch = 19, add = TRUE)

par(op)

```

---

elev

*Empirical Limited Expected Value*


---

### Description

Compute the empirical limited expected value for individual or grouped data.

### Usage

```

elev(x, ...)

## Default S3 method:
elev(x, ...)

## S3 method for class 'grouped.data'
elev(x, ...)

## S3 method for class 'elev'
print(x, digits = getOption("digits") - 2, ...)

## S3 method for class 'elev'
summary(object, ...)

## S3 method for class 'elev'
knots(Fn, ...)

```

```
## S3 method for class 'elev'
plot(x, ..., main = NULL, xlab = "x", ylab = "Empirical LEV")
```

### Arguments

**x** a vector or an object of class "grouped.data" (in which case only the first column of frequencies is used); for the methods, an object of class "elev", typically.

**digits** number of significant digits to use, see [print](#).

**Fn, object** an R object inheriting from "ogive".

**main** main title.

**xlab, ylab** labels of x and y axis.

**...** arguments to be passed to subsequent methods.

### Details

The limited expected value (LEV) at  $u$  of a random variable  $X$  is  $E[X \wedge u] = E[\min(X, u)]$ . For individual data  $x_1, \dots, x_n$ , the empirical LEV  $E_n[X \wedge u]$  is thus

$$E_n[X \wedge u] = \frac{1}{n} \left( \sum_{x_j < u} x_j + \sum_{x_j \geq u} u \right).$$

Methods of `elev` exist for individual data or for grouped data created with `grouped.data`. The formula in this case is too long to show here. See the reference for details.

### Value

For `elev`, a function of class "elev", inheriting from the "function" class.

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

### References

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (1998), *Loss Models, From Data to Decisions*, Wiley.

### See Also

[grouped.data](#) to create grouped data objects; [stepfun](#) for related documentation (even though the empirical LEV is not a step function).

**Examples**

```

data(gdental)
lev <- elev(gdental)
lev
summary(lev)
knots(lev)          # the group boundaries

lev(knots(lev))     # empirical lev at boundaries
lev(c(80, 200, 2000)) # and at other limits

plot(lev, type = "o", pch = 16)

```

emm

*Empirical Moments***Description**

Raw empirical moments for individual and grouped data.

**Usage**

```

emm(x, order = 1, ...)

## Default S3 method:
emm(x, order = 1, ...)

## S3 method for class 'grouped.data'
emm(x, order = 1, ...)

```

**Arguments**

`x` a vector or matrix of individual data, or an object of class "grouped data".

`order` order of the moment. Must be positive.

`...` further arguments passed to or from other methods.

**Details**

Arguments `...` are passed to `colMeans`; `na.rm = TRUE` may be useful for individual data with missing values.

For individual data, the  $k$ th empirical moment is  $\sum_{j=1}^n x_j^k$ .

For grouped data with group boundaries  $c_1, \dots, c_r$  and group frequencies  $n_1, \dots, n_r$ , the  $k$ th empirical moment is

$$\sum_{j=1}^r \frac{n_j(c_j^{k+1} - c_{j-1}^{k+1})}{n(k+1)(c_j - c_{j-1})},$$

where  $n = \sum_{j=1}^r n_j$ .



**Value**

A named vector or matrix of moments.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (1998), *Loss Models, From Data to Decisions*, Wiley.

**See Also**

[mean](#) and [mean.grouped.data](#) for simpler access to the first moment.

**Examples**

```
## Individual data
data(dental)
emm(dental, order = 1:3)

## Grouped data
data(gdental)
emm(gdental)
x <- grouped.data(cj = gdental[, 1],
                  nj1 = sample(1:100, nrow(gdental)),
                  nj2 = sample(1:100, nrow(gdental)))
emm(x) # same as mean(x)
```

---

ExponentialSupp

*Moments and Moment Generating Function of the Exponential Distribution*

---

**Description**

Raw moments, limited moments and moment generating function for the exponential distribution with rate rate (i.e., mean 1/rate).

**Usage**

```
mexp(order, rate = 1)
levexp(limit, rate = 1, order = 1)
mgfexp(t, rate = 1, log = FALSE)
```

**Arguments**

order	order of the moment.
limit	limit of the loss variable.
rate	vector of rates.
t	numeric vector.
log	logical; if TRUE, the cumulant generating function is returned.

**Details**

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ , the  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$  and the moment generating function is  $E[e^{tX}]$ ,  $k > -1$ .

**Value**

mexp gives the  $k$ th raw moment, levexp gives the  $k$ th moment of the limited loss variable, and mgfexp gives the moment generating function in  $t$ .

Invalid arguments will result in return value NaN, with a warning.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>, Christophe Dutang and Mathieu Pigeon.

**References**

Johnson, N. L. and Kotz, S. (1970), *Continuous Univariate Distributions, Volume 1*, Wiley.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**See Also**

[Exponential](#)

**Examples**

```
mexp(2, 3) - mexp(1, 3)^2
levexp(10, 3, order = 2)
mgfexp(1,2)
```

---

 Extract.grouped.data *Extract or Replace Parts of a Grouped Data Object*


---

## Description

Extract or replace subsets of grouped data objects.

## Usage

```
## S3 method for class 'grouped.data'
x[i, j]
## S3 replacement method for class 'grouped.data'
x[i, j] <- value
```

## Arguments

x	an object of class grouped.data.
i, j	elements to extract or replace. i, j are numeric or character or, for [ only, empty. Numeric values are coerced to integer as if by <a href="#">as.integer</a> . For replacement by [, a logical matrix is allowed, but not replacement in the group boundaries and group frequencies simultaneously.
value	a suitable replacement value.

## Details

Objects of class "grouped.data" can mostly be indexed like data frames, with the following restrictions:

1. For [, the extracted object must keep a group boundaries column and at least one group frequencies column to remain of class "grouped.data";
2. For [<-, it is not possible to replace group boundaries and group frequencies simultaneously;
3. When replacing group boundaries, `length(value) == length(i) + 1`.

`x[, 1]` will return the plain vector of group boundaries.

Replacement of non adjacent group boundaries is not possible for obvious reasons.

Otherwise, extraction and replacement should work just like for data frames.

## Value

For [ an object of class "grouped.data", a data frame or a vector.

For [<- an object of class "grouped.data".

## Note

Currently [[, [[<-, \$ and \$<- are not specifically supported, but should work as usual on group frequency columns.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>

**See Also**

[\[.data.frame\]](#) for extraction and replacement methods of data frames, [grouped.data](#) to create grouped data objects.

**Examples**

```
data(gdental)

(x <- gdental[1])      # select column 1
class(x)               # no longer a grouped.data object
class(gdental[2])     # same
gdental[, 1]           # group boundaries
gdental[, 2]           # group frequencies

gdental[1:4,]         # a subset
gdental[c(1, 3, 5),]  # avoid this

gdental[1:2, 1] <- c(0, 30, 60) # modified boundaries
gdental[, 2] <- 10           # modified frequencies
## Not run: gdental[1, ] <- 2 # not allowed
```

---

FellerPareto

*The Feller Pareto Distribution*

---

**Description**

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Feller Pareto distribution with parameters `min`, `shape1`, `shape2`, `shape3` and `scale`.

**Usage**

```
dfpareto(x, min, shape1, shape2, shape3, rate = 1, scale = 1/rate,
         log = FALSE)
pfpareto(q, min, shape1, shape2, shape3, rate = 1, scale = 1/rate,
         lower.tail = TRUE, log.p = FALSE)
qfpareto(p, min, shape1, shape2, shape3, rate = 1, scale = 1/rate,
         lower.tail = TRUE, log.p = FALSE)
rfpareto(n, min, shape1, shape2, shape3, rate = 1, scale = 1/rate)
mfpareto(order, min, shape1, shape2, shape3, rate = 1, scale = 1/rate)
levfpareto(limit, min, shape1, shape2, shape3, rate = 1, scale = 1/rate,
           order = 1)
```

**Arguments**

<code>x, q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>min</code>	lower bound of the support of the distribution.
<code>shape1, shape2, shape3, scale</code>	parameters. Must be strictly positive.
<code>rate</code>	an alternative way to specify the scale.
<code>log, log.p</code>	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
<code>order</code>	order of the moment.
<code>limit</code>	limit of the loss variable.

**Details**

The Feller-Pareto distribution with parameters  $\text{min} = \mu$ ,  $\text{shape1} = \alpha$ ,  $\text{shape2} = \gamma$ ,  $\text{shape3} = \tau$  and  $\text{scale} = \theta$ , has density:

$$f(x) = \frac{\Gamma(\alpha + \tau)}{\Gamma(\alpha)\Gamma(\tau)} \frac{\gamma((x - \mu)/\theta)^{\gamma\tau - 1}}{\theta[1 + ((x - \mu)/\theta)^\gamma]^{\alpha + \tau}}$$

for  $x > \mu$ ,  $-\infty < \mu < \infty$ ,  $\alpha > 0$ ,  $\gamma > 0$ ,  $\tau > 0$  and  $\theta > 0$ . (Here  $\Gamma(\alpha)$  is the function implemented by R's `gamma()` and defined in its help.)

The Feller-Pareto is the distribution of the random variable

$$\mu + \theta \left( \frac{1 - X}{X} \right)^{1/\gamma},$$

where  $X$  has a beta distribution with parameters  $\alpha$  and  $\tau$ .

The Feller-Pareto defines a large family of distributions encompassing the transformed beta family and many variants of the Pareto distribution. Setting  $\mu = 0$  yields the [transformed beta](#) distribution.

The Feller-Pareto distribution also has the following direct special cases:

- A [Pareto IV](#) distribution when `shape3 == 1`;
- A [Pareto III](#) distribution when `shape1 shape3 == 1`;
- A [Pareto II](#) distribution when `shape1 shape2 == 1`;
- A [Pareto I](#) distribution when `shape1 shape2 == 1` and `min = scale`.

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$  for nonnegative integer values of  $k < \alpha\gamma$ .

The  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$  for nonnegative integer values of  $k$  and  $\alpha - j/\gamma, j = 1, \dots, k$  not a negative integer.

Note that the range of admissible values for  $k$  in raw and limited moments is larger when  $\mu = 0$ .

**Value**

dfpareto gives the density, pfpareto gives the distribution function, qfpareto gives the quantile function, rfpareto generates random deviates, mfpareto gives the  $k$ th raw moment, and levfpareto gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

**Note**

levfpareto computes the limited expected value using [betaint](#).

For the Feller-Pareto and other Pareto distributions, we use the classification of Arnold (2015) with the parametrization of Klugman et al. (2012).

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Nicholas Langevin

**References**

Arnold, B.C. (2015), *Pareto Distributions*, Second Edition, CRC Press.

Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

Abramowitz, M. and Stegun, I. A. (1972), *Handbook of Mathematical Functions*, Dover.

**See Also**

[dtrbeta](#) for the transformed beta distribution.

**Examples**

```
exp(dfpareto(2, 1, 2, 3, 4, 5, log = TRUE))
p <- (1:10)/10
pfpareto(qfpareto(p, 1, 2, 3, 4, 5), 1, 2, 3, 4, 5)

## variance
mfpareto(2, 1, 2, 3, 4, 5) - mfpareto(1, 1, 2, 3, 4, 5)^2

## case with shape1 - order/shape2 > 0
levfpareto(10, 1, 2, 3, 4, scale = 1, order = 2)

## case with shape1 - order/shape2 < 0
levfpareto(20, 10, 0.1, 14, 2, scale = 1.5, order = 2)
```

---

GammaSupp	<i>Moments and Moment Generating Function of the Gamma Distribution</i>
-----------	---

---

### Description

Raw moments, limited moments and moment generating function for the Gamma distribution with parameters shape and scale.

### Usage

```
mgamma(order, shape, rate = 1, scale = 1/rate)
levgamma(limit, shape, rate = 1, scale = 1/rate, order = 1)
mgfgamma(t, shape, rate = 1, scale = 1/rate, log = FALSE)
```

### Arguments

order	order of the moment.
limit	limit of the loss variable.
rate	an alternative way to specify the scale.
shape, scale	shape and scale parameters. Must be strictly positive.
t	numeric vector.
log	logical; if TRUE, the cumulant generating function is returned.

### Details

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ , the  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$  and the moment generating function is  $E[e^{tX}]$ ,  $k > -\alpha$ .

### Value

mgamma gives the  $k$ th raw moment, levgamma gives the  $k$ th moment of the limited loss variable, and mgfgamma gives the moment generating function in t.

Invalid arguments will result in return value NaN, with a warning.

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>, Christophe Dutang and Mathieu Pigeon

### References

Johnson, N. L. and Kotz, S. (1970), *Continuous Univariate Distributions, Volume 1*, Wiley.  
 Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**See Also**[GammaDist](#)**Examples**

```
mgamma(2, 3, 4) - mgamma(1, 3, 4)^2
levgamma(10, 3, 4, order = 2)
mgfgamma(1,3,2)
```

---

gdental	<i>Grouped Dental Claims Data Set</i>
---------	---------------------------------------

---

**Description**

Grouped dental claims, that is presented in a number of claims per claim amount group form.

**Usage**

```
gdental
```

**Format**

An object of class "grouped.data" (inheriting from class "data.frame") consisting of 10 rows and 2 columns. The environment of the object contains the plain vector of `cj` of group boundaries

**Source**

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (1998), *Loss Models, From Data to Decisions*, Wiley.

**See Also**

[grouped.data](#) for a description of grouped data objects.

---

GeneralizedBeta	<i>The Generalized Beta Distribution</i>
-----------------	--

---

**Description**

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Generalized Beta distribution with parameters `shape1`, `shape2`, `shape3` and `scale`.



**Usage**

```

dgenbeta(x, shape1, shape2, shape3, rate = 1, scale = 1/rate,
         log = FALSE)
pgenbeta(q, shape1, shape2, shape3, rate = 1, scale = 1/rate,
         lower.tail = TRUE, log.p = FALSE)
qgenbeta(p, shape1, shape2, shape3, rate = 1, scale = 1/rate,
         lower.tail = TRUE, log.p = FALSE)
rgenbeta(n, shape1, shape2, shape3, rate = 1, scale = 1/rate)
mgenbeta(order, shape1, shape2, shape3, rate = 1, scale = 1/rate)
levgenbeta(limit, shape1, shape2, shape3, rate = 1, scale = 1/rate,
           order = 1)

```

**Arguments**

<code>x, q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>shape1, shape2, shape3, scale</code>	parameters. Must be strictly positive.
<code>rate</code>	an alternative way to specify the scale.
<code>log, log.p</code>	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
<code>order</code>	order of the moment.
<code>limit</code>	limit of the loss variable.

**Details**

The generalized beta distribution with parameters  $\text{shape1} = \alpha$ ,  $\text{shape2} = \beta$ ,  $\text{shape3} = \tau$  and  $\text{scale} = \theta$ , has density:

$$f(x) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} (x/\theta)^{\alpha\tau} (1 - (x/\theta)^\tau)^{\beta-1} \frac{\tau}{x}$$

for  $0 < x < \theta$ ,  $\alpha > 0$ ,  $\beta > 0$ ,  $\tau > 0$  and  $\theta > 0$ . (Here  $\Gamma(\alpha)$  is the function implemented by R's [gamma\(\)](#) and defined in its help.)

The generalized beta is the distribution of the random variable

$$\theta X^{1/\tau},$$

where  $X$  has a beta distribution with parameters  $\alpha$  and  $\beta$ .

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$  and the  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)]$ ,  $k > -\alpha\tau$ .

**Value**

dgenbeta gives the density, pgenbeta gives the distribution function, qgenbeta gives the quantile function, rgenbeta generates random deviates, mgenbeta gives the  $k$ th raw moment, and levgenbeta gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

**Note**

This is *not* the generalized three-parameter beta distribution defined on page 251 of Johnson et al, 1995.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>

**References**

Johnson, N. L., Kotz, S. and Balakrishnan, N. (1995) *Continuous Univariate Distributions, Volume 2*, Wiley.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**Examples**

```
exp(dgenbeta(2, 2, 3, 4, 0.2, log = TRUE))
p <- (1:10)/10
pgenbeta(qgenbeta(p, 2, 3, 4, 0.2), 2, 3, 4, 0.2)
mgenbeta(2, 1, 2, 3, 0.25) - mgenbeta(1, 1, 2, 3, 0.25) ^ 2
levgenbeta(10, 1, 2, 3, 0.25, order = 2)
```

**Description**

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Generalized Pareto distribution with parameters shape1, shape2 and scale.

**Usage**

```
dgenpareto(x, shape1, shape2, rate = 1, scale = 1/rate,
           log = FALSE)
pgenpareto(q, shape1, shape2, rate = 1, scale = 1/rate,
           lower.tail = TRUE, log.p = FALSE)
qgenpareto(p, shape1, shape2, rate = 1, scale = 1/rate,
           lower.tail = TRUE, log.p = FALSE)
rgenpareto(n, shape1, shape2, rate = 1, scale = 1/rate)
mgenpareto(order, shape1, shape2, rate = 1, scale = 1/rate)
levgenpareto(limit, shape1, shape2, rate = 1, scale = 1/rate,
             order = 1)
```

**Arguments**

<code>x, q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>shape1, shape2, scale</code>	parameters. Must be strictly positive.
<code>rate</code>	an alternative way to specify the scale.
<code>log, log.p</code>	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
<code>order</code>	order of the moment.
<code>limit</code>	limit of the loss variable.

**Details**

The Generalized Pareto distribution with parameters  $\text{shape1} = \alpha$ ,  $\text{shape2} = \tau$  and  $\text{scale} = \theta$  has density:

$$f(x) = \frac{\Gamma(\alpha + \tau)}{\Gamma(\alpha)\Gamma(\tau)} \frac{\theta^\alpha x^{\tau-1}}{(x + \theta)^{\alpha+\tau}}$$

for  $x > 0$ ,  $\alpha > 0$ ,  $\tau > 0$  and  $\theta > 0$ . (Here  $\Gamma(\alpha)$  is the function implemented by R's [gamma\(\)](#) and defined in its help.)

The Generalized Pareto is the distribution of the random variable

$$\theta \left( \frac{X}{1 - X} \right),$$

where  $X$  has a beta distribution with parameters  $\alpha$  and  $\tau$ .

The Generalized Pareto distribution has the following special cases:

- A [Pareto](#) distribution when `shape2 == 1`;
- An [Inverse Pareto](#) distribution when `shape1 == 1`.

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $-\tau < k < \alpha$ .

The  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ ,  $k > -\tau$  and  $\alpha - k$  not a negative integer.

**Value**

dgenpareto gives the density, pgenpareto gives the distribution function, qgenpareto gives the quantile function, rgenpareto generates random deviates, mgenpareto gives the  $k$ th raw moment, and levgenpareto gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

**Note**

levgenpareto computes the limited expected value using [betaint](#).

Distribution also known as the Beta of the Second Kind. See also Kleiber and Kotz (2003) for alternative names and parametrizations.

The Generalized Pareto distribution defined here is different from the one in Embrechts et al. (1997) and in [Wikipedia](#); see also Kleiber and Kotz (2003, section 3.12). One may most likely compute quantities for the latter using functions for the [Pareto](#) distribution with the appropriate change of parametrization.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Embrechts, P., Klüppelberg, C. and Mikisch, T. (1997), *Modelling Extremal Events for Insurance and Finance*, Springer.

Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**Examples**

```
exp(dgenpareto(3, 3, 4, 4, log = TRUE))
p <- (1:10)/10
pgenpareto(qgenpareto(p, 3, 3, 1), 3, 3, 1)
qgenpareto(.3, 3, 4, 4, lower.tail = FALSE)

## variance
mgenpareto(2, 3, 2, 1) - mgenpareto(1, 3, 2, 1)^2

## case with shape1 - order > 0
levgenpareto(10, 3, 3, scale = 1, order = 2)

## case with shape1 - order < 0
levgenpareto(10, 1.5, 3, scale = 1, order = 2)
```

---

grouped.data	<i>Grouped data</i>
--------------	---------------------

---

## Description

Creation of grouped data objects, from either a provided set of group boundaries and group frequencies, or from individual data using automatic or specified breakpoints.

## Usage

```
grouped.data(..., breaks = "Sturges", include.lowest = TRUE,
             right = TRUE, nclass = NULL, group = FALSE,
             row.names = NULL, check.rows = FALSE,
             check.names = TRUE)
```

## Arguments

...	these arguments are either of the form value or tag = value. See Details.
breaks	same as for <a href="#">hist</a> , namely one of: <ul style="list-style-type: none"> <li>• a vector giving the breakpoints between groups;</li> <li>• a function to compute the vector of breakpoints;</li> <li>• a single number giving the number of groups;</li> <li>• a character string naming an algorithm to compute the number of groups (see <a href="#">hist</a>);</li> <li>• a function to compute the number of groups.</li> </ul> <p>In the last three cases the number is a suggestion only; the breakpoints will be set to <a href="#">pretty</a> values. If breaks is a function, the first element in ... is supplied to it as the only argument.</p>
include.lowest	logical; if TRUE, a data point equal to the breaks value will be included in the first (or last, for right = FALSE) group. Used only for individual data; see Details.
right	logical; indicating if the intervals should be closed on the right (and open on the left) or vice versa.
nclass	numeric (integer); equivalent to breaks for a scalar or character argument.
group	logical; an alternative way to force grouping of individual data.
row.names, check.rows, check.names	arguments identical to those of <a href="#">data.frame</a> .

## Details

A grouped data object is a special form of data frame consisting of one column of contiguous group boundaries and one or more columns of frequencies within each group.

The function can create a grouped data object from two types of arguments.

1. Group boundaries and frequencies. This is the default mode of operation if the call has at least two elements in . . . .

The first argument will then be taken as the vector of group boundaries. This vector must be exactly one element longer than the other arguments, which will be taken as vectors of group frequencies. All arguments are coerced to data frames.

2. Individual data. This mode of operation is active if there is a single argument in . . . , or if either breaks or nclass is specified or group is TRUE.

Arguments of . . . are first grouped using `hist`. If needed, breakpoints are set using the first argument.

Missing (NA) frequencies are replaced by zeros, with a warning.

Extraction and replacement methods exist for `grouped.data` objects, but working on non adjacent groups will most likely yield useless results.

### Value

An object of class `c("grouped.data", "data.frame")` with an environment containing the vector `cj` of group boundaries.

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>, Mathieu Pigeon and Louis-Philippe Pouliot

### References

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (1998), *Loss Models, From Data to Decisions*, Wiley.

### See Also

[\[.grouped.data\]](#) for extraction and replacement methods.

[data.frame](#) for usual data frame creation and manipulation.

[hist](#) for details on the calculation of breakpoints.

### Examples

```
## Most common usage using a predetermined set of group
## boundaries and group frequencies.
cj <- c(0, 25, 50, 100, 250, 500, 1000)
nj <- c(30, 31, 57, 42, 45, 10)
(x <- grouped.data(Group = cj, Frequency = nj))
class(x)

x[, 1] # group boundaries
x[, 2] # group frequencies

## Multiple frequency columns are supported
x <- sample(1:100, 9)
y <- sample(1:100, 9)
```

```

grouped.data(cj = 1:10, nj.1 = x, nj.2 = y)

## Alternative usage with grouping of individual data.
grouped.data(x)                # automatic breakpoints
grouped.data(x, breaks = 7)    # forced number of groups
grouped.data(x, breaks = c(0,25,75,100)) # specified groups
grouped.data(x, y, breaks = c(0,25,75,100)) # multiple data sets

## Not run: ## Providing two or more data sets and automatic breakpoints is
## very error-prone since the range of the first data set has to
## include the ranges of all the other data sets.
range(x)
range(y)
grouped.data(x, y, group = TRUE)
## End(Not run)

```

---

Gumbel

*The Gumbel Distribution*


---

### Description

Density function, distribution function, quantile function, random generation and raw moments for the Gumbel extreme value distribution with parameters alpha and scale.

### Usage

```

dgumbel(x, alpha, scale, log = FALSE)
pgumbel(q, alpha, scale, lower.tail = TRUE, log.p = FALSE)
qgumbel(p, alpha, scale, lower.tail = TRUE, log.p = FALSE)
rgumbel(n, alpha, scale)
mgumbel(order, alpha, scale)
mgfgumbel(t, alpha, scale, log = FALSE)

```

### Arguments

x, q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If $\text{length}(n) > 1$ , the length is taken to be the number required.
alpha	location parameter.
scale	parameter. Must be strictly positive.
log, log.p	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
order	order of the moment. Only values 1 and 2 are supported.
t	numeric vector.

**Details**

The Gumbel distribution with parameters  $\alpha = \alpha$  and  $\text{scale} = \theta$  has distribution function:

$$F(x) = \exp[-\exp(-(x - \alpha)/\theta)]$$

for  $-\infty < x < \infty$ ,  $-\infty < \alpha < \infty$  and  $\theta > 0$ .

The mode of the distribution is in  $\alpha$ , the mean is  $\alpha + \gamma\theta$ , where  $\gamma = 0.57721566$  is the Euler-Mascheroni constant, and the variance is  $\pi^2\theta^2/6$ .

**Value**

`dgumbel` gives the density, `pgumbel` gives the distribution function, `qgumbel` gives the quantile function, `rgumbel` generates random deviates, `mgumbel` gives the  $k$ th raw moment,  $k = 1, 2$ , and `mgfgamma` gives the moment generating function in `t`.

Invalid arguments will result in return value `NaN`, with a warning.

**Note**

Distribution also known as the generalized extreme value distribution Type-I.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>

**References**

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**Examples**

```
dgumbel(c(-5, 0, 10, 20), 0.5, 2)

p <- (1:10)/10
pgumbel(qgumbel(p, 2, 3), 2, 3)

curve(pgumbel(x, 0.5, 2), from = -5, to = 20, col = "red")
curve(pgumbel(x, 1.0, 2), add = TRUE, col = "green")
curve(pgumbel(x, 1.5, 3), add = TRUE, col = "blue")
curve(pgumbel(x, 3.0, 4), add = TRUE, col = "cyan")

a <- 3; s <- 4
mgumbel(1, a, s)           # mean
a - s * digamma(1)       # same

mgumbel(2, a, s) - mgumbel(1, a, s)^2 # variance
(pi * s)^2/6             # same
```



---

hachemeister	<i>Hachemeister Data Set</i>
--------------	------------------------------

---

**Description**

Hachemeister (1975) data set giving average claim amounts in private passenger bodily injury insurance in five U.S. states over 12 quarters between July 1970 and June 1973 and the corresponding number of claims.

**Usage**

```
hachemeister
```

**Format**

A matrix with 5 rows and the following 25 columns:

state the state number;

ratio.1,...,ratio.12 the average claim amounts;

weight.1,...,weight.12 the corresponding number of claims.

**Source**

Hachemeister, C. A. (1975), *Credibility for regression models with application to trend*, Proceedings of the Berkeley Actuarial Research Conference on Credibility, Academic Press.

---

hist.grouped.data	<i>Histogram for Grouped Data</i>
-------------------	-----------------------------------

---

**Description**

This method for the generic function `hist` is mainly useful to plot the histogram of grouped data. If `plot = FALSE`, the resulting object of class "histogram" is returned for compatibility with `hist.default`, but does not contain much information not already in `x`.

**Usage**

```
## S3 method for class 'grouped.data'
hist(x, freq = NULL, probability = !freq,
     density = NULL, angle = 45, col = NULL, border = NULL,
     main = paste("Histogram of" , xname),
     xlim = range(cj), ylim = NULL, xlab = xname, ylab,
     axes = TRUE, plot = TRUE, labels = FALSE, ...)
```

**Arguments**

x	an object of class "grouped.data"; only the first column of frequencies is used.
freq	logical; if TRUE, the histogram graphic is a representation of frequencies, the counts component of the result; if FALSE, probability densities, component density, are plotted (so that the histogram has a total area of one). Defaults to TRUE <i>iff</i> group boundaries are equidistant (and probability is not specified).
probability	an <i>alias</i> for !freq, for S compatibility.
density	the density of shading lines, in lines per inch. The default value of NULL means that no shading lines are drawn. Non-positive values of density also inhibit the drawing of shading lines.
angle	the slope of shading lines, given as an angle in degrees (counter-clockwise).
col	a colour to be used to fill the bars. The default of NULL yields unfilled bars.
border	the color of the border around the bars. The default is to use the standard foreground color.
main, xlab, ylab	these arguments to title have useful defaults here.
xlim, ylim	the range of x and y values with sensible defaults. Note that xlim is <i>not</i> used to define the histogram (breaks), but only for plotting (when plot = TRUE).
axes	logical. If TRUE (default), axes are draw if the plot is drawn.
plot	logical. If TRUE (default), a histogram is plotted, otherwise a list of breaks and counts is returned.
labels	logical or character. Additionally draw labels on top of bars, if not FALSE; see <a href="#">plot.histogram</a> .
...	further graphical parameters passed to <a href="#">plot.histogram</a> and their to <a href="#">title</a> and <a href="#">axis</a> (if plot=TRUE).

**Value**

An object of class "histogram" which is a list with components:

breaks	the $r + 1$ group boundaries.
counts	$r$ integers; the frequency within each group.
density	the relative frequencies within each group $n_j/n$ , where $n_j = \text{counts}[j]$ .
intensities	same as density. Deprecated, but retained for compatibility.
mids	the $r$ group midpoints.
xname	a character string with the actual x argument name.
equidist	logical, indicating if the distances between breaks are all the same.

**Note**

The resulting value does *not* depend on the values of the arguments freq (or probability) or plot. This is intentionally different from S.

**References**

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (1998), *Loss Models, From Data to Decisions*, Wiley.

**See Also**

[hist](#) and [hist.default](#) for histograms of individual data and fancy examples.

**Examples**

```
data(gdental)
hist(gdental)
```

---

 InverseBurr

*The Inverse Burr Distribution*


---

**Description**

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Inverse Burr distribution with parameters shape1, shape2 and scale.

**Usage**

```
dinvburr(x, shape1, shape2, rate = 1, scale = 1/rate,
          log = FALSE)
pinvburr(q, shape1, shape2, rate = 1, scale = 1/rate,
          lower.tail = TRUE, log.p = FALSE)
qinvburr(p, shape1, shape2, rate = 1, scale = 1/rate,
          lower.tail = TRUE, log.p = FALSE)
rinvburr(n, shape1, shape2, rate = 1, scale = 1/rate)
minvburr(order, shape1, shape2, rate = 1, scale = 1/rate)
levinvburr(limit, shape1, shape2, rate = 1, scale = 1/rate,
            order = 1)
```

**Arguments**

x, q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If length(n) > 1, the length is taken to be the number required.
shape1, shape2, scale	parameters. Must be strictly positive.
rate	an alternative way to specify the scale.
log, log.p	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
order	order of the moment.
limit	limit of the loss variable.

### Details

The inverse Burr distribution with parameters  $\text{shape1} = \tau$ ,  $\text{shape2} = \gamma$  and  $\text{scale} = \theta$ , has density:

$$f(x) = \frac{\tau\gamma(x/\theta)^{\gamma\tau}}{x[1 + (x/\theta)^\gamma]^{\tau+1}}$$

for  $x > 0$ ,  $\tau > 0$ ,  $\gamma > 0$  and  $\theta > 0$ .

The inverse Burr is the distribution of the random variable

$$\theta \left( \frac{X}{1-X} \right)^{1/\gamma},$$

where  $X$  has a beta distribution with parameters  $\tau$  and 1.

The inverse Burr distribution has the following special cases:

- A [Loglogistic](#) distribution when  $\text{shape1} == 1$ ;
- An [Inverse Pareto](#) distribution when  $\text{shape2} == 1$ ;
- An [Inverse Paralogistic](#) distribution when  $\text{shape1} == \text{shape2}$ .

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $-\tau\gamma < k < \gamma$ .

The  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ ,  $k > -\tau\gamma$  and  $1 - k/\gamma$  not a negative integer.

### Value

`dinvburr` gives the density, `invburr` gives the distribution function, `qinvburr` gives the quantile function, `rinvburr` generates random deviates, `minvburr` gives the  $k$ th raw moment, and `levinvburr` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

### Note

`levinvburr` computes the limited expected value using [betaint](#).

Also known as the Dagum distribution. See also Kleiber and Kotz (2003) for alternative names and parametrizations.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

### References

- Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.
- Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**Examples**

```

exp(dinvburr(2, 2, 3, 1, log = TRUE))
p <- (1:10)/10
pinvburr(qinvburr(p, 2, 3, 1), 2, 3, 1)

## variance
minvburr(2, 2, 3, 1) - minvburr(1, 2, 3, 1) ^ 2

## case with 1 - order/shape2 > 0
levinvburr(10, 2, 3, 1, order = 2)

## case with 1 - order/shape2 < 0
levinvburr(10, 2, 1.5, 1, order = 2)

```

---

InverseExponential      *The Inverse Exponential Distribution*

---

**Description**

Density function, distribution function, quantile function, random generation raw moments and limited moments for the Inverse Exponential distribution with parameter scale.

**Usage**

```

dinvexp(x, rate = 1, scale = 1/rate, log = FALSE)
pinvexp(q, rate = 1, scale = 1/rate, lower.tail = TRUE, log.p = FALSE)
qinvexp(p, rate = 1, scale = 1/rate, lower.tail = TRUE, log.p = FALSE)
rinvexp(n, rate = 1, scale = 1/rate)
minvexp(order, rate = 1, scale = 1/rate)
levinvexp(limit, rate = 1, scale = 1/rate, order)

```

**Arguments**

<code>x, q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>scale</code>	parameter. Must be strictly positive.
<code>rate</code>	an alternative way to specify the scale.
<code>log, log.p</code>	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
<code>order</code>	order of the moment.
<code>limit</code>	limit of the loss variable.

**Details**

The inverse exponential distribution with parameter  $\text{scale} = \theta$  has density:

$$f(x) = \frac{\theta e^{-\theta/x}}{x^2}$$

for  $x > 0$  and  $\theta > 0$ .

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $k < 1$ , and the  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ , all  $k$ .

**Value**

`dinvexp` gives the density, `pinvexp` gives the distribution function, `qinvexp` gives the quantile function, `rinvexp` generates random deviates, `minvexp` gives the  $k$ th raw moment, and `levinvexp` calculates the  $k$ th limited moment.

Invalid arguments will result in return value NaN, with a warning.

**Note**

`levinvexp` computes the limited expected value using `gammainc` from package **expint**.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**Examples**

```
exp(dinvexp(2, 2, log = TRUE))
p <- (1:10)/10
pinvexp(qinvexp(p, 2), 2)
minvexp(0.5, 2)
```

**Description**

Density function, distribution function, quantile function, random generation, raw moments, and limited moments for the Inverse Gamma distribution with parameters shape and scale.

**Usage**

```
dinvgamma(x, shape, rate = 1, scale = 1/rate, log = FALSE)
pinvgamma(q, shape, rate = 1, scale = 1/rate,
          lower.tail = TRUE, log.p = FALSE)
qinvgamma(p, shape, rate = 1, scale = 1/rate,
          lower.tail = TRUE, log.p = FALSE)
rinvgamma(n, shape, rate = 1, scale = 1/rate)
minvgamma(order, shape, rate = 1, scale = 1/rate)
levinvgamma(limit, shape, rate = 1, scale = 1/rate,
            order = 1)
mgfinvgamma(t, shape, rate = 1, scale = 1/rate, log = FALSE)
```

**Arguments**

x, q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
shape, scale	parameters. Must be strictly positive.
rate	an alternative way to specify the scale.
log, log.p	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
order	order of the moment.
limit	limit of the loss variable.
t	numeric vector.

**Details**

The inverse gamma distribution with parameters  $\text{shape} = \alpha$  and  $\text{scale} = \theta$  has density:

$$f(x) = \frac{u^\alpha e^{-u}}{x\Gamma(\alpha)}, \quad u = \theta/x$$

for  $x > 0$ ,  $\alpha > 0$  and  $\theta > 0$ . (Here  $\Gamma(\alpha)$  is the function implemented by R's `gamma()` and defined in its help.)

The special case  $\text{shape} == 1$  is an [Inverse Exponential](#) distribution.

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $k < \alpha$ , and the  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ , all  $k$ .

The moment generating function is given by  $E[e^{tX}]$ .

**Value**

`dinvgamma` gives the density, `pinvgamma` gives the distribution function, `qinvgamma` gives the quantile function, `rinvgamma` generates random deviates, `minvgamma` gives the  $k$ th raw moment, `levinvgamma` gives the  $k$ th moment of the limited loss variable, and `mgfinvgamma` gives the moment generating function in `t`.

Invalid arguments will result in return value `NaN`, with a warning.

**Note**

levinvgamma computes the limited expected value using `gammainc` from package **expint**.

Also known as the Vinci distribution. See also Kleiber and Kotz (2003) for alternative names and parametrizations.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**Examples**

```
exp(dinvgamma(2, 3, 4, log = TRUE))
p <- (1:10)/10
pinvgamma(qinvgamma(p, 2, 3), 2, 3)
minvgamma(-1, 2, 2) ^ 2
levinvgamma(10, 2, 2, order = 1)
mgfinvgamma(-1, 3, 2)
```

---

InverseGaussian

*The Inverse Gaussian Distribution*

---

**Description**

Density function, distribution function, quantile function, random generation, raw moments, limited moments and moment generating function for the Inverse Gaussian distribution with parameters mean and shape.

**Usage**

```
dinvgauss(x, mean, shape = 1, dispersion = 1/shape,
          log = FALSE)
pinvgauss(q, mean, shape = 1, dispersion = 1/shape,
          lower.tail = TRUE, log.p = FALSE)
qinvgauss(p, mean, shape = 1, dispersion = 1/shape,
          lower.tail = TRUE, log.p = FALSE,
          tol = 1e-14, maxit = 100, echo = FALSE, trace = echo)
rinvgauss(n, mean, shape = 1, dispersion = 1/shape)
minvgauss(order, mean, shape = 1, dispersion = 1/shape)
```



```
levinvgauss(limit, mean, shape = 1, dispersion = 1/shape, order = 1)
mgfinvgauss(t, mean, shape = 1, dispersion = 1/shape, log = FALSE)
```

### Arguments

<code>x, q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>mean, shape</code>	parameters. Must be strictly positive. Infinite values are supported.
<code>dispersion</code>	an alternative way to specify the shape.
<code>log, log.p</code>	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
<code>order</code>	order of the moment. Only <code>order = 1</code> is supported by <code>levinvgauss</code> .
<code>limit</code>	limit of the loss variable.
<code>tol</code>	small positive value. Tolerance to assess convergence in the Newton computation of quantiles.
<code>maxit</code>	positive integer; maximum number of recursions in the Newton computation of quantiles.
<code>echo, trace</code>	logical; echo the recursions to screen in the Newton computation of quantiles.
<code>t</code>	numeric vector.

### Details

The inverse Gaussian distribution with parameters  $\mu = \mu$  and  $\text{dispersion} = \phi$  has density:

$$f(x) = \left( \frac{1}{2\pi\phi x^3} \right)^{1/2} \exp\left( -\frac{(x - \mu)^2}{2\mu^2\phi x} \right),$$

for  $x \geq 0$ ,  $\mu > 0$  and  $\phi > 0$ .

The limiting case  $\mu = \infty$  is an inverse chi-squared distribution (or inverse gamma with  $\text{shape} = 1/2$  and  $\text{rate} = 2\phi$ ). This distribution has no finite strictly positive, integer moments.

The limiting case  $\phi = 0$  is an infinite spike in  $x = 0$ .

If the random variable  $X$  is  $\text{IG}(\mu, \phi)$ , then  $X/\mu$  is  $\text{IG}(1, \phi\mu)$ .

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $k = 1, 2, \dots$ , the limited expected value at some limit  $d$  is  $E[\min(X, d)]$  and the moment generating function is  $E[e^{tX}]$ .

The moment generating function of the inverse gaussian is defined for  $t \leq 1/(2 * \text{mean}^2 * \phi)$ .

### Value

`dinvgauss` gives the density, `pinvgauss` gives the distribution function, `qinvgauss` gives the quantile function, `rinvgauss` generates random deviates, `minvgauss` gives the  $k$ th raw moment, `levinvgauss` gives the limited expected value, and `mgfinvgauss` gives the moment generating function in `t`.

Invalid arguments will result in return value NaN, with a warning.

**Note**

Functions `dinvgauss`, `pinvgauss` and `qinvgauss` are C implementations of functions of the same name in package **statmod**; see Giner and Smyth (2016).

Devroye (1986, chapter 4) provides a nice presentation of the algorithm to generate random variates from an inverse Gaussian distribution.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>

**References**

Giner, G. and Smyth, G. K. (2016), "**statmod**: Probability Calculations for the Inverse Gaussian Distribution", *R Journal*, vol. 8, no 1, p. 339-351. <https://journal.r-project.org/archive/2016-1/giner-smyth.pdf>

Chhikara, R. S. and Folk, T. L. (1989), *The Inverse Gaussian Distribution: Theory, Methodology and Applications*, Decker.

Devroye, L. (1986), *Non-Uniform Random Variate Generation*, Springer-Verlag. <http://luc.devroye.org/rnbookindex.html>

**See Also**

`dinvgamma` for the inverse gamma distribution.

**Examples**

```
dinvgauss(c(-1, 0, 1, 2, Inf), mean = 1.5, dis = 0.7)
dinvgauss(c(-1, 0, 1, 2, Inf), mean = Inf, dis = 0.7)
dinvgauss(c(-1, 0, 1, 2, Inf), mean = 1.5, dis = Inf) # spike at zero

## Typical graphical representations of the inverse Gaussian
## distribution. First fixed mean and varying shape; second
## varying mean and fixed shape.
col = c("red", "blue", "green", "cyan", "yellow", "black")
par = c(0.125, 0.5, 1, 2, 8, 32)
curve(dinvgauss(x, 1, par[1]), from = 0, to = 2, col = col[1])
for (i in 2:6)
  curve(dinvgauss(x, 1, par[i]), add = TRUE, col = col[i])

curve(dinvgauss(x, par[1], 1), from = 0, to = 2, col = col[1])
for (i in 2:6)
  curve(dinvgauss(x, par[i], 1), add = TRUE, col = col[i])

pinvgauss(qinvgauss((1:10)/10, 1.5, shape = 2), 1.5, 2)

minvgauss(1:4, 1.5, 2)

levinvgauss(c(0, 0.5, 1, 1.2, 10, Inf), 1.5, 2)
```

---

InverseParalogistic    *The Inverse Paralogistic Distribution*


---

**Description**

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Inverse Paralogistic distribution with parameters shape and scale.

**Usage**

```
dinvparalogis(x, shape, rate = 1, scale = 1/rate, log = FALSE)
pinvparalogis(q, shape, rate = 1, scale = 1/rate,
              lower.tail = TRUE, log.p = FALSE)
qinvparalogis(p, shape, rate = 1, scale = 1/rate,
              lower.tail = TRUE, log.p = FALSE)
rinvparalogis(n, shape, rate = 1, scale = 1/rate)
minvparalogis(order, shape, rate = 1, scale = 1/rate)
levinvparalogis(limit, shape, rate = 1, scale = 1/rate,
                order = 1)
```

**Arguments**

x, q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If length(n) > 1, the length is taken to be the number required.
shape, scale	parameters. Must be strictly positive.
rate	an alternative way to specify the scale.
log, log.p	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
order	order of the moment.
limit	limit of the loss variable.

**Details**

The inverse paralogistic distribution with parameters shape =  $\tau$  and scale =  $\theta$  has density:

$$f(x) = \frac{\tau^2(x/\theta)^{\tau^2}}{x[1 + (x/\theta)^\tau]^{\tau+1}}$$

for  $x > 0$ ,  $\tau > 0$  and  $\theta > 0$ .

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $-\tau^2 < k < \tau$ .

The  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ ,  $k > -\tau^2$  and  $1 - k/\tau$  not a negative integer.

**Value**

dinvparalogis gives the density, pinvparalogis gives the distribution function, qinvparalogis gives the quantile function, rinvparalogis generates random deviates, minvparalogis gives the  $k$ th raw moment, and levinvparalogis gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

**Note**

levinvparalogis computes the limited expected value using [betaint](#).

See Kleiber and Kotz (2003) for alternative names and parametrizations.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**Examples**

```
exp(dinvparalogis(2, 3, 4, log = TRUE))
p <- (1:10)/10
pinvparalogis(qinvparalogis(p, 2, 3), 2, 3)

## first negative moment
minvparalogis(-1, 2, 2)

## case with 1 - order/shape > 0
levinvparalogis(10, 2, 2, order = 1)

## case with 1 - order/shape < 0
levinvparalogis(10, 2/3, 2, order = 1)
```

**Description**

Density function, distribution function, quantile function, random generation raw moments and limited moments for the Inverse Pareto distribution with parameters shape and scale.

**Usage**

```
dinvpareto(x, shape, scale, log = FALSE)
pinvpareto(q, shape, scale, lower.tail = TRUE, log.p = FALSE)
qinvpareto(p, shape, scale, lower.tail = TRUE, log.p = FALSE)
rinvpareto(n, shape, scale)
minvpareto(order, shape, scale)
levinvpareto(limit, shape, scale, order = 1)
```

**Arguments**

<code>x, q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>shape, scale</code>	parameters. Must be strictly positive.
<code>log, log.p</code>	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
<code>order</code>	order of the moment.
<code>limit</code>	limit of the loss variable.

**Details**

The inverse Pareto distribution with parameters `shape =  $\tau$`  and `scale =  $\theta$`  has density:

$$f(x) = \frac{\tau\theta x^{\tau-1}}{(x + \theta)^{\tau+1}}$$

for  $x > 0$ ,  $\tau > 0$  and  $\theta > 0$ .

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $-\tau < k < 1$ .

The  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ ,  $k > -\tau$ .

**Value**

`dinvpareto` gives the density, `pinvpareto` gives the distribution function, `qinvpareto` gives the quantile function, `rinvpareto` generates random deviates, `minvpareto` gives the  $k$ th raw moment, and `levinvpareto` calculates the  $k$ th limited moment.

Invalid arguments will result in return value NaN, with a warning.

**Note**

Evaluation of `levinvpareto` is done using numerical integration.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

## References

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

## Examples

```
exp(dinvpareto(2, 3, 4, log = TRUE))
p <- (1:10)/10
pinvpareto(qinvpareto(p, 2, 3), 2, 3)
minvpareto(0.5, 1, 2)
```

---

InverseTransformedGamma

*The Inverse Transformed Gamma Distribution*

---

## Description

Density function, distribution function, quantile function, random generation, raw moments, and limited moments for the Inverse Transformed Gamma distribution with parameters shape1, shape2 and scale.

## Usage

```
dinvtrgamma(x, shape1, shape2, rate = 1, scale = 1/rate,
            log = FALSE)
pinvtrgamma(q, shape1, shape2, rate = 1, scale = 1/rate,
            lower.tail = TRUE, log.p = FALSE)
qinvtrgamma(p, shape1, shape2, rate = 1, scale = 1/rate,
            lower.tail = TRUE, log.p = FALSE)
rinvtrgamma(n, shape1, shape2, rate = 1, scale = 1/rate)
minvtrgamma(order, shape1, shape2, rate = 1, scale = 1/rate)
levinvtrgamma(limit, shape1, shape2, rate = 1, scale = 1/rate,
            order = 1)
```

## Arguments

x, q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If length(n) > 1, the length is taken to be the number required.
shape1, shape2, scale	parameters. Must be strictly positive.
rate	an alternative way to specify the scale.
log, log.p	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
order	order of the moment.
limit	limit of the loss variable.

## Details

The inverse transformed gamma distribution with parameters  $\text{shape1} = \alpha$ ,  $\text{shape2} = \tau$  and  $\text{scale} = \theta$ , has density:

$$f(x) = \frac{\tau u^\alpha e^{-u}}{x \Gamma(\alpha)}, \quad u = (\theta/x)^\tau$$

for  $x > 0$ ,  $\alpha > 0$ ,  $\tau > 0$  and  $\theta > 0$ . (Here  $\Gamma(\alpha)$  is the function implemented by R's `gamma()` and defined in its help.)

The inverse transformed gamma is the distribution of the random variable  $\theta X^{-1/\tau}$ , where  $X$  has a gamma distribution with shape parameter  $\alpha$  and scale parameter 1 or, equivalently, of the random variable  $Y^{-1/\tau}$  with  $Y$  a gamma distribution with shape parameter  $\alpha$  and scale parameter  $\theta^{-\tau}$ .

The inverse transformed gamma distribution defines a family of distributions with the following special cases:

- An **Inverse Gamma** distribution when  $\text{shape2} == 1$ ;
- An **Inverse Weibull** distribution when  $\text{shape1} == 1$ ;
- An **Inverse Exponential** distribution when  $\text{shape1} == \text{shape2} == 1$ ;

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $k < \alpha\tau$ , and the  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$  for all  $k$ .

## Value

`dinvtrgamma` gives the density, `pinvtrgamma` gives the distribution function, `qinvtrgamma` gives the quantile function, `rinvtrgamma` generates random deviates, `minvtrgamma` gives the  $k$ th raw moment, and `levinvtrgamma` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

## Note

`levinvtrgamma` computes the limited expected value using `gammainc` from package **expint**.

Distribution also known as the Inverse Generalized Gamma. See also Kleiber and Kotz (2003) for alternative names and parametrizations.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

## Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

## References

- Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.
- Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**Examples**

```
exp(dinvtrgamma(2, 3, 4, 5, log = TRUE))
p <- (1:10)/10
pinvtrgamma(qinvtrgamma(p, 2, 3, 4), 2, 3, 4)
minvtrgamma(2, 3, 4, 5)
levinvtrgamma(200, 3, 4, 5, order = 2)
```

InverseWeibull

*The Inverse Weibull Distribution***Description**

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Inverse Weibull distribution with parameters shape and scale.

**Usage**

```
dinvweibull(x, shape, rate = 1, scale = 1/rate, log = FALSE)
pinvweibull(q, shape, rate = 1, scale = 1/rate,
            lower.tail = TRUE, log.p = FALSE)
qinvweibull(p, shape, rate = 1, scale = 1/rate,
            lower.tail = TRUE, log.p = FALSE)
rinvweibull(n, shape, rate = 1, scale = 1/rate)
minvweibull(order, shape, rate = 1, scale = 1/rate)
levinvweibull(limit, shape, rate = 1, scale = 1/rate,
              order = 1)
```

**Arguments**

<code>x, q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>shape, scale</code>	parameters. Must be strictly positive.
<code>rate</code>	an alternative way to specify the scale.
<code>log, log.p</code>	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
<code>order</code>	order of the moment.
<code>limit</code>	limit of the loss variable.



**Details**

The inverse Weibull distribution with parameters  $\text{shape} = \tau$  and  $\text{scale} = \theta$  has density:

$$f(x) = \frac{\tau(\theta/x)^\tau e^{-(\theta/x)^\tau}}{x}$$

for  $x > 0$ ,  $\tau > 0$  and  $\theta > 0$ .

The special case  $\text{shape} = 1$  is an [Inverse Exponential](#) distribution.

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $k < \tau$ , and the  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ , all  $k$ .

**Value**

`dinvweibull` gives the density, `pinvweibull` gives the distribution function, `qinvweibull` gives the quantile function, `rinvweibull` generates random deviates, `minvweibull` gives the  $k$ th raw moment, and `levinvweibull` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

**Note**

`levinvweibull` computes the limited expected value using `gammainc` from package **expint**.

Distribution also known as the log-Gompertz. See also Kleiber and Kotz (2003) for alternative names and parametrizations.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**Examples**

```
exp(dinvweibull(2, 3, 4, log = TRUE))
p <- (1:10)/10
pinvweibull(qinvweibull(p, 2, 3), 2, 3)
mlgompertz(-1, 3, 3)
levinvweibull(10, 2, 3, order = 1)
```

---

 Logarithmic

*The Logarithmic Distribution*


---

### Description

Density function, distribution function, quantile function and random generation for the Logarithmic (or log-series) distribution with parameter `prob`.

### Usage

```
dlogarithmic(x, prob, log = FALSE)
plogarithmic(q, prob, lower.tail = TRUE, log.p = FALSE)
qlogarithmic(p, prob, lower.tail = TRUE, log.p = FALSE)
rlogarithmic(n, prob)
```

### Arguments

<code>x</code>	vector of (strictly positive integer) quantiles.
<code>q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>prob</code>	parameter. $0 \leq \text{prob} < 1$ .
<code>log, log.p</code>	logical; if TRUE, probabilities $p$ are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .

### Details

The logarithmic (or log-series) distribution with parameter `prob` =  $\theta$  has probability mass function

$$p(x) = \frac{a\theta^x}{x},$$

with  $a = -1/\log(1 - \theta)$  and for  $x = 1, 2, \dots, 0 \leq \theta < 1$ .

The logarithmic distribution is the limiting case of the zero-truncated negative binomial distribution with size parameter equal to 0. Note that in this context, parameter `prob` generally corresponds to the probability of *failure* of the zero-truncated negative binomial.

If an element of `x` is not integer, the result of `dlogarithmic` is zero, with a warning.

The quantile is defined as the smallest value  $x$  such that  $F(x) \geq p$ , where  $F$  is the distribution function.

**Value**

dlogarithmic gives the probability mass function, plogarithmic gives the distribution function, qllogarithmic gives the quantile function, and rlogarithmic generates random deviates.

Invalid prob will result in return value NaN, with a warning.

The length of the result is determined by n for rlogarithmic, and is the maximum of the lengths of the numerical arguments for the other functions.

**Note**

qllogarithmic is based on qbinom et al.; it uses the Cornish–Fisher Expansion to include a skewness correction to a normal approximation, followed by a search.

rlogarithmic is an implementation of the LS and LK algorithms of Kemp (1981) with automatic selection. As suggested by Devroye (1986), the LS algorithm is used when  $\text{prob} < 0.95$ , and the LK algorithm otherwise.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>

**References**

Johnson, N. L., Kemp, A. W. and Kotz, S. (2005), *Univariate Discrete Distributions, Third Edition*, Wiley.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

Kemp, A. W. (1981), “Efficient Generation of Logarithmically Distributed Pseudo-Random Variables”, *Journal of the Royal Statistical Society, Series C*, vol. 30, p. 249-253.

Devroye, L. (1986), *Non-Uniform Random Variate Generation*, Springer-Verlag. <http://luc.devroye.org/rnbookindex.html>

**See Also**

[dztnbinom](#) for the zero-truncated negative binomial distribution.

**Examples**

```
## Table 1 of Kemp (1981) [also found in Johnson et al. (2005), chapter 7]
p <- c(0.1, 0.3, 0.5, 0.7, 0.8, 0.85, 0.9, 0.95, 0.99, 0.995, 0.999, 0.9999)
round(rbind(dlogarithmic(1, p),
            dlogarithmic(2, p),
            plogarithmic(9, p, lower.tail = FALSE),
            -p/((1 - p) * log(1 - p))), 2)

qllogarithmic(plogarithmic(1:10, 0.9), 0.9)

x <- rlogarithmic(1000, 0.8)
y <- sort(unique(x))
plot(y, table(x)/length(x), type = "h", lwd = 2,
```

```

    pch = 19, col = "black", xlab = "x", ylab = "p(x)",
    main = "Empirical vs theoretical probabilities")
points(y, dlogarithmic(y, prob = 0.8),
       pch = 19, col = "red")
legend("topright", c("empirical", "theoretical"),
       lty = c(1, NA), pch = c(NA, 19), col = c("black", "red"))

```

Loggamma

*The Loggamma Distribution***Description**

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Loggamma distribution with parameters `shapelog` and `ratelog`.

**Usage**

```

dlgamma(x, shapelog, ratelog, log = FALSE)
plgamma(q, shapelog, ratelog, lower.tail = TRUE, log.p = FALSE)
qlgamma(p, shapelog, ratelog, lower.tail = TRUE, log.p = FALSE)
rlgamma(n, shapelog, ratelog)
mlgamma(order, shapelog, ratelog)
levlgamma(limit, shapelog, ratelog, order = 1)

```

**Arguments**

<code>x, q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>shapelog, ratelog</code>	parameters. Must be strictly positive.
<code>log, log.p</code>	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
<code>order</code>	order of the moment.
<code>limit</code>	limit of the loss variable.

**Details**

The loggamma distribution with parameters `shapelog` =  $\alpha$  and `ratelog` =  $\lambda$  has density:

$$f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} \frac{(\log x)^{\alpha-1}}{x^{\lambda+1}}$$

for  $x > 1$ ,  $\alpha > 0$  and  $\lambda > 0$ . (Here  $\Gamma(\alpha)$  is the function implemented by R's `gamma()` and defined in its help.)

The loggamma is the distribution of the random variable  $e^{-X}$ , where  $X$  has a gamma distribution with shape parameter  $alpha$  and scale parameter  $1/\lambda$ .

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$  and the  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ ,  $k < \lambda$ .

### Value

`dlgamma` gives the density, `plgamma` gives the distribution function, `qlgamma` gives the quantile function, `rlgamma` generates random deviates, `mlgamma` gives the  $k$ th raw moment, and `levlgamma` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

### Note

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

### References

Hogg, R. V. and Klugman, S. A. (1984), *Loss Distributions*, Wiley.

### Examples

```
exp(dlgamma(2, 3, 4, log = TRUE))
p <- (1:10)/10
plgamma(qlgamma(p, 2, 3), 2, 3)
mlgamma(2, 3, 4) - mlgamma(1, 3, 4)^2
levlgamma(10, 3, 4, order = 2)
```

### Description

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Loglogistic distribution with parameters shape and scale.

**Usage**

```

dllogis(x, shape, rate = 1, scale = 1/rate, log = FALSE)
pllogis(q, shape, rate = 1, scale = 1/rate,
        lower.tail = TRUE, log.p = FALSE)
qllogis(p, shape, rate = 1, scale = 1/rate,
        lower.tail = TRUE, log.p = FALSE)
rllogis(n, shape, rate = 1, scale = 1/rate)
mllogis(order, shape, rate = 1, scale = 1/rate)
levllogis(limit, shape, rate = 1, scale = 1/rate,
          order = 1)

```

**Arguments**

<code>x, q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>shape, scale</code>	parameters. Must be strictly positive.
<code>rate</code>	an alternative way to specify the scale.
<code>log, log.p</code>	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
<code>order</code>	order of the moment.
<code>limit</code>	limit of the loss variable.

**Details**

The loglogistic distribution with parameters  $\text{shape} = \gamma$  and  $\text{scale} = \theta$  has density:

$$f(x) = \frac{\gamma(x/\theta)^\gamma}{x[1 + (x/\theta)^\gamma]^2}$$

for  $x > 0$ ,  $\gamma > 0$  and  $\theta > 0$ .

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $-\gamma < k < \gamma$ .

The  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ ,  $k > -\gamma$  and  $1 - k/\gamma$  not a negative integer.

**Value**

`dllogis` gives the density, `pllogis` gives the distribution function, `qllogis` gives the quantile function, `rllogis` generates random deviates, `mllogis` gives the  $k$ th raw moment, and `levllogis` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

**Note**

levllogis computes the limited expected value using [betaint](#).

Also known as the Fisk distribution. See also Kleiber and Kotz (2003) for alternative names and parametrizations.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**See Also**

[dpareto3](#) for an equivalent distribution with a location parameter.

**Examples**

```
exp(dllogis(2, 3, 4, log = TRUE))
p <- (1:10)/10
pllogis(qllogis(p, 2, 3), 2, 3)

## mean
mllogis(1, 2, 3)

## case with 1 - order/shape > 0
levllogis(10, 2, 3, order = 1)

## case with 1 - order/shape < 0
levllogis(10, 2/3, 3, order = 1)
```

**Description**

Raw moments and limited moments for the Lognormal distribution whose logarithm has mean equal to meanlog and standard deviation equal to sdlog.

**Usage**

```
mlnorm(order, meanlog = 0, sdlog = 1)
levlnorm(limit, meanlog = 0, sdlog = 1, order = 1)
```

**Arguments**

order	order of the moment.
limit	limit of the loss variable.
meanlog, sdlog	mean and standard deviation of the distribution on the log scale with default values of 0 and 1 respectively.

**Value**

`mlnorm` gives the  $k$ th raw moment and `levlnorm` gives the  $k$ th moment of the limited loss variable. Invalid arguments will result in return value NaN, with a warning.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**See Also**

[Lognormal](#) for details on the lognormal distribution and functions `[dpqr]lnorm`.

**Examples**

```
mlnorm(2, 3, 4) - mlnorm(1, 3, 4)^2
levlnorm(10, 3, 4, order = 2)
```

---

mde

*Minimum Distance Estimation*


---

**Description**

Minimum distance fitting of univariate distributions, allowing parameters to be held fixed if desired.

**Usage**

```
mde(x, fun, start, measure = c("CvM", "chi-square", "LAS"),
    weights = NULL, ...)
```



**Arguments**

x	a vector or an object of class "grouped data" (in which case only the first column of frequencies is used).
fun	function returning a cumulative distribution (for measure = "CvM" and measure = "chi-square") or a limited expected value (for measure = "LAS") evaluated at its first argument.
start	a named list giving the parameters to be optimized with initial values
measure	either "CvM" for the Cramer-von Mises method, "chi-square" for the modified chi-square method, or "LAS" for the layer average severity method.
weights	weights; see details.
...	Additional parameters, either for fun or for optim. In particular, it can be used to specify bounds via lower or upper or both. If arguments of fun are included they will be held fixed.

**Details**

The Cramer-von Mises method ("CvM") minimizes the squared difference between the theoretical cdf and the empirical cdf at the data points (for individual data) or the ogive at the knots (for grouped data).

The modified chi-square method ("chi-square") minimizes the modified chi-square statistic for grouped data, that is the squared difference between the expected and observed frequency within each group.

The layer average severity method ("LAS") minimizes the squared difference between the theoretical and empirical limited expected value within each group for grouped data.

All sum of squares can be weighted. If arguments weights is missing, weights default to 1 for measure = "CvM" and measure = "LAS"; for measure = "chi-square", weights default to  $1/n_j$ , where  $n_j$  is the frequency in group  $j = 1, \dots, r$ .

Optimization is performed using `optim`. For one-dimensional problems the Nelder-Mead method is used and for multi-dimensional problems the BFGS method, unless arguments named lower or upper are supplied when L-BFGS-B is used or method is supplied explicitly.

**Value**

An object of class "mde", a list with two components:

estimate	the parameter estimates, and
distance	the distance.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (1998), *Loss Models, From Data to Decisions*, Wiley.

**Examples**

```
## Individual data example
data(dental)
mde(dental, pexp, start = list(rate = 1/200), measure = "CvM")

## Example 2.21 of Klugman et al. (1998)
data(gdental)
mde(gdental, pexp, start = list(rate = 1/200), measure = "CvM")
mde(gdental, pexp, start = list(rate = 1/200), measure = "chi-square")
mde(gdental, levexp, start = list(rate = 1/200), measure = "LAS")

## Two-parameter distribution example
try(mde(gdental, ppareto, start = list(shape = 3, scale = 600),
       measure = "CvM")) # no convergence

## Working in log scale often solves the problem
ppareto.log <- function(x, shape, scale)
  ppareto(x, exp(shape), exp(scale))

( p <- mde(gdental, ppareto.log, start = list(shape = log(3),
       scale = log(600)), measure = "CvM") )
exp(p$estimate)
```

---

mean.grouped.data      *Arithmetic Mean*

---

**Description**

Mean of grouped data objects.

**Usage**

```
## S3 method for class 'grouped.data'
mean(x, ...)
```

**Arguments**

x                    an object of class "grouped.data".  
 ...                  further arguments passed to or from other methods.

**Details**

The mean of grouped data with group boundaries  $c_1, \dots, c_r$  and group frequencies  $n_1, \dots, n_r$  is

$$\sum_{j=1}^r \frac{c_{j-1} + c_j}{2} n_j.$$

**Value**

A named vector of means.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>

**References**

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (1998), *Loss Models, From Data to Decisions*, Wiley.

**See Also**

[grouped.data](#) to create grouped data objects; [emm](#) to compute higher moments.

**Examples**

```
data(gdental)
mean(gdental)
```

---

NormalSupp

*Moments and Moment generating function of the Normal Distribution*

---

**Description**

Raw moments and moment generating function for the normal distribution with mean equal to mean and standard deviation equal to sd.

**Usage**

```
mnorm(order, mean = 0, sd = 1)
mgfnorm(t, mean = 0, sd = 1, log = FALSE)
```

**Arguments**

order	vector of integers; order of the moment.
mean	vector of means.
sd	vector of standard deviations.
t	numeric vector.
log	logical; if TRUE, the cumulant generating function is returned.

**Details**

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$  and the moment generating function is  $E[e^{tX}]$ .

Only integer moments are supported.

**Value**

`mnorm` gives the  $k$ th raw moment and `mgfnorm` gives the moment generating function in `t`.  
Invalid arguments will result in return value `NaN`, with a warning.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>, Christophe Dutang

**References**

Johnson, N. L. and Kotz, S. (1970), *Continuous Univariate Distributions, Volume 1*, Wiley.

**See Also**

[Normal](#)

**Examples**

```
mgfnorm(0:4,1,2)
mnorm(3)
```

---

ogive

*Ogive for Grouped Data*

---

**Description**

Compute a smoothed empirical distribution function for grouped data.

**Usage**

```
ogive(x, ...)

## Default S3 method:
ogive(x, y = NULL, breaks = "Sturges", nclass = NULL, ...)

## S3 method for class 'grouped.data'
ogive(x, ...)

## S3 method for class 'ogive'
print(x, digits = getOption("digits") - 2, ...)

## S3 method for class 'ogive'
summary(object, ...)

## S3 method for class 'ogive'
knots(Fn, ...)

## S3 method for class 'ogive'
plot(x, main = NULL, xlab = "x", ylab = "F(x)", ...)
```

**Arguments**

x	for the generic and all but the default method, an object of class "grouped.data"; for the default method, a vector of individual data if y is NULL, a vector of group boundaries otherwise.
y	a vector of group frequencies.
breaks, nclass	arguments passed to <a href="#">grouped.data</a> ; used only for individual data (when y is NULL).
digits	number of significant digits to use, see <a href="#">print</a> .
Fn, object	an R object inheriting from "ogive".
main	main title.
xlab, ylab	labels of x and y axis.
...	arguments to be passed to subsequent methods.

**Details**

The ogive is a linear interpolation of the empirical cumulative distribution function.

The equation of the ogive is

$$G_n(x) = \frac{(c_j - x)F_n(c_{j-1}) + (x - c_{j-1})F_n(c_j)}{c_j - c_{j-1}}$$

for  $c_{j-1} < x \leq c_j$  and where  $c_0, \dots, c_r$  are the  $r + 1$  group boundaries and  $F_n$  is the empirical distribution function of the sample.

**Value**

For ogive, a function of class "ogive", inheriting from the "function" class.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (1998), *Loss Models, From Data to Decisions*, Wiley.

**See Also**

[grouped.data](#) to create grouped data objects; [quantile.grouped.data](#) for the inverse function; [approxfun](#), which is used to compute the ogive; [stepfun](#) for related documentation (even though the ogive is not a step function).

**Examples**

```
## Most common usage: create ogive from grouped data object.
Fn <- ogive(gdental)
Fn
summary(Fn)
knots(Fn)                # the group boundaries

Fn(knots(Fn))            # true values of the empirical cdf
Fn(c(80, 200, 2000))    # linear interpolations

plot(Fn)                 # graphical representation

## Alternative 1: create ogive directly from individual data
## without first creating a grouped data object.
ogive(dental)           # automatic class boundaries
ogive(dental, breaks = c(0, 50, 200, 500, 1500, 2000))

## Alternative 2: create ogive from set of group boundaries and
## group frequencies.
cj <- c(0, 25, 50, 100, 250, 500, 1000)
nj <- c(30, 31, 57, 42, 45, 10)
ogive(cj, nj)
```

---

Paralogistic

*The Paralogistic Distribution*


---

**Description**

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Paralogistic distribution with parameters shape and scale.

**Usage**

```
dparalogis(x, shape, rate = 1, scale = 1/rate, log = FALSE)
pparalogis(q, shape, rate = 1, scale = 1/rate,
            lower.tail = TRUE, log.p = FALSE)
qparalogis(p, shape, rate = 1, scale = 1/rate,
            lower.tail = TRUE, log.p = FALSE)
rparalogis(n, shape, rate = 1, scale = 1/rate)
mparalogis(order, shape, rate = 1, scale = 1/rate)
levparalogis(limit, shape, rate = 1, scale = 1/rate,
              order = 1)
```

**Arguments**

x, q                    vector of quantiles.  
p                        vector of probabilities.

n	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
shape, scale	parameters. Must be strictly positive.
rate	an alternative way to specify the scale.
log, log.p	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
order	order of the moment.
limit	limit of the loss variable.

### Details

The paralogistic distribution with parameters  $\text{shape} = \alpha$  and  $\text{scale} = \theta$  has density:

$$f(x) = \frac{\alpha^2 (x/\theta)^\alpha}{x[1 + (x/\theta)^\alpha]^{\alpha+1}}$$

for  $x > 0$ ,  $\alpha > 0$  and  $\theta > 0$ .

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $-\alpha < k < \alpha^2$ .

The  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ ,  $k > -\alpha$  and  $\alpha - k/\alpha$  not a negative integer.

### Value

`dparalogis` gives the density, `pparalogis` gives the distribution function, `qparalogis` gives the quantile function, `rparalogis` generates random deviates, `mparalogis` gives the  $k$ th raw moment, and `levparalogis` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

### Note

`levparalogis` computes the limited expected value using `betaint`.

See Kleiber and Kotz (2003) for alternative names and parametrizations.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

### References

- Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.
- Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**Examples**

```

exp(dparalogis(2, 3, 4, log = TRUE))
p <- (1:10)/10
pparalogis(qparalogis(p, 2, 3), 2, 3)

## variance
mparalogis(2, 2, 3) - mparalogis(1, 2, 3)^2

## case with shape - order/shape > 0
levparalogis(10, 2, 3, order = 2)

## case with shape - order/shape < 0
levparalogis(10, 1.25, 3, order = 2)

```

---

Pareto

*The Pareto Distribution*


---

**Description**

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Pareto distribution with parameters shape and scale.

**Usage**

```

dpareto(x, shape, scale, log = FALSE)
ppareto(q, shape, scale, lower.tail = TRUE, log.p = FALSE)
qpareto(p, shape, scale, lower.tail = TRUE, log.p = FALSE)
rpareto(n, shape, scale)
mpareto(order, shape, scale)
levpareto(limit, shape, scale, order = 1)

```

**Arguments**

x, q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If length(n) > 1, the length is taken to be the number required.
shape, scale	parameters. Must be strictly positive.
log, log.p	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
order	order of the moment.
limit	limit of the loss variable.



**Details**

The Pareto distribution with parameters  $\text{shape} = \alpha$  and  $\text{scale} = \theta$  has density:

$$f(x) = \frac{\alpha\theta^\alpha}{(x + \theta)^{\alpha+1}}$$

for  $x > 0$ ,  $\alpha > 0$  and  $\theta$ .

There are many different definitions of the Pareto distribution in the literature; see Arnold (2015) or Kleiber and Kotz (2003). In the nomenclature of **actuar**, The “Pareto distribution” does not have a location parameter. The version with a location parameter is the [Pareto II](#).

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $-1 < k < \alpha$ .

The  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ ,  $k > -1$  and  $\alpha - k$  not a negative integer.

**Value**

`dpareto` gives the density, `ppareto` gives the distribution function, `qpareto` gives the quantile function, `rpareto` generates random deviates, `mpareto` gives the  $k$ th raw moment, and `levpareto` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

**Note**

`levpareto` computes the limited expected value using [betaint](#).

The version of the Pareto defined for  $x > \theta$  is named Single Parameter Pareto, or Pareto I, in **actuar**.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**See Also**

[dpareto2](#) for an equivalent distribution with location parameter.

[dpareto1](#) for the Single Parameter Pareto distribution.

"distributions" package vignette for details on the interrelations between the continuous size distributions in **actuar** and complete formulas underlying the above functions.

**Examples**

```

exp(dpareto(2, 3, 4, log = TRUE))
p <- (1:10)/10
ppareto(qpareto(p, 2, 3), 2, 3)

## variance
mpareto(2, 4, 1) - mpareto(1, 4, 1)^2

## case with shape - order > 0
levpareto(10, 3, scale = 1, order = 2)

## case with shape - order < 0
levpareto(10, 1.5, scale = 1, order = 2)

```

Pareto2

*The Pareto II Distribution***Description**

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Pareto II distribution with parameters min, shape and scale.

**Usage**

```

dpareto2(x, min, shape, rate = 1, scale = 1/rate,
         log = FALSE)
ppareto2(q, min, shape, rate = 1, scale = 1/rate,
         lower.tail = TRUE, log.p = FALSE)
qpareto2(p, min, shape, rate = 1, scale = 1/rate,
         lower.tail = TRUE, log.p = FALSE)
rpareto2(n, min, shape, rate = 1, scale = 1/rate)
mpareto2(order, min, shape, rate = 1, scale = 1/rate)
levpareto2(limit, min, shape, rate = 1, scale = 1/rate,
           order = 1)

```

**Arguments**

x, q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If length(n) > 1, the length is taken to be the number required.
min	lower bound of the support of the distribution.
shape, scale	parameters. Must be strictly positive.
rate	an alternative way to specify the scale.
log, log.p	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
order	order of the moment.
limit	limit of the loss variable.

### Details

The Pareto II (or “type II”) distribution with parameters  $\text{min} = \mu$ ,  $\text{shape} = \alpha$  and  $\text{scale} = \theta$  has density:

$$f(x) = \frac{\alpha}{\theta[1 + (x - \mu)/\theta]^{\alpha+1}}$$

for  $x > \mu$ ,  $-\infty < \mu < \infty$ ,  $\alpha > 0$  and  $\theta > 0$ .

The Pareto II is the distribution of the random variable

$$\mu + \theta \left( \frac{X}{1 - X} \right),$$

where  $X$  has a beta distribution with parameters 1 and  $\alpha$ . It derives from the [Feller-Pareto](#) distribution with  $\tau = \gamma = 1$ . Setting  $\mu = 0$  yields the familiar [Pareto](#) distribution.

The [Pareto I](#) (or Single parameter Pareto) distribution is a special case of the Pareto II with  $\text{min} = \text{scale}$ .

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$  for nonnegative integer values of  $k < \alpha$ .

The  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$  for nonnegative integer values of  $k$  and  $\alpha - j$ ,  $j = 1, \dots, k$  not a negative integer.

### Value

`dpareto2` gives the density, `ppareto2` gives the distribution function, `qpareto2` gives the quantile function, `rpareto2` generates random deviates, `mpareto2` gives the  $k$ th raw moment, and `levpareto2` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

### Note

`levpareto2` computes the limited expected value using [betaint](#).

For Pareto distributions, we use the classification of Arnold (2015) with the parametrization of Klugman et al. (2012).

The “distributions” package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>

### References

- Arnold, B.C. (2015), *Pareto Distributions*, Second Edition, CRC Press.
- Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.
- Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**See Also**

[dpareto](#) for the Pareto distribution without a location parameter.

**Examples**

```
exp(dpareto2(1, min = 10, 3, 4, log = TRUE))
p <- (1:10)/10
ppareto2(qpareto2(p, min = 10, 2, 3), min = 10, 2, 3)

## variance
mpareto2(2, min = 10, 4, 1) - mpareto2(1, min = 10, 4, 1)^2

## case with shape - order > 0
levpareto2(10, min = 10, 3, scale = 1, order = 2)

## case with shape - order < 0
levpareto2(10, min = 10, 1.5, scale = 1, order = 2)
```

---

Pareto3

*The Pareto III Distribution*


---

**Description**

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Pareto III distribution with parameters `min`, `shape` and `scale`.

**Usage**

```
dpareto3(x, min, shape, rate = 1, scale = 1/rate,
         log = FALSE)
ppareto3(q, min, shape, rate = 1, scale = 1/rate,
         lower.tail = TRUE, log.p = FALSE)
qpareto3(p, min, shape, rate = 1, scale = 1/rate,
         lower.tail = TRUE, log.p = FALSE)
rpareto3(n, min, shape, rate = 1, scale = 1/rate)
mpareto3(order, min, shape, rate = 1, scale = 1/rate)
levpareto3(limit, min, shape, rate = 1, scale = 1/rate,
           order = 1)
```

**Arguments**

<code>x</code> , <code>q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>min</code>	lower bound of the support of the distribution.
<code>shape</code> , <code>scale</code>	parameters. Must be strictly positive.

rate	an alternative way to specify the scale.
log, log.p	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
order	order of the moment.
limit	limit of the loss variable.

### Details

The Pareto III (or “type III”) distribution with parameters  $\text{min} = \mu$ ,  $\text{shape} = \gamma$  and  $\text{scale} = \theta$  has density:

$$f(x) = \frac{\gamma((x - \mu)/\theta)^{\gamma-1}}{\theta[1 + ((x - \mu)/\theta)^\gamma]^2}$$

for  $x > \mu$ ,  $-\infty < \mu < \infty$ ,  $\gamma > 0$  and  $\theta > 0$ .

The Pareto III is the distribution of the random variable

$$\mu + \theta \left( \frac{X}{1 - X} \right)^{1/\gamma},$$

where  $X$  has a uniform distribution on  $(0, 1)$ . It derives from the [Feller-Pareto](#) distribution with  $\alpha = \tau = 1$ . Setting  $\mu = 0$  yields the [loglogistic](#) distribution.

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$  for nonnegative integer values of  $k < \gamma$ .

The  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$  for nonnegative integer values of  $k$  and  $1 - j/\gamma$ ,  $j = 1, \dots, k$  not a negative integer.

### Value

`dpareto3` gives the density, `ppareto3` gives the distribution function, `qpareto3` gives the quantile function, `rpareto3` generates random deviates, `mpareto3` gives the  $k$ th raw moment, and `levpareto3` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

### Note

`levpareto3` computes the limited expected value using [betaint](#).

For Pareto distributions, we use the classification of Arnold (2015) with the parametrization of Klugman et al. (2012).

The “distributions” package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>

## References

- Arnold, B.C. (2015), *Pareto Distributions*, Second Edition, CRC Press.
- Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.
- Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

## See Also

[dllogis](#) for the loglogistic distribution.

## Examples

```
exp(dpareto3(1, min = 10, 3, 4, log = TRUE))
p <- (1:10)/10
ppareto3(qpareto3(p, min = 10, 2, 3), min = 10, 2, 3)

## mean
mpareto3(1, min = 10, 2, 3)

## case with 1 - order/shape > 0
levpareto3(20, min = 10, 2, 3, order = 1)

## case with 1 - order/shape < 0
levpareto3(20, min = 10, 2/3, 3, order = 1)
```

---

Pareto4

*The Pareto IV Distribution*

---

## Description

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Pareto IV distribution with parameters `min`, `shape1`, `shape2` and `scale`.

## Usage

```
dpareto4(x, min, shape1, shape2, rate = 1, scale = 1/rate,
         log = FALSE)
ppareto4(q, min, shape1, shape2, rate = 1, scale = 1/rate,
         lower.tail = TRUE, log.p = FALSE)
qpareto4(p, min, shape1, shape2, rate = 1, scale = 1/rate,
         lower.tail = TRUE, log.p = FALSE)
rpareto4(n, min, shape1, shape2, rate = 1, scale = 1/rate)
mpareto4(order, min, shape1, shape2, rate = 1, scale = 1/rate)
levpareto4(limit, min, shape1, shape2, rate = 1, scale = 1/rate,
           order = 1)
```

**Arguments**

<code>x, q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>min</code>	lower bound of the support of the distribution.
<code>shape1, shape2, scale</code>	parameters. Must be strictly positive.
<code>rate</code>	an alternative way to specify the scale.
<code>log, log.p</code>	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
<code>order</code>	order of the moment.
<code>limit</code>	limit of the loss variable.

**Details**

The Pareto IV (or “type IV”) distribution with parameters  $\text{min} = \mu$ ,  $\text{shape1} = \alpha$ ,  $\text{shape2} = \gamma$  and  $\text{scale} = \theta$  has density:

$$f(x) = \frac{\alpha\gamma((x - \mu)/\theta)^{\gamma-1}}{\theta[1 + ((x - \mu)/\theta)^\gamma]^{\alpha+1}}$$

for  $x > \mu$ ,  $-\infty < \mu < \infty$ ,  $\alpha > 0$ ,  $\gamma > 0$  and  $\theta > 0$ .

The Pareto IV is the distribution of the random variable

$$\mu + \theta \left( \frac{X}{1 - X} \right)^{1/\gamma},$$

where  $X$  has a beta distribution with parameters 1 and  $\alpha$ . It derives from the [Feller-Pareto](#) distribution with  $\tau = 1$ . Setting  $\mu = 0$  yields the [Burr](#) distribution.

The Pareto IV distribution also has the following direct special cases:

- A [Pareto III](#) distribution when  $\text{shape1} == 1$ ;
- A [Pareto II](#) distribution when  $\text{shape1} == 1$ .

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$  for nonnegative integer values of  $k < \alpha\gamma$ .

The  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$  for nonnegative integer values of  $k$  and  $\alpha - j/\gamma, j = 1, \dots, k$  not a negative integer.

**Value**

`dpareto4` gives the density, `ppareto4` gives the distribution function, `qpareto4` gives the quantile function, `rpareto4` generates random deviates, `mpareto4` gives the  $k$ th raw moment, and `levpareto4` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

**Note**

levpareto4 computes the limited expected value using [betaint](#).

For Pareto distributions, we use the classification of Arnold (2015) with the parametrization of Klugman et al. (2012).

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>

**References**

Arnold, B.C. (2015), *Pareto Distributions*, Second Edition, CRC Press.

Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**See Also**

[dburr](#) for the Burr distribution.

**Examples**

```
exp(dpareto4(1, min = 10, 2, 3, log = TRUE))
p <- (1:10)/10
ppareto4(qpareto4(p, min = 10, 2, 3, 2), min = 10, 2, 3, 2)

## variance
mpareto4(2, min = 10, 2, 3, 1) - mpareto4(1, min = 10, 2, 3, 1) ^ 2

## case with shape1 - order/shape2 > 0
levpareto4(10, min = 10, 2, 3, 1, order = 2)

## case with shape1 - order/shape2 < 0
levpareto4(10, min = 10, 1.5, 0.5, 1, order = 2)
```

**Description**

Density, distribution function, random generation, raw moments and moment generating function for the (continuous) Phase-type distribution with parameters prob and rates.



**Usage**

```

dphtype(x, prob, rates, log = FALSE)
pphtype(q, prob, rates, lower.tail = TRUE, log.p = FALSE)
rphtype(n, prob, rates)
mphtype(order, prob, rates)
mgfphtype(t, prob, rates, log = FALSE)

```

**Arguments**

<code>x, q</code>	vector of quantiles.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>prob</code>	vector of initial probabilities for each of the transient states of the underlying Markov chain. The initial probability of the absorbing state is $1 - \text{sum}(\text{prob})$ .
<code>rates</code>	square matrix of the rates of transition among the states of the underlying Markov chain.
<code>log, log.p</code>	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
<code>order</code>	order of the moment.
<code>t</code>	numeric vector.

**Details**

The phase-type distribution with parameters  $\text{prob} = \pi$  and  $\text{rates} = \mathbf{T}$  has density:

$$f(x) = \pi e^{\mathbf{T}x} \mathbf{t}$$

for  $x \geq 0$  and  $f(0) = 1 - \pi e$ , where  $e$  is a column vector with all components equal to one,  $\mathbf{t} = -\mathbf{T}e$  is the exit rates vector and  $e^{\mathbf{T}x}$  denotes the matrix exponential of  $\mathbf{T}x$ . The matrix exponential of a matrix  $\mathbf{M}$  is defined as the Taylor series

$$e^{\mathbf{M}} = \sum_{n=0}^{\infty} \frac{\mathbf{M}^n}{n!}.$$

The parameters of the distribution must satisfy  $\pi e \leq 1$ ,  $\mathbf{T}_{ii} < 0$ ,  $\mathbf{T}_{ij} \geq 0$  and  $\mathbf{T}e \leq 0$ .

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$  and the moment generating function is  $E[e^{tX}]$ .

**Value**

`dphasetype` gives the density, `pphasetype` gives the distribution function, `rphasetype` generates random deviates, `mphasetype` gives the  $k$ th raw moment, and `mgfpphasetype` gives the moment generating function in  $x$ .

Invalid arguments will result in return value NaN, with a warning.

**Note**

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Christophe Dutang

**References**

[https://en.wikipedia.org/wiki/Phase-type\\_distribution](https://en.wikipedia.org/wiki/Phase-type_distribution)

Neuts, M. F. (1981), *Generating random variates from a distribution of phase type*, WSC '81: Proceedings of the 13th conference on Winter simulation, IEEE Press.

**Examples**

```
## Erlang(3, 2) distribution
T <- cbind(c(-2, 0, 0), c(2, -2, 0), c(0, 2, -2))
pi <- c(1,0,0)
x <- 0:10

dphtype(x, pi, T) # density
dgamma(x, 3, 2) # same
pphtype(x, pi, T) # cdf
pgamma(x, 3, 2) # same

rphtype(10, pi, T) # random values
mphtype(1, pi, T) # expected value

curve(mgfphtype(x, pi, T), from = -10, to = 1)
```

---

PoissonInverseGaussian

*The Poisson-Inverse Gaussian Distribution*

---

**Description**

Density function, distribution function, quantile function and random generation for the Poisson-inverse Gaussian discrete distribution with parameters mean and shape.

**Usage**

```
dpoisinvgauss(x, mean, shape = 1, dispersion = 1/shape,
log = FALSE)
ppoisinvgauss(q, mean, shape = 1, dispersion = 1/shape,
lower.tail = TRUE, log.p = FALSE)
qpoisinvgauss(p, mean, shape = 1, dispersion = 1/shape,
lower.tail = TRUE, log.p = FALSE)
rpoisinvgauss(n, mean, shape = 1, dispersion = 1/shape)
```

**Arguments**

x	vector of (positive integer) quantiles.
q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
mean, shape	parameters. Must be strictly positive. Infinite values are supported.
dispersion	an alternative way to specify the shape.
log, log.p	logical; if TRUE, probabilities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .

**Details**

The Poisson-inverse Gaussian distribution is the result of the continuous mixture between a Poisson distribution and an inverse Gaussian, that is, the distribution with probability mass function

$$p(x) = \int_0^{\infty} \frac{\lambda^x e^{-\lambda}}{x!} g(\lambda; \mu, \phi) d\lambda,$$

where  $g(\lambda; \mu, \phi)$  is the density function of the inverse Gaussian distribution with parameters mean =  $\mu$  and dispersion =  $\phi$  (see [dinvgauss](#)).

The resulting probability mass function is

$$p(x) = \sqrt{\frac{2}{\pi\phi}} \frac{e^{(\phi\mu)^{-1}}}{x!} \left( \sqrt{2\phi \left( 1 + \frac{1}{2\phi\mu^2} \right)} \right)^{-(x-\frac{1}{2})} K_{x-\frac{1}{2}} \left( \sqrt{\frac{2}{\phi} \left( 1 + \frac{1}{2\phi\mu^2} \right)} \right),$$

for  $x = 0, 1, \dots$ ,  $\mu > 0$ ,  $\phi > 0$  and where  $K_\nu(x)$  is the modified Bessel function of the third kind implemented by R's [besselK\(\)](#) and defined in its help.

The limiting case  $\mu = \infty$  has well defined probability mass and distribution functions, but has no finite strictly positive, integer moments. The pmf in this case reduces to

$$p(x) = \sqrt{\frac{2}{\pi\phi}} \frac{1}{x!} (\sqrt{2\phi})^{-(x-\frac{1}{2})} K_{x-\frac{1}{2}}(\sqrt{2/\phi}).$$

The limiting case  $\phi = 0$  is a degenerate distribution in  $x = 0$ .

If an element of  $x$  is not integer, the result of `dpoisinvgauss` is zero, with a warning.

The quantile is defined as the smallest value  $x$  such that  $F(x) \geq p$ , where  $F$  is the distribution function.

**Value**

`dpoisinvgauss` gives the probability mass function, `ppoisinvgauss` gives the distribution function, `qpoisinvgauss` gives the quantile function, and `rpoisinvgauss` generates random deviates.

Invalid arguments will result in return value `NaN`, with a warning.

The length of the result is determined by `n` for `rpoisinvgauss`, and is the maximum of the lengths of the numerical arguments for the other functions.

**Note**

[dpqr]pig are aliases for [dpqr]poisinvgauss.

qpoisinvgauss is based on qbinom et al.; it uses the Cornish–Fisher Expansion to include a skewness correction to a normal approximation, followed by a search.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>

**References**

Holla, M. S. (1966), “On a Poisson-Inverse Gaussian Distribution”, *Metrika*, vol. 15, p. 377-384.

Johnson, N. L., Kemp, A. W. and Kotz, S. (2005), *Univariate Discrete Distributions, Third Edition*, Wiley.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

Shaban, S. A., (1981) “Computation of the poisson-inverse gaussian distribution”, *Communications in Statistics - Theory and Methods*, vol. 10, no. 14, p. 1389-1399.

**See Also**

[dpois](#) for the Poisson distribution, [dinvgauss](#) for the inverse Gaussian distribution.

**Examples**

```
## Tables I and II of Shaban (1981)
x <- 0:2
sapply(c(0.4, 0.8, 1), dpoisinvgauss, x = x, mean = 0.1)
sapply(c(40, 80, 100, 130), dpoisinvgauss, x = x, mean = 1)

qpoisinvgauss(ppoisinvgauss(0:10, 1, dis = 2.5), 1, dis = 2.5)

x <- rpoisinvgauss(1000, 1, dis = 2.5)
y <- sort(unique(x))
plot(y, table(x)/length(x), type = "h", lwd = 2,
     pch = 19, col = "black", xlab = "x", ylab = "p(x)",
     main = "Empirical vs theoretical probabilities")
points(y, dpoisinvgauss(y, 1, dis = 2.5),
       pch = 19, col = "red")
legend("topright", c("empirical", "theoretical"),
      lty = c(1, NA), pch = c(NA, 19), col = c("black", "red"))
```

---

`quantile.aggregateDist`*Quantiles of Aggregate Claim Amount Distribution*

---

## Description

Quantile and Value-at-Risk methods for objects of class "aggregateDist".

## Usage

```
## S3 method for class 'aggregateDist'
quantile(x,
         probs = c(0.25, 0.5, 0.75, 0.9, 0.95, 0.975, 0.99, 0.995),
         smooth = FALSE, names = TRUE, ...)

## S3 method for class 'aggregateDist'
VaR(x, conf.level = c(0.9, 0.95, 0.99),
    smooth = FALSE, names = TRUE, ...)
```

## Arguments

<code>x</code>	an object of class "aggregateDist".
<code>probs, conf.level</code>	numeric vector of probabilities with values in $[0, 1)$ .
<code>smooth</code>	logical; when TRUE and <code>x</code> is a step function, quantiles are linearly interpolated between knots.
<code>names</code>	logical; if true, the result has a <code>names</code> attribute. Set to FALSE for speedup with many probs.
<code>...</code>	further arguments passed to or from other methods.

## Details

The quantiles are taken directly from the cumulative distribution function defined in `x`. Linear interpolation is available for step functions.

## Value

A numeric vector, named if `names` is TRUE.

## Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Louis-Philippe Pouliot

## See Also

[aggregateDist](#)

**Examples**

```

model.freq <- expression(data = rpois(3))
model.sev <- expression(data = rlnorm(10, 1.5))
Fs <- aggregateDist("simulation", model.freq, model.sev, nb.simul = 1000)
quantile(Fs, probs = c(0.25, 0.5, 0.75))
VaR(Fs)

```

---

quantile.grouped.data *Quantiles of Grouped Data*

---

**Description**

Sample quantiles corresponding to the given probabilities for objects of class "grouped.data".

**Usage**

```

## S3 method for class 'grouped.data'
quantile(x, probs = seq(0, 1, 0.25),
        names = TRUE, ...)

```

**Arguments**

x	an object of class "grouped.data".
probs	numeric vector of probabilities with values in [0, 1].
names	logical; if true, the result has a names attribute. Set to FALSE for speedup with many probs.
...	further arguments passed to or from other methods.

**Details**

The quantile function is the inverse of the ogive, that is a linear interpolation of the empirical quantile function.

The equation of the quantile function is

$$x = \frac{c_j(F_n(c_{j-1}) - q) + c_{j-1}(q - F_n(c_j))}{F_n(c_j) - F_n(c_{j-1})}$$

for  $0 \leq q \leq c_j$  and where  $c_0, \dots, c_r$  are the  $r + 1$  group boundaries and  $F_n$  is the empirical distribution function of the sample.

**Value**

A numeric vector, named if names is TRUE.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>

**See Also**

[ogive](#) for the smoothed empirical distribution of which `quantile.grouped.data` is an inverse; [grouped.data](#) to create grouped data objects.

**Examples**

```
data(gdental)
quantile(gdental)
Fn <- ogive(gdental)
Fn(quantile(gdental)) # inverse function
```

---

rcompound

*Simulation from Compound Models*


---

**Description**

`rcompound` generates random variates from a compound model.

`rcomppois` is a simplified version for a common case.

**Usage**

```
rcompound(n, model.freq, model.sev, SIMPLIFY = TRUE)
```

```
rcomppois(n, lambda, model.sev, SIMPLIFY = TRUE)
```

**Arguments**

<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>model.freq</code> , <code>model.sev</code>	expressions specifying the frequency and severity simulation models with the number of variates omitted (see details).
<code>lambda</code>	Poisson parameter.
<code>SIMPLIFY</code>	boolean; if <code>FALSE</code> the frequency and severity variates are returned along with the aggregate variates.

**Details**

`rcompound` generates variates from a random variable of the form

$$S = X_1 + \dots + X_N,$$

where  $N$  is the frequency random variable and  $X_1, X_2, \dots$  are the severity random variables. The latter are mutually independent, identically distributed and independent from  $N$ .

`model.freq` and `model.sev` specify the simulation models for the frequency and the severity random variables, respectively. A model is a complete call to a random number generation function,

with the number of variates omitted. This is similar to [rcomphierarc](#), but the calls need not be wrapped into [expression](#). Either argument may also be the name of an object containing an expression, in which case the object will be evaluated in the evaluation frame to retrieve the expression. The argument of the random number generation functions for the number of variates to simulate **must** be named `n`.

`rcomppois` generates variates from the common Compound Poisson model, that is when random variable  $N$  is Poisson distributed with mean  $\lambda$ .

### Value

When `SIMPLIFY = TRUE`, a vector of aggregate amounts  $S_1, \dots, S_n$ .

When `SIMPLIFY = FALSE`, a list of three elements:

aggregate	vector of aggregate amounts $S_1, \dots, S_n$ ;
frequency	vector of frequencies $N_1, \dots, N_n$ ;
severity	vector of severities $X_1, X_2, \dots$

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>

### See Also

[rcomphierarc](#) to simulate from compound hierarchical models.

### Examples

```
## Compound Poisson model with gamma severity.
rcompound(10, rpois(2), rgamma(2, 3))
rcomppois(10, 2, rgamma(2, 3))      # same

## Frequencies and individual claim amounts along with aggregate
## values.
rcomppois(10, 2, rgamma(2, 3), SIMPLIFY = FALSE)

## Wrapping the simulation models into expression() is allowed, but
## not needed.
rcompound(10, expression(rpois(2)), expression(rgamma(2, 3)))

## Not run: ## Speed comparison between rcompound() and rcomphierarc().
## [Also note the simpler syntax for rcompound().]
system.time(rcompound(1e6, rpois(2), rgamma(2, 3)))
system.time(rcomphierarc(1e6, expression(rpois(2)), expression(rgamma(2, 3))))
## End(Not run)
## The severity can itself be a compound model. It makes sense
## in such a case to use a zero-truncated frequency distribution
## for the second level model.
rcomppois(10, 2,
           rcompound(rztnbinom(1.5, 0.7), rlnorm(1.2, 1)))
```



---

 rmixture

*Simulation from Discrete Mixtures*


---

### Description

Generate random variates from a discrete mixture of distributions.

### Usage

```
rmixture(n, probs, models, shuffle = TRUE)
```

### Arguments

n	number of random variates to generate. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
probs	numeric non-negative vector specifying the probability for each model; is internally normalized to sum 1. Infinite and missing values are not allowed. Values are recycled as necessary to match the length of models.
models	vector of expressions specifying the simulation models with the number of variates omitted (see details). Models are recycled as necessary to match the length of probs.
shuffle	logical; should the random variates from the distributions be shuffled?

### Details

`rmixture` generates variates from a discrete mixture, that is random variable with a probability density function of the form

$$f(x) = p_1 f_1(x) + \dots + p_n f_n(x),$$

where  $f_1, \dots, f_n$  are densities and  $\sum_{i=1}^n p_i = 1$ .

The values in `probs` will be internally normalized to be used as probabilities  $p_1 + \dots + p_n$ .

The specification of simulation models uses the syntax of `rcomphierarc`. Models  $f_1, \dots, f_n$  are expressed in a semi-symbolic fashion using an object of mode `expression` where each element is a complete call to a random number generation function, with the number of variates omitted.

The argument of the random number generation functions for the number of variates to simulate **must** be named `n`.

If `shuffle` is `FALSE`, the output vector contains all the random variates from the first model, then all the random variates from the second model, and so on. If the order of the variates is irrelevant, this cuts the time to generate the variates roughly in half.

### Value

A vector of random variates from the mixture with density  $f(x)$ .

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>

**See Also**

[rcompound](#) to simulate from compound models.

[rcomphierarc](#) to simulate from compound hierarchical models.

**Examples**

```
## Mixture of two exponentials (with means 1/3 and 1/7) with equal
## probabilities.
rmixture(10, 0.5, expression(rexp(3), rexp(7)))
rmixture(10, 42, expression(rexp(3), rexp(7))) # same

## Mixture of two lognormals with different probabilities.
rmixture(10, probs = c(0.55, 0.45),
        models = expression(rlnorm(3.6, 0.6),
                             rlnorm(4.6, 0.3)))
```

---

ruin

*Probability of Ruin*

---

**Description**

Calculation of infinite time probability of ruin in the models of Cramér-Lundberg and Sparre Andersen, that is with exponential or phase-type (including mixtures of exponentials, Erlang and mixture of Erlang) claims interarrival time.

**Usage**

```
ruin(claims = c("exponential", "Erlang", "phase-type"), par.claims,
     wait = c("exponential", "Erlang", "phase-type"), par.wait,
     premium.rate = 1, tol = sqrt(.Machine$double.eps),
     maxit = 200L, echo = FALSE)

## S3 method for class 'ruin'
plot(x, from = NULL, to = NULL, add = FALSE,
     xlab = "u", ylab = expression(psi(u)),
     main = "Probability of Ruin", xlim = NULL, ...)
```

**Arguments**

`claims` character; the type of claim severity distribution.

`wait` character; the type of claim interarrival (wait) time distribution.

`par.claims`, `par.wait` named list containing the parameters of the distribution (see details).

premium.rate	numeric vector of length 1; the premium rate.
tol, maxit, echo	respectively the tolerance level of the stopping criteria, the maximum number of iterations and whether or not to echo the procedure when the transition rates matrix is determined iteratively. Ignored if wait = "exponential".
x	an object of class "ruin".
from, to	the range over which the function will be plotted.
add	logical; if TRUE add to already existing plot.
xlim	numeric of length 2; if specified, it serves as default for c(from, to).
xlab, ylab	label of the x and y axes, respectively.
main	main title.
...	further graphical parameters accepted by <a href="#">curve</a> .

### Details

The names of the parameters in `par.claims` and `par.wait` must be the same as in [dexp](#), [dgamma](#) or [dphype](#), as appropriate. A model will be a mixture of exponential or Erlang distributions (but not phase-type) when the parameters are vectors of length  $> 1$  and the parameter list contains a vector weights of the coefficients of the mixture.

Parameters are recycled when needed. Their names can be abbreviated.

Combinations of exponentials as defined in Dufresne and Gerber (1988) are *not* supported.

Ruin probabilities are evaluated using [pphype](#) except when both distributions are exponential, in which case an explicit formula is used.

When `wait != "exponential"` (Sparre Andersen model), the transition rate matrix  $Q$  of the distribution of the probability of ruin is determined iteratively using a fixed point-like algorithm. The stopping criteria used is

$$\max \left\{ \sum_{j=1}^n |Q_{ij} - Q'_{ij}| \right\} < \text{tol},$$

where  $Q$  and  $Q'$  are two successive values of the matrix.

### Value

A function of class "ruin" inheriting from the "function" class to compute the probability of ruin given initial surplus levels. The function has arguments:

u	numeric vector of initial surplus levels;
survival	logical; if FALSE (default), probabilities are $\psi(u)$ , otherwise, $\phi(u) = 1 - \psi(u)$ ;
lower.tail	an alias for !survival.

### Author(s)

Vincent Goulet <[vincent.goulet@act.ulaval.ca](mailto:vincent.goulet@act.ulaval.ca)>, and Christophe Dutang

## References

- Asmussen, S. and Rolski, T. (1991), Computational methods in risk theory: A matrix algorithmic approach, *Insurance: Mathematics and Economics* **10**, 259–274.
- Dufresne, F. and Gerber, H. U. (1988), Three methods to calculate the probability of ruin, *Astin Bulletin* **19**, 71–90.
- Gerber, H. U. (1979), *An Introduction to Mathematical Risk Theory*, Huebner Foundation.

## Examples

```
## Case with an explicit formula: exponential claims and exponential
## interarrival times.
psi <- ruin(claims = "e", par.claims = list(rate = 5),
           wait = "e", par.wait = list(rate = 3))

psi
psi(0:10)
plot(psi, from = 0, to = 10)

## Mixture of two exponentials for claims, exponential interarrival
## times (Gerber 1979)
psi <- ruin(claims = "e", par.claims = list(rate = c(3, 7), w = 0.5),
           wait = "e", par.wait = list(rate = 3), pre = 1)
u <- 0:10
psi(u)
(24 * exp(-u) + exp(-6 * u))/35 # same

## Phase-type claims, exponential interarrival times (Asmussen and
## Rolski 1991)
p <- c(0.5614, 0.4386)
r <- matrix(c(-8.64, 0.101, 1.997, -1.095), 2, 2)
lambda <- 1/(1.1 * mphtype(1, p, r))
psi <- ruin(claims = "p", par.claims = list(prob = p, rates = r),
           wait = "e", par.wait = list(rate = lambda))

psi
plot(psi, xlim = c(0, 50))

## Phase-type claims, mixture of two exponentials for interarrival times
## (Asmussen and Rolski 1991)
a <- (0.4/5 + 0.6) * lambda
ruin(claims = "p", par.claims = list(prob = p, rates = r),
     wait = "e", par.wait = list(rate = c(5 * a, a), weights =
                                   c(0.4, 0.6)),
     maxit = 225L)
```

---

severity

*Manipulation of Individual Claim Amounts*

---

## Description

severity is a generic function created to manipulate individual claim amounts. The function invokes particular *methods* which depend on the `class` of the first argument.

**Usage**

```
severity(x, ...)  
  
## Default S3 method:  
severity(x, bycol = FALSE, drop = TRUE, ...)
```

**Arguments**

x	an R object.
bycol	logical; whether to “unroll” horizontally (FALSE) or vertically (TRUE)
...	further arguments to be passed to or from other methods.
drop	logical; if TRUE, the result is coerced to the lowest possible dimension.

**Details**

Currently, the default method is equivalent to [unroll](#). This is liable to change since the link between the name and the use of the function is rather weak.

**Value**

A vector or matrix.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Louis-Philippe Pouliot

**See Also**

[severity.portfolio](#) for the original motivation of these functions.

**Examples**

```
x <- list(c(1:3), c(1:8), c(1:4), c(1:3))  
(mat <- matrix(x, 2, 2))  
severity(mat)  
severity(mat, bycol = TRUE)
```

---

simul

*Simulation from Compound Hierarchical Models*

---

**Description**

Simulate data for insurance applications allowing hierarchical structures and separate models for the frequency and severity of claims distributions.

rcomphierarc is an alias for simul.

**Usage**

```

simul(nodes, model.freq = NULL, model.sev = NULL, weights = NULL)

rcomphierarc(nodes, model.freq = NULL, model.sev = NULL, weights = NULL)

## S3 method for class 'portfolio'
print(x, ...)

```

**Arguments**

<code>nodes</code>	a vector or a named list giving the number of "nodes" at each level in the hierarchy of the portfolio. The nodes are listed from top (portfolio) to bottom (usually the years of experience).
<code>model.freq</code>	a named vector of expressions specifying the frequency of claims model (see details); if NULL, only claim amounts are simulated.
<code>model.sev</code>	a named vector of expressions specifying the severity of claims model (see details); if NULL, only claim numbers are simulated.
<code>weights</code>	a vector of weights.
<code>x</code>	a portfolio object.
<code>...</code>	potential further arguments required by generic.

**Details**

The order and the names of the elements in `nodes`, `model.freq` and `model.sev` must match. At least one of `model.freq` and `model.sev` must be non NULL.

`nodes` may be a basic vector, named or not, for non hierarchical models. The rule above still applies, so `model.freq` and `model.sev` should not be named if `nodes` is not. However, for non hierarchical models, [rcompound](#) is faster and has a simpler interface.

`nodes` specifies the hierarchical layout of the portfolio. Each element of the list is a vector of the number of nodes at a given level. Vectors are recycled as necessary.

`model.freq` and `model.sev` specify the simulation models for claim numbers and claim amounts, respectively. A model is expressed in a semi-symbolic fashion using an object of mode [expression](#). Each element of the object must be named and should be a complete call to a random number generation function, with the number of variates omitted. Hierarchical (or mixtures of) models are achieved by replacing one or more parameters of a distribution at a given level by any combination of the names of the levels above. If no mixing is to take place at a level, the model for this level can be NULL.

The argument of the random number generation functions for the number of variates to simulate **must** be named `n`.

Weights will be used wherever the name "weights" appears in a model. It is the user's responsibility to ensure that the length of `weights` will match the number of nodes when weights are to be used. Normally, there should be one weight per node at the lowest level of the model.

Data is generated in lexicographic order, that is by row in the output matrix.

**Value**

An object of class "portfolio". A print method for this class displays the models used in the simulation as well as the frequency of claims for each year and entity in the portfolio.

An object of class "portfolio" is a list containing the following components:

data	a two dimension list where each element is a vector of claim amounts;
weights	the vector of weights given in argument reshaped as a matrix matching element data, or NULL;
classification	a matrix of integers where each row is a unique set of subscripts identifying an entity in the portfolio (e.g. integers $i$ , $j$ and $k$ for data $X_{ijkt}$ );
nodes	the nodes argument, appropriately recycled;
model.freq	the frequency model as given in argument;
model.sev	the severity model as given in argument.

It is recommended to manipulate objects of class "portfolio" by means of the corresponding methods of functions aggregate, frequency and severity.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>, Sébastien Auclair and Louis-Philippe Pouliot

**References**

Goulet, V. and Pouliot, L.-P. (2008), Simulation of compound hierarchical models in R, *North American Actuarial Journal* **12**, 401–412.

**See Also**

[simul.summaries](#) for the functions to create the matrices of aggregate claim amounts, frequencies and individual claim amounts.

[rcompound](#) for a simpler and much faster way to generate variates from standard, non hierarchical, compound models.

**Examples**

```
## Two level (contracts and years) portfolio with frequency model
## Nit|Theta_i ~ Poisson(Theta_i), Theta_i ~ Gamma(2, 3) and severity
## model X ~ Lognormal(5, 1)
simul(nodes = list(contract = 10, year = 5),
      model.freq = expression(contract = rgamma(2, 3),
                             year = rpois(contract)),
      model.sev = expression(contract = NULL,
                             year = rlnorm(5, 1)))

## Model with weights and mixtures for both frequency and severity
## models
nodes <- list(entity = 8, year = c(5, 4, 4, 5, 3, 5, 4, 5))
mf <- expression(entity = rgamma(2, 3),
```

```

        year = rpois(weights * entity))
ms <- expression(entity = rnorm(5, 1),
                 year = rlnorm(entity, 1))
wit <- sample(2:10, 35, replace = TRUE)
pf <- simul(nodes, mf, ms, wit)
pf # print method
weights(pf) # extraction of weights
aggregate(pf)[, -1]/weights(pf)[, -1] # ratios

## Four level hierarchical model for frequency only
nodes <- list(sector = 3, unit = c(3, 4),
             employer = c(3, 4, 3, 4, 2, 3, 4), year = 5)
mf <- expression(sector = rexp(1),
                 unit = rexp(sector),
                 employer = rgamma(unit, 1),
                 year = rpois(employer))
pf <- simul(nodes, mf, NULL)
pf # print method
aggregate(pf) # aggregate claim amounts
frequency(pf) # frequencies
severity(pf) # individual claim amounts

## Standard, non hierarchical, compound model with simplified
## syntax (function rcompound() is much faster for such cases)
simul(10,
      model.freq = expression(rpois(2)),
      model.sev = expression(rgamma(2, 3)))

```

---

simul.summaries

*Summary Statistics of a Portfolio*


---

## Description

Methods for [class](#) "portfolio" objects.

aggregate splits portfolio data into subsets and computes summary statistics for each.

frequency computes the frequency of claims for subsets of portfolio data.

severity extracts the individual claim amounts.

weights extracts the matrix of weights.

## Usage

```

## S3 method for class 'portfolio'
aggregate(x, by = names(x$nodes), FUN = sum,
         classification = TRUE, prefix = NULL, ...)

```

```

## S3 method for class 'portfolio'
frequency(x, by = names(x$nodes),
         classification = TRUE, prefix = NULL, ...)

```



```
## S3 method for class 'portfolio'
severity(x, by = head(names(x$node), -1), splitcol = NULL,
         classification = TRUE, prefix = NULL, ...)

## S3 method for class 'portfolio'
weights(object, classification = TRUE, prefix = NULL, ...)
```

### Arguments

<code>x</code> , object	an object of class "portfolio", typically created with <code>simul</code> .
<code>by</code>	character vector of grouping elements using the level names of the portfolio in <code>x</code> . The names can be abbreviated.
<code>FUN</code>	the function to be applied to data subsets.
<code>classification</code>	boolean; if TRUE, the node identifier columns are included in the output.
<code>prefix</code>	characters to prefix column names with; if NULL, sensible defaults are used when appropriate.
<code>splitcol</code>	columns of the data matrix to extract separately; usual matrix indexing methods are supported.
<code>...</code>	optional arguments to FUN, or passed to or from other methods.

### Details

By default, `aggregate.portfolio` computes the aggregate claim amounts for the grouping specified in `by`. Any other statistic based on the individual claim amounts can be used through argument `FUN`.

`frequency.portfolio` is equivalent to using `aggregate.portfolio` with argument `FUN` equal to `if (identical(x,NA)) NA else length(x)`.

`severity.portfolio` extracts individual claim amounts of a portfolio by groupings using the default method of `severity`. Argument `splitcol` allows to get the individual claim amounts of specific columns separately.

`weights.portfolio` extracts the weight matrix of a portfolio.

### Value

A matrix or vector depending on the groupings specified in `by`.

For the `aggregate` and `frequency` methods: if at least one level other than the last one is used for grouping, the result is a matrix obtained by binding the appropriate node identifiers extracted from `x$classification` if `classification = TRUE`, and the summaries per grouping. If the last level is used for grouping, the column names of `x$data` are retained; if the last level is not used for grouping, the column name is replaced by the deparsed name of `FUN`. If only the last level is used (column summaries), a named vector is returned.

For the `severity` method: a list of two elements:

<code>main</code>	NULL or a matrix of claim amounts for the columns not specified in <code>splitcol</code> , with the appropriate node identifiers extracted from <code>x\$classification</code> if <code>classification = TRUE</code> ;
-------------------	--

`split` same as above, but for the columns specified in `splitcol`.

For the `weights` method: the weight matrix of the portfolio with node identifiers if `classification = TRUE`.

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>, Louis-Philippe Pouliot.

### See Also

[simul](#)

### Examples

```
nodes <- list(sector = 3, unit = c(3, 4),
             employer = c(3, 4, 3, 4, 2, 3, 4), year = 5)
model.freq <- expression(sector = rexp(1),
                        unit = rexp(sector),
                        employer = rgamma(unit, 1),
                        year = rpois(employer))
model.sev <- expression(sector = rnorm(6, 0.1),
                       unit = rnorm(sector, 1),
                       employer = rnorm(unit, 1),
                       year = rlnorm(employer, 1))
pf <- simul(nodes, model.freq, model.sev)

aggregate(pf)           # aggregate claim amount by employer and year
aggregate(pf, classification = FALSE) # same, without node identifiers
aggregate(pf, by = "sector")      # by sector
aggregate(pf, by = "y")          # by year
aggregate(pf, by = c("s", "u"), mean) # average claim amount

frequency(pf)          # number of claims
frequency(pf, prefix = "freq.") # more explicit column names

severity(pf)           # claim amounts by row
severity(pf, by = "year") # by column
severity(pf, by = c("s", "u")) # by unit
severity(pf, splitcol = "year.5") # last year separate
severity(pf, splitcol = 5) # same
severity(pf, splitcol = c(FALSE, FALSE, FALSE, FALSE, TRUE)) # same

weights(pf)

## For portfolios with weights, the following computes loss ratios.
## Not run: aggregate(pf, classif = FALSE) / weights(pf, classif = FALSE)
```

---

 SingleParameterPareto *The Single-parameter Pareto Distribution*


---

**Description**

Density function, distribution function, quantile function, random generation, raw moments, and limited moments for the Single-parameter Pareto distribution with parameter shape.

**Usage**

```
dpareto1(x, shape, min, log = FALSE)
ppareto1(q, shape, min, lower.tail = TRUE, log.p = FALSE)
qpareto1(p, shape, min, lower.tail = TRUE, log.p = FALSE)
rpareto1(n, shape, min)
mpareto1(order, shape, min)
levpareto1(limit, shape, min, order = 1)
```

**Arguments**

x, q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If length(n) > 1, the length is taken to be the number required.
shape	parameter. Must be strictly positive.
min	lower bound of the support of the distribution.
log, log.p	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
order	order of the moment.
limit	limit of the loss variable.

**Details**

The single-parameter Pareto, or Pareto I, distribution with parameter shape =  $\alpha$  has density:

$$f(x) = \frac{\alpha\theta^\alpha}{x^{\alpha+1}}$$

for  $x > \theta$ ,  $\alpha > 0$  and  $\theta > 0$ .

Although there appears to be two parameters, only shape is a true parameter. The value of  $\min = \theta$  must be set in advance.

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $k < \alpha$  and the  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ ,  $x \geq \theta$ .

**Value**

`dpareto1` gives the density, `ppareto1` gives the distribution function, `qpareto1` gives the quantile function, `rpareto1` generates random deviates, `mpareto1` gives the  $k$ th raw moment, and `levpareto1` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

**Note**

For Pareto distributions, we use the classification of Arnold (2015) with the parametrization of Klugman et al. (2012).

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Arnold, B.C. (2015), *Pareto Distributions*, Second Edition, CRC Press.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**See Also**

[dpareto](#) for the two-parameter Pareto distribution.

**Examples**

```
exp(dpareto1(5, 3, 4, log = TRUE))
p <- (1:10)/10
ppareto1(qpareto1(p, 2, 3), 2, 3)
mpareto1(2, 3, 4) - mpareto(1, 3, 4) ^ 2
levpareto(10, 3, 4, order = 2)
```

**Description**

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Transformed Beta distribution with parameters `shape1`, `shape2`, `shape3` and `scale`.

**Usage**

```
dtrbeta(x, shape1, shape2, shape3, rate = 1, scale = 1/rate,
        log = FALSE)
ptrbeta(q, shape1, shape2, shape3, rate = 1, scale = 1/rate,
        lower.tail = TRUE, log.p = FALSE)
qtrbeta(p, shape1, shape2, shape3, rate = 1, scale = 1/rate,
        lower.tail = TRUE, log.p = FALSE)
rtrbeta(n, shape1, shape2, shape3, rate = 1, scale = 1/rate)
mtrbeta(order, shape1, shape2, shape3, rate = 1, scale = 1/rate)
levtrbeta(limit, shape1, shape2, shape3, rate = 1, scale = 1/rate,
          order = 1)
```

**Arguments**

`x`, `q`            vector of quantiles.

`p`                    vector of probabilities.

`n`                    number of observations. If `length(n) > 1`, the length is taken to be the number required.

`shape1`, `shape2`, `shape3`, `scale`            parameters. Must be strictly positive.

`rate`                an alternative way to specify the scale.

`log`, `log.p`        logical; if TRUE, probabilities/densities  $p$  are returned as  $\log(p)$ .

`lower.tail`        logical; if TRUE (default), probabilities are  $P[X \leq x]$ , otherwise,  $P[X > x]$ .

`order`                order of the moment.

`limit`                limit of the loss variable.

**Details**

The transformed beta distribution with parameters  $\text{shape1} = \alpha$ ,  $\text{shape2} = \gamma$ ,  $\text{shape3} = \tau$  and  $\text{scale} = \theta$ , has density:

$$f(x) = \frac{\Gamma(\alpha + \tau)}{\Gamma(\alpha)\Gamma(\tau)} \frac{\gamma(x/\theta)^{\gamma\tau}}{x[1 + (x/\theta)^\gamma]^{\alpha+\tau}}$$

for  $x > 0$ ,  $\alpha > 0$ ,  $\gamma > 0$ ,  $\tau > 0$  and  $\theta > 0$ . (Here  $\Gamma(\alpha)$  is the function implemented by R's [gamma\(\)](#) and defined in its help.)

The transformed beta is the distribution of the random variable

$$\theta \left( \frac{X}{1-X} \right)^{1/\gamma},$$

where  $X$  has a beta distribution with parameters  $\tau$  and  $\alpha$ .

The transformed beta distribution defines a family of distributions with the following special cases:

- A [Burr](#) distribution when  $\text{shape3} == 1$ ;
- A [loglogistic](#) distribution when  $\text{shape1} == \text{shape3} == 1$ ;
- A [paralogistic](#) distribution when  $\text{shape3} == 1$  and  $\text{shape2} == \text{shape1}$ ;

- A [generalized Pareto](#) distribution when `shape2 == 1`;
- A [Pareto](#) distribution when `shape2 == shape3 == 1`;
- An [inverse Burr](#) distribution when `shape1 == 1`;
- An [inverse Pareto](#) distribution when `shape2 == shape1 == 1`;
- An [inverse paralogistic](#) distribution when `shape1 == 1` and `shape3 == shape2`.

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ ,  $-\tau\gamma < k < \alpha\gamma$ .

The  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ ,  $k > -\tau\gamma$  and  $\alpha - k/\gamma$  not a negative integer.

### Value

`dtrbeta` gives the density, `ptrbeta` gives the distribution function, `qtrbeta` gives the quantile function, `rtrbeta` generates random deviates, `mtrbeta` gives the  $k$ th raw moment, and `levtrbeta` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

### Note

`levtrbeta` computes the limited expected value using [betaint](#).

Distribution also known as the Generalized Beta of the Second Kind and Pearson Type VI. See also Kleiber and Kotz (2003) for alternative names and parametrizations.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

### References

Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

### See Also

[dfpareto](#) for an equivalent distribution with a location parameter.

### Examples

```
exp(dtrbeta(2, 2, 3, 4, 5, log = TRUE))
p <- (1:10)/10
ptrbeta(qtrbeta(p, 2, 3, 4, 5), 2, 3, 4, 5)
qpearson6(0.3, 2, 3, 4, 5, lower.tail = FALSE)

## variance
```

```

mtrbeta(2, 2, 3, 4, 5) - mtrbeta(1, 2, 3, 4, 5)^2

## case with shape1 - order/shape2 > 0
levtrbeta(10, 2, 3, 4, scale = 1, order = 2)

## case with shape1 - order/shape2 < 0
levtrbeta(10, 1/3, 0.75, 4, scale = 0.5, order = 2)

```

---

TransformedGamma

*The Transformed Gamma Distribution*


---

### Description

Density function, distribution function, quantile function, random generation, raw moments and limited moments for the Transformed Gamma distribution with parameters shape1, shape2 and scale.

### Usage

```

dtrgamma(x, shape1, shape2, rate = 1, scale = 1/rate,
         log = FALSE)
ptrgamma(q, shape1, shape2, rate = 1, scale = 1/rate,
         lower.tail = TRUE, log.p = FALSE)
qtrgamma(p, shape1, shape2, rate = 1, scale = 1/rate,
         lower.tail = TRUE, log.p = FALSE)
rtrgamma(n, shape1, shape2, rate = 1, scale = 1/rate)
mtrgamma(order, shape1, shape2, rate = 1, scale = 1/rate)
levtrgamma(limit, shape1, shape2, rate = 1, scale = 1/rate,
           order = 1)

```

### Arguments

<code>x, q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>shape1, shape2, scale</code>	parameters. Must be strictly positive.
<code>rate</code>	an alternative way to specify the scale.
<code>log, log.p</code>	logical; if TRUE, probabilities/densities $p$ are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
<code>order</code>	order of the moment.
<code>limit</code>	limit of the loss variable.

### Details

The transformed gamma distribution with parameters  $\text{shape1} = \alpha$ ,  $\text{shape2} = \tau$  and  $\text{scale} = \theta$  has density:

$$f(x) = \frac{\tau u^\alpha e^{-u}}{x \Gamma(\alpha)}, \quad u = (x/\theta)^\tau$$

for  $x > 0$ ,  $\alpha > 0$ ,  $\tau > 0$  and  $\theta > 0$ . (Here  $\Gamma(\alpha)$  is the function implemented by R's `gamma()` and defined in its help.)

The transformed gamma is the distribution of the random variable  $\theta X^{1/\tau}$ , where  $X$  has a gamma distribution with shape parameter  $\alpha$  and scale parameter 1 or, equivalently, of the random variable  $Y^{1/\tau}$  with  $Y$  a gamma distribution with shape parameter  $\alpha$  and scale parameter  $\theta^\tau$ .

The transformed gamma probability distribution defines a family of distributions with the following special cases:

- A **Gamma** distribution when  $\text{shape2} == 1$ ;
- A **Weibull** distribution when  $\text{shape1} == 1$ ;
- An **Exponential** distribution when  $\text{shape2} == \text{shape1} == 1$ .

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$  and the  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ ,  $k > -\alpha\tau$ .

### Value

`dtrgamma` gives the density, `ptrgamma` gives the distribution function, `qtrgamma` gives the quantile function, `rtrgamma` generates random deviates, `mtrgamma` gives the  $k$ th raw moment, and `levtrgamma` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

### Note

Distribution also known as the Generalized Gamma. See also Kleiber and Kotz (2003) for alternative names and parametrizations.

The "distributions" package vignette provides the interrelations between the continuous size distributions in **actuar** and the complete formulas underlying the above functions.

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

### References

- Kleiber, C. and Kotz, S. (2003), *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley.
- Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.



**Examples**

```
exp(dtrgamma(2, 3, 4, 5, log = TRUE))
p <- (1:10)/10
ptrgamma(qtrgamma(p, 2, 3, 4), 2, 3, 4)
mtrgamma(2, 3, 4, 5) - mtrgamma(1, 3, 4, 5) ^ 2
levtrgamma(10, 3, 4, 5, order = 2)
```

UniformSupp

---

*Moments and Moment Generating Function of the Uniform Distribution*

---

**Description**

Raw moments, limited moments and moment generating function for the Uniform distribution from min to max.

**Usage**

```
munif(order, min = 0, max = 1)
levunif(limit, min = 0, max = 1, order = 1)
mgfunif(t, min = 0, max = 1, log = FALSE)
```

**Arguments**

order	order of the moment.
min, max	lower and upper limits of the distribution. Must be finite.
limit	limit of the random variable.
t	numeric vector.
log	logical; if TRUE, the cumulant generating function is returned.

**Details**

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$ , the  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$  and the moment generating function is  $E[e^{tX}]$ .

**Value**

munif gives the  $k$ th raw moment, levunif gives the  $k$ th moment of the limited random variable, and mgfunif gives the moment generating function in t.

Invalid arguments will result in return value NaN, with a warning.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>, Christophe Dutang

## References

[https://en.wikipedia.org/wiki/Uniform\\_distribution\\_%28continuous%29](https://en.wikipedia.org/wiki/Uniform_distribution_%28continuous%29)

## See Also

[Uniform](#).

## Examples

```
munif(-1)
munif(1:5)
levunif(3, order=1:5)
levunif(3, 2, 4)
mgfunif(1, 1, 2)
```

---

unroll

*Display a Two-Dimension Version of a Matrix of Vectors*

---

## Description

Displays all values of a matrix of vectors by “unrolling” the object vertically or horizontally.

## Usage

```
unroll(x, bycol = FALSE, drop = TRUE)
```

## Arguments

x	a list of vectors with a <code>dim</code> attribute of length 0, 1 or 2.
bycol	logical; whether to unroll horizontally (FALSE) or vertically (TRUE).
drop	logical; if TRUE, the result is coerced to the lowest possible dimension.

## Details

`unroll` returns a matrix where elements of `x` are concatenated (“unrolled”) by row (`bycol = FALSE`) or by column (`bycol = TRUE`). `NA` is used to make rows/columns of equal length.

Vectors and one dimensional arrays are coerced to **row** matrices.

## Value

A vector or matrix.

## Author(s)

Vincent Goulet <[vincent.goulet@act.ulaval.ca](mailto:vincent.goulet@act.ulaval.ca)> and Louis-Philippe Pouliot

**See Also**

This function was originally written for use in [severity.portfolio](#).

**Examples**

```
x <- list(c(1:3), c(1:8), c(1:4), c(1:3))
(mat <- matrix(x, 2, 2))
```

```
unroll(mat)
unroll(mat, bycol = TRUE)
```

```
unroll(mat[1, ])
unroll(mat[1, ], drop = FALSE)
```

---

VaR

*Value at Risk*

---

**Description**

Value at Risk.

**Usage**

```
VaR(x, ...)
```

**Arguments**

`x` an R object.  
`...` further arguments passed to or from other methods.

**Details**

This is a generic function with, currently, only a method for objects of class "aggregateDist".

**Value**

An object of class numeric.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Tommy Ouellet

**See Also**

[VaR.aggregateDist](#), [aggregateDist](#)

**Description**

Raw moments and limited moments for the Weibull distribution with parameters shape and scale.

**Usage**

```
mweibull(order, shape, scale = 1)
levweibull(limit, shape, scale = 1, order = 1)
```

**Arguments**

order	order of the moment.
limit	limit of the loss variable.
shape, scale	shape and scale parameters, the latter defaulting to 1.

**Details**

The  $k$ th raw moment of the random variable  $X$  is  $E[X^k]$  and the  $k$ th limited moment at some limit  $d$  is  $E[\min(X, d)^k]$ ,  $k > -\tau$ .

**Value**

`mweibull` gives the  $k$ th raw moment and `levweibull` gives the  $k$ th moment of the limited loss variable.

Invalid arguments will result in return value NaN, with a warning.

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca> and Mathieu Pigeon

**References**

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**See Also**

[Weibull](#) for details on the Weibull distribution and functions `[dpqr]weibull`.

**Examples**

```
mweibull(2, 3, 4) - mweibull(1, 3, 4)^2
levweibull(10, 3, 4, order = 2)
```

**Description**

Density function, distribution function, quantile function and random generation for the Zero-Modified Binomial distribution with parameters size and prob, and probability at zero  $p_0$ .

**Usage**

```
dzmbinom(x, size, prob, p0, log = FALSE)
pzmbinom(q, size, prob, p0, lower.tail = TRUE, log.p = FALSE)
qzmbinom(p, size, prob, p0, lower.tail = TRUE, log.p = FALSE)
rzmbinom(n, size, prob, p0)
```

**Arguments**

x	vector of (strictly positive integer) quantiles.
q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If length(n) > 1, the length is taken to be the number required.
size	number of trials (strictly positive integer).
prob	probability of success on each trial. $0 \leq \text{prob} \leq 1$ .
p0	probability mass at zero. $0 \leq p_0 \leq 1$ .
log, log.p	logical; if TRUE, probabilities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .

**Details**

The zero-modified binomial distribution with size =  $n$ , prob =  $p$  and  $p_0 = p_0$  is a discrete mixture between a degenerate distribution at zero and a (standard) binomial. The probability mass function is  $p(0) = p_0$  and

$$p(x) = \frac{(1 - p_0)}{(1 - (1 - p)^n)} f(x)$$

for  $x = 1, \dots, n$ ,  $0 < p \leq 1$  and  $0 \leq p_0 \leq 1$ , where  $f(x)$  is the probability mass function of the binomial. The cumulative distribution function is

$$P(x) = p_0 + (1 - p_0) \left( \frac{F(x) - F(0)}{1 - F(0)} \right)$$

The mean is  $(1 - p_0)\mu$  and the variance is  $(1 - p_0)\sigma^2 + p_0(1 - p_0)\mu^2$ , where  $\mu$  and  $\sigma^2$  are the mean and variance of the zero-truncated binomial.

In the terminology of Klugman et al. (2012), the zero-modified binomial is a member of the  $(a, b, 1)$  class of distributions with  $a = -p/(1 - p)$  and  $b = (n + 1)p/(1 - p)$ .

The special case  $p_0 = 0$  is the zero-truncated binomial.

If an element of  $x$  is not integer, the result of `dzmbinom` is zero, with a warning.

The quantile is defined as the smallest value  $x$  such that  $P(x) \geq p$ , where  $P$  is the distribution function.

### Value

`dzmbinom` gives the probability mass function, `pzmbinom` gives the distribution function, `qzmbinom` gives the quantile function, and `rzmbinom` generates random deviates.

Invalid `size`, `prob` or `p0` will result in return value `NaN`, with a warning.

The length of the result is determined by `n` for `rzmbinom`, and is the maximum of the lengths of the numerical arguments for the other functions.

### Note

Functions `{d,p,q}zmbinom` use `{d,p,q}binom` for all but the trivial input values and  $p(0)$ .

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>

### References

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

### See Also

[dbinom](#) for the binomial distribution.

[dztbinom](#) for the zero-truncated binomial distribution.

### Examples

```
dzmbinom(1:5, size = 5, prob = 0.4, p0 = 0.2)
(1-0.2) * dbinom(1:5, 5, 0.4)/pbinom(0, 5, 0.4, lower = FALSE) # same

## simple relation between survival functions
pzmbinom(0:5, 5, 0.4, p0 = 0.2, lower = FALSE)
(1-0.2) * pbinom(0:5, 5, 0.4, lower = FALSE) /
  pbinom(0, 5, 0.4, lower = FALSE) # same

qzmbinom(pzmbinom(1:10, 10, 0.6, p0 = 0.1), 10, 0.6, p0 = 0.1)

n <- 8; p <- 0.3; p0 <- 0.025
x <- 0:n
title <- paste("ZM Binomial(", n, ", ", " ", p, ", ", p0 = " ", p0,
              ") and Binomial(", n, ", ", " ", p, ") PDF",
              sep = " ")
plot(x, dzmbinom(x, n, p, p0), type = "h", lwd = 2, ylab = "p(x)",
     main = title)
```

```
points(x, dbinom(x, n, p), pch = 19, col = "red")
legend("topright", c("ZT binomial probabilities", "Binomial probabilities"),
      col = c("black", "red"), lty = c(1, 0), lwd = 2, pch = c(NA, 19))
```

ZeroModifiedGeometric *The Zero-Modified Geometric Distribution*

## Description

Density function, distribution function, quantile function and random generation for the Zero-Modified Geometric distribution with parameter prob and arbitrary probability at zero p0.

## Usage

```
dzmgeom(x, prob, p0, log = FALSE)
pzmgeom(q, prob, p0, lower.tail = TRUE, log.p = FALSE)
qzmgeom(p, prob, p0, lower.tail = TRUE, log.p = FALSE)
rzmgeom(n, prob, p0)
```

## Arguments

x	vector of (strictly positive integer) quantiles.
q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If length(n) > 1, the length is taken to be the number required.
prob	parameter. $0 < \text{prob} \leq 1$ .
p0	probability mass at zero. $0 \leq p_0 \leq 1$ .
log, log.p	logical; if TRUE, probabilities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .

## Details

The zero-modified geometric distribution with  $\text{prob} = p$  and  $p_0 = p_0$  is a discrete mixture between a degenerate distribution at zero and a (standard) geometric. The probability mass function is  $p(0) = p_0$  and

$$p(x) = \frac{(1 - p_0)}{(1 - p)} f(x)$$

for  $x = 1, 2, \dots$ ,  $0 < p < 1$  and  $0 \leq p_0 \leq 1$ , where  $f(x)$  is the probability mass function of the geometric. The cumulative distribution function is

$$P(x) = p_0 + (1 - p_0) \left( \frac{F(x) - F(0)}{1 - F(0)} \right)$$

The mean is  $(1 - p_0)\mu$  and the variance is  $(1 - p_0)\sigma^2 + p_0(1 - p_0)\mu^2$ , where  $\mu$  and  $\sigma^2$  are the mean and variance of the zero-truncated geometric.

In the terminology of Klugman et al. (2012), the zero-modified geometric is a member of the  $(a, b, 1)$  class of distributions with  $a = 1 - p$  and  $b = 0$ .

The special case  $p\theta = 0$  is the zero-truncated geometric.

If an element of  $x$  is not integer, the result of `dzmgeom` is zero, with a warning.

The quantile is defined as the smallest value  $x$  such that  $P(x) \geq p$ , where  $P$  is the distribution function.

### Value

`dzmgeom` gives the (log) probability mass function, `pzmgeom` gives the (log) distribution function, `qzmgeom` gives the quantile function, and `rzmgeom` generates random deviates.

Invalid `prob` or `p0` will result in return value `NaN`, with a warning.

The length of the result is determined by `n` for `rzmgeom`, and is the maximum of the lengths of the numerical arguments for the other functions.

### Note

Functions `{d,p,q}zmgeom` use `{d,p,q}geom` for all but the trivial input values and  $p(0)$ .

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>

### References

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

### See Also

[dgeom](#) for the geometric distribution.

[dztgeom](#) for the zero-truncated geometric distribution.

[dzmnbinom](#) for the zero-modified negative binomial, of which the zero-modified geometric is a special case.

### Examples

```
p <- 1/(1 + 0.5)
dzmgeom(1:5, prob = p, p0 = 0.6)
(1-0.6) * dgeom(1:5, p)/pgeom(0, p, lower = FALSE) # same

## simple relation between survival functions
pzmgeom(0:5, p, p0 = 0.2, lower = FALSE)
(1-0.2) * pgeom(0:5, p, lower = FALSE)/pgeom(0, p, lower = FALSE) # same

qzmgeom(pzmgeom(0:10, 0.3, p0 = 0.6), 0.3, p0 = 0.6)
```



---

 ZeroModifiedLogarithmic

*The Zero-Modified Logarithmic Distribution*


---

## Description

Density function, distribution function, quantile function and random generation for the Zero-Modified Logarithmic (or log-series) distribution with parameter *prob* and arbitrary probability at zero *p0*.

## Usage

```

dzmlogarithmic(x, prob, p0, log = FALSE)
pzmlogarithmic(q, prob, p0, lower.tail = TRUE, log.p = FALSE)
qzmlogarithmic(p, prob, p0, lower.tail = TRUE, log.p = FALSE)
rzmlogarithmic(n, prob, p0)
  
```

## Arguments

<i>x</i>	vector of (strictly positive integer) quantiles.
<i>q</i>	vector of quantiles.
<i>p</i>	vector of probabilities.
<i>n</i>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<i>prob</i>	parameter. $0 \leq \text{prob} < 1$ .
<i>p0</i>	probability mass at zero. $0 \leq p0 \leq 1$ .
<i>log, log.p</i>	logical; if TRUE, probabilities <i>p</i> are returned as $\log(p)$ .
<i>lower.tail</i>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .

## Details

The zero-modified logarithmic distribution with *prob* = *p* and *p0* = *p0* is a discrete mixture between a degenerate distribution at zero and a (standard) logarithmic. The probability mass function is  $p(0) = p_0$  and

$$p(x) = (1 - p_0)f(x)$$

for  $x = 1, 2, \dots$ ,  $0 < p < 1$  and  $0 \leq p_0 \leq 1$ , where  $f(x)$  is the probability mass function of the logarithmic. The cumulative distribution function is

$$P(x) = p_0 + (1 - p_0)F(x)$$

The special case  $p_0 = 0$  is the standard logarithmic.

The zero-modified logarithmic distribution is the limiting case of the zero-modified negative binomial distribution with size parameter equal to 0. Note that in this context, parameter *prob* generally corresponds to the probability of *failure* of the zero-truncated negative binomial.

If an element of  $x$  is not integer, the result of `dzmlogarithmic` is zero, with a warning.

The quantile is defined as the smallest value  $x$  such that  $F(x) \geq p$ , where  $F$  is the distribution function.

### Value

`dzmlogarithmic` gives the probability mass function, `pzmlogarithmic` gives the distribution function, `qzmlogarithmic` gives the quantile function, and `rzmlogarithmic` generates random deviates.

Invalid `prob` or `p0` will result in return value `NaN`, with a warning.

The length of the result is determined by `n` for `rzmlogarithmic`, and is the maximum of the lengths of the numerical arguments for the other functions.

### Note

Functions `{d,p,q}zmlogarithmic` use `{d,p,q}logarithmic` for all but the trivial input values and  $p(0)$ .

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>

### References

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

### See Also

[dlogarithmic](#) for the logarithmic distribution.

[dztnbinom](#) for the zero modified negative binomial distribution.

### Examples

```
p <- 1/(1 + 0.5)
dzmlogarithmic(1:5, prob = p, p0 = 0.6)
(1-0.6) * dlogarithmic(1:5, p)/plogarithmic(0, p, lower = FALSE) # same

## simple relation between survival functions
pzmlogarithmic(0:5, p, p0 = 0.2, lower = FALSE)
(1-0.2) * plogarithmic(0:5, p, lower = FALSE)/plogarithmic(0, p, lower = FALSE) # same

qzmlogarithmic(pzmlogarithmic(0:10, 0.3, p0 = 0.6), 0.3, p0 = 0.6)
```

---

ZeroModifiedNegativeBinomial

*The Zero-Modified Negative Binomial Distribution*


---

## Description

Density function, distribution function, quantile function and random generation for the Zero-Modified Negative Binomial distribution with parameters `size` and `prob`, and arbitrary probability at zero `p0`.

## Usage

```
dzmnbinom(x, size, prob, p0, log = FALSE)
pzmnbinom(q, size, prob, p0, lower.tail = TRUE, log.p = FALSE)
qzmnbinom(p, size, prob, p0, lower.tail = TRUE, log.p = FALSE)
rzmnbinom(n, size, prob, p0)
```

## Arguments

<code>x</code>	vector of (strictly positive integer) quantiles.
<code>q</code>	vector of quantiles.
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations. If <code>length(n) &gt; 1</code> , the length is taken to be the number required.
<code>size</code>	target for number of successful trials, or dispersion parameter. Must be positive, need not be integer.
<code>prob</code>	parameter. $0 < \text{prob} \leq 1$ .
<code>p0</code>	probability mass at zero. $0 \leq p0 \leq 1$ .
<code>log, log.p</code>	logical; if TRUE, probabilities <code>p</code> are returned as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .

## Details

The zero-modified negative binomial distribution with `size = r`, `prob = p` and `p0 = p0` is a discrete mixture between a degenerate distribution at zero and a (standard) negative binomial. The probability mass function is  $p(0) = p_0$  and

$$p(x) = \frac{(1 - p_0)}{(1 - p^r)} f(x)$$

for  $x = 1, 2, \dots, r \geq 0$ ,  $0 < p < 1$  and  $0 \leq p_0 \leq 1$ , where  $f(x)$  is the probability mass function of the negative binomial. The cumulative distribution function is

$$P(x) = p_0 + (1 - p_0) \left( \frac{F(x) - F(0)}{1 - F(0)} \right)$$

The mean is  $(1 - p_0)\mu$  and the variance is  $(1 - p_0)\sigma^2 + p_0(1 - p_0)\mu^2$ , where  $\mu$  and  $\sigma^2$  are the mean and variance of the zero-truncated negative binomial.

In the terminology of Klugman et al. (2012), the zero-modified negative binomial is a member of the  $(a, b, 1)$  class of distributions with  $a = 1 - p$  and  $b = (r - 1)(1 - p)$ .

The special case  $p_0 == 0$  is the zero-truncated negative binomial.

The limiting case  $size == 0$  is the zero-modified logarithmic distribution with parameters  $1 - prob$  and  $p_0$ .

Unlike the standard negative binomial functions, parametrization through the mean  $\mu$  is not supported to avoid ambiguity as to whether  $\mu$  is the mean of the underlying negative binomial or the mean of the zero-modified distribution.

If an element of  $x$  is not integer, the result of `dzmnbinom` is zero, with a warning.

The quantile is defined as the smallest value  $x$  such that  $P(x) \geq p$ , where  $P$  is the distribution function.

### Value

`dzmnbinom` gives the (log) probability mass function, `pzmnbino` gives the (log) distribution function, `qzmnbinom` gives the quantile function, and `rzmnbinom` generates random deviates.

Invalid `size`, `prob` or `p0` will result in return value `NaN`, with a warning.

The length of the result is determined by `n` for `rzmnbinom`, and is the maximum of the lengths of the numerical arguments for the other functions.

### Note

Functions `{d,p,q}zmnbinom` use `{d,p,q}nbinom` for all but the trivial input values and  $p(0)$ .

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>

### References

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

### See Also

[dnbinom](#) for the negative binomial distribution.

[dztnbinom](#) for the zero-truncated negative binomial distribution.

[dzmgeom](#) for the zero-modified geometric and [dzmlogarithmic](#) for the zero-modified logarithmic, which are special cases of the zero-modified negative binomial.

**Examples**

```
## Example 6.3 of Klugman et al. (2012)
p <- 1/(1 + 0.5)
dzmnbinom(1:5, size = 2.5, prob = p, p0 = 0.6)
(1-0.6) * dnbinom(1:5, 2.5, p)/pnbinom(0, 2.5, p, lower = FALSE) # same

## simple relation between survival functions
pzmnbinom(0:5, 2.5, p, p0 = 0.2, lower = FALSE)
(1-0.2) * pnbinom(0:5, 2.5, p, lower = FALSE) /
  pnbinom(0, 2.5, p, lower = FALSE) # same

qzmnbinom(pzmnbinom(0:10, 2.5, 0.3, p0 = 0.1), 2.5, 0.3, p0 = 0.1)
```

---

ZeroModifiedPoisson    *The Zero-Modified Poisson Distribution*

---

**Description**

Density function, distribution function, quantile function, random generation for the Zero-Modified Poisson distribution with parameter lambda and arbitrary probability at zero p0.

**Usage**

```
dzmpois(x, lambda, p0, log = FALSE)
pzmmpois(q, lambda, p0, lower.tail = TRUE, log.p = FALSE)
qzmmpois(p, lambda, p0, lower.tail = TRUE, log.p = FALSE)
rzmpois(n, lambda, p0)
```

**Arguments**

x	vector of (strictly positive integer) quantiles.
q	vector of quantiles.
p	vector of probabilities.
n	number of values to return.
lambda	vector of (non negative) means.
p0	probability mass at zero. $0 \leq p0 \leq 1$ .
log, log.p	logical; if TRUE, probabilities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .

**Details**

The zero-modified Poisson distribution is a discrete mixture between a degenerate distribution at zero and a (standard) Poisson. The probability mass function is  $p(0) = p_0$  and

$$p(x) = \frac{(1 - p_0)}{(1 - e^{-\lambda})} f(x)$$

for  $x = 1, 2, \dots$ ,  $\lambda > 0$  and  $0 \leq p_0 \leq 1$ , where  $f(x)$  is the probability mass function of the Poisson. The cumulative distribution function is

$$P(x) = p_0 + (1 - p_0) \left( \frac{F(x) - F(0)}{1 - F(0)} \right).$$

The mean is  $(1 - p_0)\mu$  and the variance is  $(1 - p_0)\sigma^2 + p_0(1 - p_0)\mu^2$ , where  $\mu$  and  $\sigma^2$  are the mean and variance of the zero-truncated Poisson.

In the terminology of Klugman et al. (2012), the zero-modified Poisson is a member of the  $(a, b, 1)$  class of distributions with  $a = 0$  and  $b = \lambda$ .

The special case  $p_0 = 0$  is the zero-truncated Poisson.

If an element of  $x$  is not integer, the result of `dzmpois` is zero, with a warning.

The quantile is defined as the smallest value  $x$  such that  $P(x) \geq p$ , where  $P$  is the distribution function.

### Value

`dzmpois` gives the (log) probability mass function, `pzmpois` gives the (log) distribution function, `qzmpois` gives the quantile function, and `rzmpois` generates random deviates.

Invalid `lambda` or `p0` will result in return value `NaN`, with a warning.

The length of the result is determined by `n` for `rzmpois`, and is the maximum of the lengths of the numerical arguments for the other functions.

### Note

Functions `{d,p,q}zmpois` use `{d,p,q}pois` for all but the trivial input values and  $p(0)$ .

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>

### References

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

### See Also

[dpois](#) for the standard Poisson distribution.

[dztpois](#) for the zero-truncated Poisson distribution.

### Examples

```
dzmpois(0:5, lambda = 1, p0 = 0.2)
(1-0.2) * dpois(0:5, lambda = 1)/ppois(0, 1, lower = FALSE) # same

## simple relation between survival functions
pzmpois(0:5, 1, p0 = 0.2, lower = FALSE)
```

```
(1-0.2) * ppois(0:5, 1, lower = FALSE) /
  ppois(0, 1, lower = FALSE) # same

qzmpois(pzmpois(0:10, 1, p0 = 0.7), 1, p0 = 0.7)
```

## ZeroTruncatedBinomial *The Zero-Truncated Binomial Distribution*

### Description

Density function, distribution function, quantile function and random generation for the Zero-Truncated Binomial distribution with parameters size and prob.

### Usage

```
dztbinom(x, size, prob, log = FALSE)
pztbinom(q, size, prob, lower.tail = TRUE, log.p = FALSE)
qztbinom(p, size, prob, lower.tail = TRUE, log.p = FALSE)
rztbinom(n, size, prob)
```

### Arguments

x	vector of (strictly positive integer) quantiles.
q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If length(n) > 1, the length is taken to be the number required.
size	number of trials (strictly positive integer).
prob	probability of success on each trial. $0 \leq \text{prob} \leq 1$ .
log, log.p	logical; if TRUE, probabilities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .

### Details

The zero-truncated binomial distribution with size =  $n$  and prob =  $p$  has probability mass function

$$p(x) = \binom{n}{x} \frac{p^x (1-p)^{n-x}}{1 - (1-p)^n}$$

for  $x = 1, \dots, n$  and  $0 < p \leq 1$ , and  $p(1) = 1$  when  $p = 0$ . The cumulative distribution function is

$$P(x) = \frac{F(x) - F(0)}{1 - F(0)},$$

where  $F(x)$  is the distribution function of the standard binomial.

The mean is  $np/(1 - (1-p)^n)$  and the variance is  $np[(1-p) - (1-p+np)(1-p)^n]/[1 - (1-p)^n]^2$ .

In the terminology of Klugman et al. (2012), the zero-truncated binomial is a member of the  $(a, b, 1)$  class of distributions with  $a = -p/(1-p)$  and  $b = (n+1)p/(1-p)$ .

If an element of  $x$  is not integer, the result of `dztbinom` is zero, with a warning.

The quantile is defined as the smallest value  $x$  such that  $P(x) \geq p$ , where  $P$  is the distribution function.

### Value

`dztbinom` gives the probability mass function, `pztbinom` gives the distribution function, `qztbinom` gives the quantile function, and `rztbinom` generates random deviates.

Invalid `size` or `prob` will result in return value `NaN`, with a warning.

The length of the result is determined by `n` for `rztbinom`, and is the maximum of the lengths of the numerical arguments for the other functions.

### Note

Functions `{d,p,q}zbtinom` use `{d,p,q}binom` for all but the trivial input values and  $p(0)$ .

`rztbinom` uses the simple inversion algorithm suggested by Peter Dalgaard on the r-help mailing list on 1 May 2005 (<https://stat.ethz.ch/pipermail/r-help/2005-May/070680.html>).

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>

### References

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

### See Also

[dbinom](#) for the binomial distribution.

### Examples

```
dztbinom(1:5, size = 5, prob = 0.4)
dbinom(1:5, 5, 0.4)/pbinom(0, 5, 0.4, lower = FALSE) # same

pztbinom(1, 2, prob = 0)          # point mass at 1

qztbinom(pztbinom(1:10, 10, 0.6), 10, 0.6)

n <- 8; p <- 0.3
x <- 0:n
title <- paste("ZT Binomial(", n, ", ", p,
              ") and Binomial(", n, ", ", p, ") PDF",
              sep = "")
plot(x, dztbinom(x, n, p), type = "h", lwd = 2, ylab = "p(x)",
     main = title)
```



```
points(x, dbinom(x, n, p), pch = 19, col = "red")
legend("topright", c("ZT binomial probabilities", "Binomial probabilities"),
      col = c("black", "red"), lty = c(1, 0), lwd = 2, pch = c(NA, 19))
```

ZeroTruncatedGeometric

*The Zero-Truncated Geometric Distribution***Description**

Density function, distribution function, quantile function and random generation for the Zero-Truncated Geometric distribution with parameter prob.

**Usage**

```
dztgeom(x, prob, log = FALSE)
pztgeom(q, prob, lower.tail = TRUE, log.p = FALSE)
qztgeom(p, prob, lower.tail = TRUE, log.p = FALSE)
rztgeom(n, prob)
```

**Arguments**

x	vector of (strictly positive integer) quantiles.
q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If length(n) > 1, the length is taken to be the number required.
prob	parameter. $0 < \text{prob} \leq 1$ .
log, log.p	logical; if TRUE, probabilities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .

**Details**

The zero-truncated geometric distribution with  $\text{prob} = p$  has probability mass function

$$p(x) = p(1 - p)^{x-1}$$

for  $x = 1, 2, \dots$  and  $0 < p < 1$ , and  $p(1) = 1$  when  $p = 1$ . The cumulative distribution function is

$$P(x) = \frac{F(x) - F(0)}{1 - F(0)},$$

where  $F(x)$  is the distribution function of the standard geometric.

The mean is  $1/p$  and the variance is  $(1 - p)/p^2$ .

In the terminology of Klugman et al. (2012), the zero-truncated geometric is a member of the  $(a, b, 1)$  class of distributions with  $a = 1 - p$  and  $b = 0$ .

If an element of  $x$  is not integer, the result of `dztgeom` is zero, with a warning.

The quantile is defined as the smallest value  $x$  such that  $P(x) \geq p$ , where  $P$  is the distribution function.

**Value**

dztgeom gives the (log) probability mass function, pztgeom gives the (log) distribution function, qztgeom gives the quantile function, and rztgeom generates random deviates.

Invalid prob will result in return value NaN, with a warning.

The length of the result is determined by n for rztgeom, and is the maximum of the lengths of the numerical arguments for the other functions.

**Note**

Functions {d,p,q}ztgeom use {d,p,q}geom for all but the trivial input values and  $p(0)$ .

rztgeom uses the simple inversion algorithm suggested by Peter Dalgaard on the r-help mailing list on 1 May 2005 (<https://stat.ethz.ch/pipermail/r-help/2005-May/070680.html>).

**Author(s)**

Vincent Goulet <vincent.goulet@act.ulaval.ca>

**References**

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

**See Also**

[dgeom](#) for the geometric distribution.

[dztnbinom](#) for the zero-truncated negative binomial, of which the zero-truncated geometric is a special case.

**Examples**

```
p <- 1/(1 + 0.5)
dztgeom(c(1, 2, 3), prob = p)
dgeom(c(1, 2, 3), p)/pgeom(0, p, lower = FALSE) # same
dgeom(c(1, 2, 3) - 1, p)                        # same

pztgeom(1, prob = 1)          # point mass at 1

qztgeom(pztgeom(1:10, 0.3), 0.3)
```

---

ZeroTruncatedNegativeBinomial

*The Zero-Truncated Negative Binomial Distribution*


---

## Description

Density function, distribution function, quantile function and random generation for the Zero-Truncated Negative Binomial distribution with parameters size and prob.

## Usage

```
dztnbinom(x, size, prob, log = FALSE)
pztnbinom(q, size, prob, lower.tail = TRUE, log.p = FALSE)
qztnbinom(p, size, prob, lower.tail = TRUE, log.p = FALSE)
rztnbinom(n, size, prob)
```

## Arguments

x	vector of (strictly positive integer) quantiles.
q	vector of quantiles.
p	vector of probabilities.
n	number of observations. If length(n) > 1, the length is taken to be the number required.
size	target for number of successful trials, or dispersion parameter. Must be positive, need not be integer.
prob	parameter. $0 < \text{prob} \leq 1$ .
log, log.p	logical; if TRUE, probabilities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .

## Details

The zero-truncated negative binomial distribution with size =  $r$  and prob =  $p$  has probability mass function

$$p(x) = \frac{\Gamma(x+r)p^r(1-p)^x}{\Gamma(r)x!(1-p^r)}$$

for  $x = 1, 2, \dots$ ,  $r \geq 0$  and  $0 < p < 1$ , and  $p(1) = 1$  when  $p = 1$ . The cumulative distribution function is

$$P(x) = \frac{F(x) - F(0)}{1 - F(0)},$$

where  $F(x)$  is the distribution function of the standard negative binomial.

The mean is  $r(1-p)/(p(1-p^r))$  and the variance is  $[r(1-p)(1 - (1+r(1-p))p^r)]/[p(1-p^r)]^2$ .

In the terminology of Klugman et al. (2012), the zero-truncated negative binomial is a member of the  $(a, b, 1)$  class of distributions with  $a = 1 - p$  and  $b = (r - 1)(1 - p)$ .

The limiting case `size == 0` is the [logarithmic](#) distribution with parameter `1 - prob`.

Unlike the standard negative binomial functions, parametrization through the mean `mu` is not supported to avoid ambiguity as to whether `mu` is the mean of the underlying negative binomial or the mean of the zero-truncated distribution.

If an element of `x` is not integer, the result of `dztnbinom` is zero, with a warning.

The quantile is defined as the smallest value  $x$  such that  $P(x) \geq p$ , where  $P$  is the distribution function.

### Value

`dztnbinom` gives the (log) probability mass function, `pztnbinom` gives the (log) distribution function, `qztnbinom` gives the quantile function, and `rztnbinom` generates random deviates.

Invalid `size` or `prob` will result in return value `NaN`, with a warning.

The length of the result is determined by `n` for `rztnbinom`, and is the maximum of the lengths of the numerical arguments for the other functions.

### Note

Functions `{d,p,q}ztnbinom` use `{d,p,q}nbinom` for all but the trivial input values and  $p(0)$ .

`rztnbinom` uses the simple inversion algorithm suggested by Peter Dalgaard on the r-help mailing list on 1 May 2005 (<https://stat.ethz.ch/pipermail/r-help/2005-May/070680.html>).

### Author(s)

Vincent Goulet <[vincent.goulet@act.ulaval.ca](mailto:vincent.goulet@act.ulaval.ca)>

### References

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

### See Also

[dnbinom](#) for the negative binomial distribution.

[dztgeom](#) for the zero-truncated geometric and [dlogarithmic](#) for the logarithmic, which are special cases of the zero-truncated negative binomial.

### Examples

```
## Example 6.3 of Klugman et al. (2012)
p <- 1/(1 + 0.5)
dztnbinom(c(1, 2, 3), size = 2.5, prob = p)
dnbinom(c(1, 2, 3), 2.5, p)/pnbinom(0, 2.5, p, lower = FALSE) # same

pztnbinom(1, 2, prob = 1)      # point mass at 1
dztnbinom(2, size = 1, 0.25)  # == dztgeom(2, 0.25)
dztnbinom(2, size = 0, 0.25)  # == dlogarithmic(2, 0.75)
```

```

qztnbinom(pztnbinom(1:10, 2.5, 0.3), 2.5, 0.3)

x <- rztnbinom(1000, size = 2.5, prob = 0.4)
y <- sort(unique(x))
plot(y, table(x)/length(x), type = "h", lwd = 2,
     pch = 19, col = "black", xlab = "x", ylab = "p(x)",
     main = "Empirical vs theoretical probabilities")
points(y, dztnbinom(y, size = 2.5, prob = 0.4),
       pch = 19, col = "red")
legend("topright", c("empirical", "theoretical"),
      lty = c(1, NA), lwd = 2, pch = c(NA, 19), col = c("black", "red"))

```

ZeroTruncatedPoisson *The Zero-Truncated Poisson Distribution*

## Description

Density function, distribution function, quantile function, random generation for the Zero-Truncated Poisson distribution with parameter lambda.

## Usage

```

dztpois(x, lambda, log = FALSE)
pztpois(q, lambda, lower.tail = TRUE, log.p = FALSE)
qztpois(p, lambda, lower.tail = TRUE, log.p = FALSE)
rztpois(n, lambda)

```

## Arguments

x	vector of (strictly positive integer) quantiles.
q	vector of quantiles.
p	vector of probabilities.
n	number of values to return.
lambda	vector of (non negative) means.
log, log.p	logical; if TRUE, probabilities $p$ are returned as $\log(p)$ .
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .

## Details

The zero-truncated Poisson distribution has probability mass function

$$p(x) = \frac{e^{-\lambda} \lambda^x}{x!(1 - e^{-\lambda})} = \frac{\lambda^x}{x!(e^\lambda - 1)}$$

for  $x = 1, 2, \dots$ , and  $p(1) = 1$  when  $\lambda = 0$ . The cumulative distribution function is

$$P(x) = \frac{F(x) - F(0)}{1 - F(0)},$$

where  $F(x)$  is the distribution function of the standard Poisson.

The mean is  $\lambda/(1 - e^{-\lambda})^2$  and the variance is  $\lambda[1 - (\lambda + 1)e^{-\lambda}]/(1 - e^{-\lambda})^2$ .

In the terminology of Klugman et al. (2012), the zero-truncated Poisson is a member of the  $(a, b, 1)$  class of distributions with  $a = 0$  and  $b = \lambda$ .

If an element of  $x$  is not integer, the result of `dztpois` is zero, with a warning.

The quantile is defined as the smallest value  $x$  such that  $P(x) \geq p$ , where  $P$  is the distribution function.

### Value

`dztpois` gives the (log) probability mass function, `pztpois` gives the (log) distribution function, `qztpois` gives the quantile function, and `rztpois` generates random deviates.

Invalid `lambda` will result in return value `NaN`, with a warning.

The length of the result is determined by `n` for `rztpois`, and is the maximum of the lengths of the numerical arguments for the other functions.

### Note

Functions `{d,p,q}ztpois` use `{d,p,q}pois` for all but the trivial input values and  $p(0)$ .

`rztpois` uses the simple inversion algorithm suggested by Peter Dalgaard on the r-help mailing list on 1 May 2005 (<https://stat.ethz.ch/pipermail/r-help/2005-May/070680.html>).

### Author(s)

Vincent Goulet <vincent.goulet@act.ulaval.ca>

### References

Klugman, S. A., Panjer, H. H. and Willmot, G. E. (2012), *Loss Models, From Data to Decisions, Fourth Edition*, Wiley.

### See Also

[dpois](#) for the standard Poisson distribution.

### Examples

```
dztpois(1:5, lambda = 1)
dpois(1:5, lambda = 1)/ppois(0, 1, lower = FALSE) # same

pztpois(1, lambda = 0)          # point mass at 1

qztpois(pztpois(1:10, 1), 1)

x <- seq(0, 8)
plot(x, dztpois(x, 2), type = "h", lwd = 2, ylab = "p(x)",
     main = "Zero-Truncated Poisson(2) and Poisson(2) PDF")
points(x, dpois(x, 2), pch = 19, col = "red")
legend("topright", c("ZT Poisson probabilities", "Poisson probabilities"),
```

```
col = c("black", "red"), lty = c(1, 0), lwd = 2, pch = c(NA, 19))
```

# Index

- \* **array**
  - Extract.grouped.data, 35
- \* **classes**
  - grouped.data, 45
- \* **datagen**
  - rcompound, 95
  - rmixture, 97
  - severity, 100
  - simul, 101
- \* **datasets**
  - dental, 27
  - gdental, 40
  - hachemeister, 49
- \* **distribution**
  - aggregateDist, 6
  - betaint, 11
  - BetaMoments, 13
  - Burr, 14
  - ChisqSupp, 16
  - discretize, 28
  - ExponentialSupp, 33
  - FellerPareto, 36
  - GammaSupp, 39
  - GeneralizedBeta, 40
  - GeneralizedPareto, 42
  - Gumbel, 47
  - hist.grouped.data, 49
  - InverseBurr, 51
  - InverseExponential, 53
  - InverseGamma, 54
  - InverseGaussian, 56
  - InverseParalogistic, 59
  - InversePareto, 60
  - InverseTransformedGamma, 62
  - InverseWeibull, 64
  - Logarithmic, 66
  - Loggamma, 68
  - Loglogistic, 69
  - LognormalMoments, 71
  - mde, 72
  - NormalSupp, 75
  - Paralogistic, 78
  - Pareto, 80
  - Pareto2, 82
  - Pareto3, 84
  - Pareto4, 86
  - PhaseType, 88
  - PoissonInverseGaussian, 90
  - SingleParameterPareto, 107
  - TransformedBeta, 108
  - TransformedGamma, 111
  - UniformSupp, 113
  - WeibullMoments, 116
  - ZeroModifiedBinomial, 117
  - ZeroModifiedGeometric, 119
  - ZeroModifiedLogarithmic, 121
  - ZeroModifiedNegativeBinomial, 123
  - ZeroModifiedPoisson, 125
  - ZeroTruncatedBinomial, 127
  - ZeroTruncatedGeometric, 129
  - ZeroTruncatedNegativeBinomial, 131
  - ZeroTruncatedPoisson, 133
- \* **dplot**
  - elev, 30
  - hist.grouped.data, 49
  - ogive, 76
- \* **hplot**
  - elev, 30
  - hist.grouped.data, 49
  - ogive, 76
- \* **htest**
  - mde, 72
- \* **manip**
  - Extract.grouped.data, 35
  - severity, 100
  - unroll, 114
- \* **math**
  - betaint, 11



- \* **methods**
  - grouped.data, 45
  - simul.summaries, 104
- \* **models**
  - aggregateDist, 6
  - cm, 18
  - coverage, 24
  - discretize, 28
  - ruin, 98
  - simul.summaries, 104
- \* **optimize**
  - adjCoef, 4
- \* **univar**
  - adjCoef, 4
  - CTE, 26
  - emm, 32
  - mean.grouped.data, 74
  - quantile.aggregateDist, 93
  - quantile.grouped.data, 94
  - VaR, 115
- [.data.frame, 36
- [.grouped.data, 46
- [.grouped.data (Extract.grouped.data), 35
- [<- .grouped.data (Extract.grouped.data), 35
- adjCoef, 4
- aggregate.portfolio (simul.summaries), 104
- aggregateDist, 6, 26, 27, 29, 93, 115
- approxfun, 77
- as.data.frame, 18
- as.integer, 35
- axis, 50
- besselK, 91
- Beta, 14
- betaint, 11, 16, 38, 44, 52, 60, 71, 79, 81, 83, 85, 88, 110
- BetaMoments, 13
- Burr, 14, 87, 109
- ChisqSupp, 16
- Chisquare, 17
- class, 100, 103, 104
- cm, 18
- colMeans, 32
- Coverage (coverage), 24
- coverage, 24
- CTE, 26
- CTE.aggregateDist, 10
- curve, 28, 99
- data.frame, 45, 46
- dbeta, 22
- dbinom, 8, 22, 118, 128
- dburr, 88
- dburr (Burr), 14
- dental, 27
- dexp, 99
- dfpareto, 110
- dfpareto (FellerPareto), 36
- dgamma, 22, 99
- dgenbeta (GeneralizedBeta), 40
- dgenpareto (GeneralizedPareto), 42
- dgeom, 8, 120, 130
- dgumbel (Gumbel), 47
- diff.aggregateDist (aggregateDist), 6
- dim, 114
- dinvburr (InverseBurr), 51
- dinvexp (InverseExponential), 53
- dinvgamma, 58
- dinvgamma (InverseGamma), 54
- dinvgauss, 91, 92
- dinvgauss (InverseGaussian), 56
- dinvparalogis (InverseParalogistic), 59
- dinvpareto (InversePareto), 60
- dinvtrgamma (InverseTransformedGamma), 62
- dinvweibull (InverseWeibull), 64
- discretise (discretize), 28
- discretize, 10, 28
- dlgamma (Loggamma), 68
- dlgompertz (InverseWeibull), 64
- dllogis, 86
- dllogis (Loglogistic), 69
- dlogarithmic, 8, 122, 132
- dlogarithmic (Logarithmic), 66
- dnbinom, 8, 22, 124, 132
- dnorm, 22
- dparalogis (Paralogistic), 78
- dpareto, 84, 108
- dpareto (Pareto), 80
- dpareto1, 81
- dpareto1 (SingleParameterPareto), 107
- dpareto2, 81
- dpareto2 (Pareto2), 82

- dpareto3, [71](#)
- dpareto3 (Pareto3), [84](#)
- dpareto4, [16](#)
- dpareto4 (Pareto4), [86](#)
- dpearson6 (TransformedBeta), [108](#)
- dphtype, [99](#)
- dphtype (PhaseType), [88](#)
- dpig (PoissonInverseGaussian), [90](#)
- dpois, [8](#), [92](#), [126](#), [134](#)
- dpoisinvgauss (PoissonInverseGaussian), [90](#)
- dtrbeta, [38](#)
- dtrbeta (TransformedBeta), [108](#)
- dtrgamma (TransformedGamma), [111](#)
- dzmbinom (ZeroModifiedBinomial), [117](#)
- dzmgeom, [124](#)
- dzmgeom (ZeroModifiedGeometric), [119](#)
- dzmlogarithmic, [124](#)
- dzmlogarithmic (ZeroModifiedLogarithmic), [121](#)
- dzmnbinom, [120](#)
- dzmnbinom (ZeroModifiedNegativeBinomial), [123](#)
- dzmpois (ZeroModifiedPoisson), [125](#)
- dztbinom, [118](#)
- dztbinom (ZeroTruncatedBinomial), [127](#)
- dztgeom, [120](#), [132](#)
- dztgeom (ZeroTruncatedGeometric), [129](#)
- dztnbinom, [67](#), [122](#), [124](#), [130](#)
- dztnbinom (ZeroTruncatedNegativeBinomial), [131](#)
- dztpois, [126](#)
- dztpois (ZeroTruncatedPoisson), [133](#)
  
- elev, [30](#)
- emm, [32](#), [75](#)
- Exponential, [34](#), [112](#)
- ExponentialSupp, [33](#)
- expression, [96](#), [97](#), [102](#)
- Extract.grouped.data, [35](#)
  
- Feller-Pareto, [83](#), [85](#), [87](#)
- FellerPareto, [36](#)
- formals, [25](#)
- format, [19](#)
- formula, [18](#), [23](#)
  
- frequency.portfolio (simul.summaries), [104](#)
- function, [31](#), [77](#)
  
- Gamma, [112](#)
- gamma, [12](#), [37](#), [41](#), [43](#), [55](#), [63](#), [68](#), [109](#), [112](#)
- GammaDist, [40](#)
- GammaSupp, [39](#)
- gdental, [40](#)
- generalized Pareto, [110](#)
- GeneralizedBeta, [40](#)
- GeneralizedPareto, [42](#)
- grouped.data, [31](#), [36](#), [40](#), [45](#), [75](#), [77](#), [95](#)
- Gumbel, [47](#)
  
- hachemeister, [49](#)
- hist, [45](#), [46](#), [49](#), [51](#)
- hist.default, [49](#), [51](#)
- hist.grouped.data, [49](#)
  
- inverse Burr, [110](#)
- Inverse Exponential, [55](#), [63](#), [65](#)
- Inverse Gamma, [63](#)
- Inverse Paralogistic, [52](#)
- inverse paralogistic, [110](#)
- Inverse Pareto, [43](#), [52](#)
- inverse Pareto, [110](#)
- Inverse Weibull, [63](#)
- InverseBurr, [51](#)
- InverseExponential, [53](#)
- InverseGamma, [54](#)
- InverseGaussian, [56](#)
- InverseParalogistic, [59](#)
- InversePareto, [60](#)
- InverseTransformedGamma, [62](#)
- InverseWeibull, [64](#)
  
- knots.elev (elev), [30](#)
- knots.ogive (ogive), [76](#)
  
- levbeta (BetaMoments), [13](#)
- levburr (Burr), [14](#)
- levchisq (ChisqSupp), [16](#)
- levexp (ExponentialSupp), [33](#)
- levfpareto (FellerPareto), [36](#)
- levgamma (GammaSupp), [39](#)
- levgenbeta (GeneralizedBeta), [40](#)
- levgenpareto (GeneralizedPareto), [42](#)
- levinvburr (InverseBurr), [51](#)

- levinvexp (InverseExponential), 53
- levinvgamma (InverseGamma), 54
- levinvgauss (InverseGaussian), 56
- levinvparalogis (InverseParalogistic), 59
- levinvpareto (InversePareto), 60
- levinvtrgamma (InverseTransformedGamma), 62
- levinvweibull (InverseWeibull), 64
- levlgamma (Loggamma), 68
- levlgompertz (InverseWeibull), 64
- levllogis (Loglogistic), 69
- levlnorm (LognormalMoments), 71
- levparalogis (Paralogistic), 78
- levpareto (Pareto), 80
- levpareto1 (SingleParameterPareto), 107
- levpareto2 (Pareto2), 82
- levpareto3 (Pareto3), 84
- levpareto4 (Pareto4), 86
- levpearson6 (TransformedBeta), 108
- levtrbeta, 12
- levtrbeta (TransformedBeta), 108
- levtrgamma (TransformedGamma), 111
- levunif (UniformSupp), 113
- levweibull (WeibullMoments), 116
- lines, 4
- lm, 18, 19, 23
- lm.fit, 20
- lm.wfit, 20
- log-series (Logarithmic), 66
- Logarithmic, 66
- logarithmic, 132
- Loggamma, 68
- Loglogistic, 15, 52, 69
- loglogistic, 85, 109
- Lognormal, 72
- LognormalMoments, 71
- mbeta (BetaMoments), 13
- mburr (Burr), 14
- mchisq (ChisqSupp), 16
- Mde (mde), 72
- mde, 72
- mean, 33
- mean.aggregateDist, 10
- mean.aggregateDist (aggregateDist), 6
- mean.grouped.data, 33, 74
- mexp (ExponentialSupp), 33
- mfpareto (FellerPareto), 36
- mgamma (GammaSupp), 39
- mgenbeta (GeneralizedBeta), 40
- mgenpareto (GeneralizedPareto), 42
- mgfchisq (ChisqSupp), 16
- mgfexp (ExponentialSupp), 33
- mgfgamma (GammaSupp), 39
- mgfgumbel (Gumbel), 47
- mgfinvgamma (InverseGamma), 54
- mgfinvgauss (InverseGaussian), 56
- mgfnorm (NormalSupp), 75
- mgfphtype (PhaseType), 88
- mgfunif (UniformSupp), 113
- mgumbel (Gumbel), 47
- minvburr (InverseBurr), 51
- minvexp (InverseExponential), 53
- minvgamma (InverseGamma), 54
- minvgauss (InverseGaussian), 56
- minvparalogis (InverseParalogistic), 59
- minvpareto (InversePareto), 60
- minvtrgamma (InverseTransformedGamma), 62
- minvweibull (InverseWeibull), 64
- mlgamma (Loggamma), 68
- mlgompertz (InverseWeibull), 64
- mllogis (Loglogistic), 69
- mlnorm (LognormalMoments), 71
- mnorm (NormalSupp), 75
- mparalogis (Paralogistic), 78
- mpareto (Pareto), 80
- mpareto1 (SingleParameterPareto), 107
- mpareto2 (Pareto2), 82
- mpareto3 (Pareto3), 84
- mpareto4 (Pareto4), 86
- mpearson6 (TransformedBeta), 108
- mphtype (PhaseType), 88
- mtrbeta (TransformedBeta), 108
- mtrgamma (TransformedGamma), 111
- munif (UniformSupp), 113
- mweibull (WeibullMoments), 116
- Normal, 76
- NormalSupp, 75
- ogive, 76, 95
- optim, 73
- Paralogistic, 15, 78
- paralogistic, 109
- Pareto, 15, 43, 44, 80, 83, 110

- Pareto I, [37](#), [83](#)
- Pareto II, [37](#), [81](#), [87](#)
- Pareto III, [37](#), [87](#)
- Pareto IV, [37](#)
- Pareto2, [82](#)
- Pareto3, [84](#)
- Pareto4, [86](#)
- pburr (Burr), [14](#)
- Pearson6 (TransformedBeta), [108](#)
- pfpareto (FellerPareto), [36](#)
- pgenbeta (GeneralizedBeta), [40](#)
- pgenpareto (GeneralizedPareto), [42](#)
- pgumbel (Gumbel), [47](#)
- PhaseType, [88](#)
- PIG (PoissonInverseGaussian), [90](#)
- pinvburr (InverseBurr), [51](#)
- pinvexp (InverseExponential), [53](#)
- pinvgamma (InverseGamma), [54](#)
- pinvgauss (InverseGaussian), [56](#)
- pinvparalogis (InverseParalogistic), [59](#)
- pinvpareto (InversePareto), [60](#)
- pinvtrgamma (InverseTransformedGamma), [62](#)
- pinvweibull (InverseWeibull), [64](#)
- plgamma (Loggamma), [68](#)
- plgompertz (InverseWeibull), [64](#)
- pllogis (Loglogistic), [69](#)
- plogarithmic (Logarithmic), [66](#)
- plot, [4](#)
- plot.adjCoef (adjCoef), [4](#)
- plot.aggregateDist (aggregateDist), [6](#)
- plot.elev (elev), [30](#)
- plot.histogram, [50](#)
- plot.ogive (ogive), [76](#)
- plot.ruin (ruin), [98](#)
- PoissonInverseGaussian, [90](#)
- pparalogis (Paralogistic), [78](#)
- ppareto (Pareto), [80](#)
- ppareto1 (SingleParameterPareto), [107](#)
- ppareto2 (Pareto2), [82](#)
- ppareto3 (Pareto3), [84](#)
- ppareto4 (Pareto4), [86](#)
- ppearson6 (TransformedBeta), [108](#)
- pphtype, [99](#)
- pphtype (PhaseType), [88](#)
- ppig (PoissonInverseGaussian), [90](#)
- ppoisinvgauss (PoissonInverseGaussian), [90](#)
- predict.cm (cm), [18](#)
- predict.lm, [21](#), [23](#)
- pretty, [45](#)
- print, [31](#), [77](#)
- print.aggregateDist (aggregateDist), [6](#)
- print.cm (cm), [18](#)
- print.elev (elev), [30](#)
- print.ogive (ogive), [76](#)
- print.portfolio (simul), [101](#)
- print.summary.cm (cm), [18](#)
- ptrbeta (TransformedBeta), [108](#)
- ptrgamma (TransformedGamma), [111](#)
- pzmbinom (ZeroModifiedBinomial), [117](#)
- pzmgeom (ZeroModifiedGeometric), [119](#)
- pzmlogarithmic (ZeroModifiedLogarithmic), [121](#)
- pzmnbinom (ZeroModifiedNegativeBinomial), [123](#)
- pzmpois (ZeroModifiedPoisson), [125](#)
- pztbinom (ZeroTruncatedBinomial), [127](#)
- pztgeom (ZeroTruncatedGeometric), [129](#)
- pztnbinom (ZeroTruncatedNegativeBinomial), [131](#)
- pztpois (ZeroTruncatedPoisson), [133](#)
- qburr (Burr), [14](#)
- qfpareto (FellerPareto), [36](#)
- qgenbeta (GeneralizedBeta), [40](#)
- qgenpareto (GeneralizedPareto), [42](#)
- qgumbel (Gumbel), [47](#)
- qinvburr (InverseBurr), [51](#)
- qinvexp (InverseExponential), [53](#)
- qinvgamma (InverseGamma), [54](#)
- qinvgauss (InverseGaussian), [56](#)
- qinvparalogis (InverseParalogistic), [59](#)
- qinvpareto (InversePareto), [60](#)
- qinvtrgamma (InverseTransformedGamma), [62](#)
- qinvweibull (InverseWeibull), [64](#)
- qlgamma (Loggamma), [68](#)
- qlgompertz (InverseWeibull), [64](#)
- qllogis (Loglogistic), [69](#)
- qlogarithmic (Logarithmic), [66](#)
- qparalogis (Paralogistic), [78](#)
- qpareto (Pareto), [80](#)
- qpareto1 (SingleParameterPareto), [107](#)
- qpareto2 (Pareto2), [82](#)

- qpareto3 (Pareto3), 84
- qpareto4 (Pareto4), 86
- qpearson6 (TransformedBeta), 108
- qpig (PoissonInverseGaussian), 90
- qpoisinvgauss (PoissonInverseGaussian), 90
- qtrbeta (TransformedBeta), 108
- qtrgamma (TransformedGamma), 111
- quantile.aggregateDist, 10, 93
- quantile.grouped.data, 77, 94
- qzmbinom (ZeroModifiedBinomial), 117
- qzmggeom (ZeroModifiedGeometric), 119
- qzmlogarithmic (ZeroModifiedLogarithmic), 121
- qzmnbinom (ZeroModifiedNegativeBinomial), 123
- qzmpois (ZeroModifiedPoisson), 125
- qztbinom (ZeroTruncatedBinomial), 127
- qztgeom (ZeroTruncatedGeometric), 129
- qztnbinom (ZeroTruncatedNegativeBinomial), 131
- qztpois (ZeroTruncatedPoisson), 133
- rburr (Burr), 14
- rcomphierarc, 7, 9, 10, 96–98
- rcomphierarc (simul), 101
- rcomphierarc.summaries (simul.summaries), 104
- rcompound, 95, 98, 102, 103
- rcomppois (rcompound), 95
- rffpareto (FellerPareto), 36
- rgenbeta (GeneralizedBeta), 40
- rgenpareto (GeneralizedPareto), 42
- rgumbel (Gumbel), 47
- rinvburr (InverseBurr), 51
- rinvexp (InverseExponential), 53
- rinvgamma (InverseGamma), 54
- rinvgauss (InverseGaussian), 56
- rinvparalogis (InverseParalogistic), 59
- rinvpareto (InversePareto), 60
- rinvtrgamma (InverseTransformedGamma), 62
- rinvweibull (InverseWeibull), 64
- rlgamma (Loggamma), 68
- rlgompertz (InverseWeibull), 64
- rllogis (Loglogistic), 69
- rlogarithmic (Logarithmic), 66
- rmixture, 97
- rparalogis (Paralogistic), 78
- rpareto (Pareto), 80
- rpareto1 (SingleParameterPareto), 107
- rpareto2 (Pareto2), 82
- rpareto3 (Pareto3), 84
- rpareto4 (Pareto4), 86
- rpearson6 (TransformedBeta), 108
- rphtype (PhaseType), 88
- rpig (PoissonInverseGaussian), 90
- rpoisinvgauss (PoissonInverseGaussian), 90
- rtrbeta (TransformedBeta), 108
- rtrgamma (TransformedGamma), 111
- ruin, 98
- rzmbinom (ZeroModifiedBinomial), 117
- rzmggeom (ZeroModifiedGeometric), 119
- rzmlogarithmic (ZeroModifiedLogarithmic), 121
- rzmnbinom (ZeroModifiedNegativeBinomial), 123
- rzmpois (ZeroModifiedPoisson), 125
- rztbinom (ZeroTruncatedBinomial), 127
- rztgeom (ZeroTruncatedGeometric), 129
- rztnbinom (ZeroTruncatedNegativeBinomial), 131
- rztpois (ZeroTruncatedPoisson), 133
- severity, 100, 105
- severity.portfolio, 101, 115
- severity.portfolio (simul.summaries), 104
- simpf (simul), 101
- simul, 101, 105, 106
- simul.summaries, 103, 104
- SingleParameterPareto, 107
- stepfun, 31, 77
- subset, 19, 23
- summary.aggregateDist (aggregateDist), 6
- summary.cm (cm), 18
- summary.elev (elev), 30
- summary.ogive (ogive), 76
- terms, 20
- title, 50
- transformed beta, 37
- TransformedBeta, 108

TransformedGamma, [111](#)  
 TVaR (CTE), [26](#)

Uniform, [114](#)  
 UniformSupp, [113](#)  
 unroll, [101](#), [114](#)

VaR, [27](#), [115](#)  
 VaR.aggregateDist, [115](#)  
 VaR.aggregateDist  
     (quantile.aggregateDist), [93](#)

Weibull, [112](#), [116](#)  
 WeibullMoments, [116](#)  
 weights.portfolio (simul.summaries), [104](#)

ZeroModifiedBinomial, [117](#)  
 ZeroModifiedGeometric, [119](#)  
 ZeroModifiedLogarithmic, [121](#)  
 ZeroModifiedNegativeBinomial, [123](#)  
 ZeroModifiedPoisson, [125](#)  
 ZeroTruncatedBinomial, [127](#)  
 ZeroTruncatedGeometric, [129](#)  
 ZeroTruncatedNegativeBinomial, [131](#)  
 ZeroTruncatedPoisson, [133](#)  
 ZMBinomial (ZeroModifiedBinomial), [117](#)  
 Zmgeometric (ZeroModifiedGeometric), [119](#)  
 ZMLogarithmic  
     (ZeroModifiedLogarithmic), [121](#)  
 ZMNegativeBinomial  
     (ZeroModifiedNegativeBinomial),  
     [123](#)  
 ZMNegBinomial  
     (ZeroModifiedNegativeBinomial),  
     [123](#)  
 Zmpoisson (ZeroModifiedPoisson), [125](#)  
 ZTBinomial (ZeroTruncatedBinomial), [127](#)  
 ZTGeometric (ZeroTruncatedGeometric),  
     [129](#)  
 ZTNegativeBinomial  
     (ZeroTruncatedNegativeBinomial),  
     [131](#)  
 ZTNegBinomial  
     (ZeroTruncatedNegativeBinomial),  
     [131](#)  
 ZTPoisson (ZeroTruncatedPoisson), [133](#)