

# Package ‘soilphysics’

December 7, 2020

**Type** Package

**Title** Soil Physical Analysis

**Version** 4.0

**Date** 2020-12-06

**Description** Basic and model-based soil physical analyses.

**Author** Anderson Rodrigo da Silva [aut, cre]  
(<<https://orcid.org/0000-0003-2518-542X>>),  
Renato Paiva de Lima [aut] (<<https://orcid.org/0000-0003-0524-439X>>)

**Maintainer** Anderson Rodrigo da Silva <[anderson.agro@hotmail.com](mailto:anderson.agro@hotmail.com)>

**URL** <https://arsilva87.github.io/soilphysics/>

**Imports** utils, stats, graphics, boot, grDevices, datasets, MASS,  
shiny, shinydashboard, fields

**Suggests** knitr, rmarkdown, rpanel, tcltk, tkrplot

**VignetteBuilder** knitr

**License** GPL (>= 2)

**NeedsCompilation** no

**Repository** CRAN

**Date/Publication** 2020-12-07 05:20:03 UTC

## R topics documented:

soilphysics-package . . . . .	3
aggreg.stability . . . . .	4
aggreg.stability_App . . . . .	5
bulkDensity . . . . .	6
compaction . . . . .	7
compressive_properties . . . . .	7
compressive_properties2 . . . . .	9
compressive_properties3 . . . . .	10
compressive_properties4 . . . . .	12
compressive_properties5 . . . . .	13

criticalmoisture	15
fitbusscher	16
fitlbc	18
fitsoilwater	19
fitsoilwater2	20
fitsoilwater3	21
fitsoilwater4	22
fitsoilwater5	23
fun2form	25
getInitiallbc	26
hemc	27
hydraulicCutOff	28
hydraulicCutOff2	29
iwc	31
Kr_h	33
Kr_theta	34
liquidlimit	36
llwr	37
llwrPTF	40
LLWR_App	42
llwr_llmpr	43
LLWR_LLMPR_App	46
maxcurv	47
particle.sedimentation	48
particle.sedimentation_App	50
plotCIsigmaP	50
PredComp	52
psd	52
r	53
Rsq	54
r_App	55
sigmaP	56
simSigmaP	59
Sindex	60
skp1994	62
SoilAggregate	62
soilDeformation	63
soilStrength	66
soilStrength2	68
soilStrength3	69
soilStrength4	71
soilStrength5	72
soilwater	73
soilwater2	74
soilwater3	76
soilwater4	77
soilwater5	78
SSlbc	79

<i>soilphysics-package</i>	3
stressTraffic . . . . .	80
voidratio . . . . .	82
<b>Index</b>	<b>84</b>

---

soilphysics-package     *Soil Physical Analysis*

---

## Description

Basic and model-based soil physical analyses.

## Details

Package: soilphysics  
Type: Package  
Version: 4.0  
Date: 2020-12-06  
License: GPL (>= 2)

Functions for modelling the load bearing capacity and the penetration resistance, and for predicting the stress applied by agricultural machines in the soil profile. The package allows one to model the soil water retention through six different models. There are some useful and easy-to-use functions to perform parameter estimation of these models. Methods to obtain the preconsolidation stress are available, such as the standard of Casagrande (1936) and so on. It is possible to quantify soil water availability for plants through the Least Limiting Water Range approach as well as the Integral Water Capacity. Moreover, it is possible to determine the water suction at the point of hydraulic cut-off. Also, users can deal with the high-energy-moisture-characteristics (HEMC) methodology proposed by Pierson and Mulla (1989), which is used to analyze the aggregate stability. There is a function to determine the soil critical moisture and the maximum bulk density for one or more samples, based on the Proctor (1933) compaction test. Other utilities like a function to calculate the soil liquid limit, the void ratio and to determine the maximum curvature point are available.

## Note

*soilphysics* is an ongoing project. We welcome any and all criticism, comments and suggestions.

## Author(s)

Anderson Rodrigo da Silva, Renato Paiva de Lima  
Maintainer: Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

## References

da Silva, A.R.; de Lima, R.P. (2015) soilphysics: an R package to determine soil preconsolidation pressure. *Computers and Geosciences*, 84: 54-60.

de Lima, R.P.; da Silva, A.R.; da Silva, A.P.; Leao, T.P.; Mosaddeghi, M.R. (2016) soilphysics: an R package for calculating soil water availability to plants by different soil physical indices. *Computers and Electronics in Agriculture*, 120: 63-71.

da Silva, A.R.; Lima, R.P. (2017) Determination of maximum curvature point with the R package soilphysics. *International Journal of Current Research*, 9: 45241-45245.

De Lima, R.P.; Da Silva, A.R.; Da Silva, A.P. (2021) soilphysics: An R package for simulation of soil compaction induced by agricultural field traffic. *SOIL and TILLAGE RESEARCH*, 206: 104824.

---

aggreg.stability      *Soil Aggregate-Size Distribution*

---

## Description

It calculates the mean weight diameter (MWD), the geometric mean diameter (GMD) and the soil aggregates size distribution per class based on the mass of the aggregates retained in each sieve from a total soil mass used for the soil aggregate stability test.

## Usage

```
aggreg.stability(sample.id = NA, dm.classes, aggre.mass)
```

## Arguments

sample.id	optional; a character vector containing the sample names.
dm.classes	a numeric vector containing the aggregates classes, in mm.
aggre.mass	a data.frame consisting of columns with soil aggregates mass (g) of each one of the corresponding dm.classes.

## Details

The user must arrange a data.frame with lines representing the samples and the columns representing the mass of the aggregates retained in each one of the meshes (corresponding to each size class) in the aggregate stability test.

$$\text{MWD} = \sum_{i=1}^n (x_i m_i)$$

where  $n$  is the number of aggregate size classes;  $x_i$  is mean size class  $i$ ; and  $m_i$  is the soil water-stable aggregate mass at the class =  $i$

$$\text{GMD} = e^{\sum_{i=1}^n \frac{m_i}{m} \ln(x_i)}$$

where  $n$  is the number of aggregate size classes;  $x_i$  is mean size class  $i$ ; and  $m_i$  is soil water-stable aggregate mass at the class =  $i$ ;  $m$  is the total soil mass used in the test.

### Value

A data.frame containing valor of MWD, GMD, total soil mass (total.mass) used in the aggregate stability test and the percentage of soil aggregate size distribution per class.

### Author(s)

Renato Paiva de Lima <renato\_agro\_@hotmail.com>

### References

W. Kemper, W. Chepil. (1965). Size distribution of aggregates. C. Black (Ed.). *Methods of Soil Analysis*, American Society Agronomy, Madison. pp. 499-510.

Yoder, R. A. (1936). A direct method of aggregate analysis of soils and a study of the physical nature of erosion losses. *Journal of the American Society of Agronomy*, 28:337-351.

### Examples

```
data(SoilAggregate)
classes <- c(3, 1.5, 0.75, 0.375, 0.178, 0.053)
aggreg.stability(sample.id = SoilAggregate[,1],
  dm.classes = classes, aggre.mass = SoilAggregate[, -1])

# End (not run)
```

---

aggreg.stability\_App *A shiny for Soil Aggregate-Size Distribution*

---

### Description

A shiny for Soil Aggregate-Size Distribution

**Usage**

```
aggreg.stability_App()
```

**Arguments**

Available on the interactive web page

**Value**

shiny app

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com>

**See Also**

[stressTraffic](#)

---

bulkDensity

*Soil Bulk Density Data Set*

---

**Description**

This data set refers to five observations of soil bulk density and soil moisture per sample. There are four soil samples.

**Usage**

```
data(bulkDensity)
```

**Format**

A data frame with 20 observations on the following 3 variables.

Id a factor with levels s1 s2 s3 s4, the 'ID' of each soil sample.

MOIS a numeric vector containing soil moisture values ( $\text{cm}^3 / \text{cm}^3$ ).

BULK a numeric vector containing soil bulk density values ( $\text{g} / \text{cm}^3$ ).

**Source**

Simulated data.

**Examples**

```
data(bulkDensity)
summary(bulkDensity)
```

---

`compaction`*Soil Compaction Data Set*

---

**Description**

This data set refers to physical soil variables related to soil compaction.

**Usage**

```
data(compaction)
```

**Format**

A data frame with 50 observations on the following 4 variables.

PR a numeric vector containing soil penetration resistance values (MPa).

BD a numeric vector containing soil bulk density values (g / cm<sup>3</sup>).

Mois a numeric vector containing soil moisture values (cm<sup>3</sup> / cm<sup>3</sup>).

PS a numeric vector containing soil preconsolidation stress values (kPa).

**Source**

Simulated data.

**Examples**

```
data(compaction)
summary(compaction)
```

---

`compressive_properties`*Estimation of compressive properties by Defossez et al. (2003)*

---

**Description**

It calculates the compressive parameters N and lambda using the pedo-transfer function from Defossez et al. (2003)

**Usage**

```
compressive_properties(water.content, soil=c("Loess", "Calcareous"))
```

**Arguments**

water.content a numeric vector containing the values of gravimetric water content, %  
 soil the soil group 'Loess' or 'Calcareous'. See examples

**Details**

In Defossez et al. (2003), the recompression index, kappa, was assumed as 0.0058 for both soil group.

**Value**

N the specific volume at  $p = 1kPa$ , N  
 CI the compression index, lambda

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com> Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Defossez, P., Richard, G., Boizard, H., & O'Sullivan, M. F., 2003. Modeling change in soil compaction due to agricultural traffic as function of soil water content. *Geoderma*, 116: 89–105.

**See Also**

[stressTraffic](#)

**Examples**

```
# EXAMPLE 1 - For Loess and Calcareous soil

water.content <- 25
compressive_properties(water.content=water.content, soil="Loess")
compressive_properties(water.content=water.content, soil="Calcareous")

# EXAMPLE 2 - For Loess soil

water.content <- seq(from=5,to=30,len=20)
out <- compressive_properties(water.content=water.content, soil="Loess")
plot(x=water.content ,y=out$N, ylab="N", xlab="Bulk density") # plot for N
plot(x=water.content ,y=out$CI, ylab="Lambda", xlab="Bulk density") # plot for compression index

# EXAMPLE 3 - For Calcareous soil

water.content <- seq(from=5,to=30,len=20)
out <- compressive_properties(water.content=water.content, soil="Calcareous")
plot(x=water.content ,y=out$N, ylab="N", xlab="Bulk density") # plot for N
plot(x=water.content ,y=out$CI, ylab="Lambda", xlab="Bulk density") # plot for compression index

# End (not run)
```



---

`compressive_properties2`*Estimation of compressive properties by Keller and Arvidsson (2007)*

---

**Description**

It calculates the compressive parameters N and lambda using the pedo-transfer function from Keller and Arvidsson (2007)

**Usage**

```
compressive_properties2(particle.density, bulk.density)
```

**Arguments**

`particle.density` a numeric vector containing the values of particle density,  $Mgm^{-3}$   
`bulk.density` a numeric vector containing the values of bulk density,  $Mgm^{-3}$

**Details**

In Keller and Arvidsson (2007), the recompression index, kappa, was found as 0.042 for all soil.

**Value**

N the specific volume at  $p = 1kPa$ , N  
CI the compression index, lambda

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com> Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Keller, T., Arvidsson, J., 2007. Compressive properties of some Swedish and Danish structured agricultural soils measured in uniaxial compression tests. *European Journal of Soil Science* , 58: 1373-1381.

**See Also**

[stressTraffic](#)

**Examples**

```
# EXAMPLE 1

compressive_properties2(particle.density=2.65, bulk.density=1.5)

# EXAMPLE 2

BD <- seq(from=1.2,to=1.8, by=0.01) # range of bulk density from 1.2 to 1.8
out <- compressive_properties2(particle.density=2.65, bulk.density=BD)

plot(x=BD,y=out$N, ylab="N", xlab="Bulk density") # for N
plot(x=BD,y=out$CI, ylab="Compression index (CI)", xlab="Bulk density") # for compression index

# End (not run)
```

---

compressive\_properties3

*Estimation of compressive properties by de Lima et al. (2018)*

---

**Description**

It calculates the compressive parameters N, lambda and kappa using the pedo-transfer function from de Lima et al. (2018)

**Usage**

```
compressive_properties3(bulk.density, matric.suction, soil=c("SandyLoam", "SandyClayLoam"))
```

**Arguments**

`bulk.density` a numeric vector containing the values of bulk density,  $Mgm^{-3}$   
`matric.suction` a numeric vector containing the values of matric suction, hPa  
`soil` the soil texture group 'SandyLoam' or 'SandyClayLoam'. See examples

**Details**

Pedo-transfer function developed under no-till condition. See de Lima et al. (2018)

**Value**

N the specific volume at  $p = 1kPa$ , N  
 CI the compression index, lambda  
 k the recompression index, kappa

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com> Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

## References

de Lima, R. P., da Silva, A. P., Giarola, N. F., da Silva, A. R., Rolim, M. M., Keller, T., 2018. Impact of initial bulk density and matric suction on compressive properties of two Oxisols under no-till. *Soil and Tillage Research*, 175: 168-177.

## See Also

[stressTraffic](#)

## Examples

```
# EXAMPLE 1

compressive_properties3(bulk.density=1.5, matric.suction=100, soil="SandyLoam")
compressive_properties3(bulk.density=1.5, matric.suction=100, soil="SandyClayLoam")

# EXAMPLE 2 for SandyLoam soil

matric.suction <- seq(from=30,to=1000,len=100)
out <- compressive_properties3(bulk.density=1.5,
  matric.suction=matric.suction, soil="SandyLoam")
plot(x=matric.suction,y=out$N, ylab="N",
  xlab="Matric suction (hPa)", log="x") # plot for N
# plot for lambda
plot(x=matric.suction,y=out$lambda, ylab="lambda",
  xlab="Matric suction (hPa)", log="x")
# plot for kappa
plot(x=matric.suction,y=out$k, ylab="kappa",
  xlab="Matric suction (hPa)", log="x")

# EXAMPLE 3 for SandyClayLoam soil

matric.suction <- seq(from=30,to=1000,len=100)
out <- compressive_properties3(bulk.density=1.5,
  matric.suction=matric.suction,
  soil="SandyClayLoam")
plot(x=matric.suction,y=out$N, ylab="N",
  xlab="Matric suction (hPa)", log="x") # plot for N
# plot for lambda
plot(x=matric.suction,y=out$lambda,
  ylab="lambda", xlab="Matric suction (hPa)", log="x")
# plot for kappa
plot(x=matric.suction,y=out$k,
  ylab="kappa", xlab="Matric suction (hPa)", log="x")

# End (not run)
```

---

`compressive_properties4`*Estimation of compressive properties by de Lima et al. (2020)*

---

**Description**

It calculates the compressive parameters N, lambda and kappa using the pedo-transfer function from de Lima et al. (2020)

**Usage**

```
compressive_properties4(matric.suction, soil=c("PloughLayer", "PloughPan"))
```

**Arguments**

`matric.suction` a numeric vector containing the values of matric suction, hPa.  
`soil` the soil compaction state 'PloughLayer' or 'PloughPan'. See the examples.

**Details**

Pedo-transfer function developed for a sandy loam soil texture. See de Lima et al. (2018)

**Value**

N the specific volume at  $p = 1kPa$ , N  
CI the compression index, lambda  
k the recompression index, kappa

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com> Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

de Lima, R. P., Rolim, M. M., da C. Dantas, D., da Silva, A. R., Mendonca, E. A., 2020. Compressive properties and least limiting water range of plough layer and plough pan in sugarcane fields. *Soil Use and Management*, 00: 1-12.

**See Also**

[stressTraffic](#)

**Examples**

```

# EXAMPLE 1

compressive_properties4(matric.suction=100, soil="PloughLayer")
compressive_properties4(matric.suction=100, soil="PloughPan")

# EXAMPLE 2 for "PloughLayer"

matric.suction <- seq(from=10,to=10000,len=100)
out <- compressive_properties4(matric.suction=matric.suction, soil="PloughLayer")
plot(x=matric.suction,y=out$N, ylab="N",
     xlab="Matric suction (hPa)", log="x") # plot for N
# plot for lambda
plot(x=matric.suction,y=out$lambda, ylab="lambda",
     xlab="Matric suction (hPa)", log="x")
# plot for kappa
plot(x=matric.suction,y=out$k, ylab="kappa",
     xlab="Matric suction (hPa)", log="x")

# EXAMPLE 3 for "PloughPan"

matric.suction <- seq(from=10,to=10000,len=100)
out <- compressive_properties4(matric.suction=matric.suction,
                             soil="PloughPan")

# plot for N
plot(x=matric.suction,y=out$N,
     ylab="N", xlab="Matric suction (hPa)", log="x")
# plot for lambda
plot(x=matric.suction,y=out$lambda,
     ylab="lambda", xlab="Matric suction (hPa)", log="x")
# plot for kappa
plot(x=matric.suction,y=out$k, ylab="kappa",
     xlab="Matric suction (hPa)", log="x")

# End (not run)

```

---

compressive\_properties5

*Estimation of compressive properties by O'Sullivan et al. (1999)*

---

**Description**

It calculates the compressive parameters N, lambda and kappa using the pedo-transfer function from O'Sullivan et al. (1999)

**Usage**

```
compressive_properties5(water.content, soil=c("SandyLoam","ClayLoam"))
```

**Arguments**

`water.content` a numeric vector containing the values of gravimetric water content, %  
`soil` the the soil texture group 'SandyLoam' or 'ClayLoam'. See examples.

**Details**

See O'Sullivan et al. (1999).

**Value**

`N` the specific volume at  $p = 1kPa$ , `N`  
`CI` the compression index, `lambda`  
`k` the recompression index, `kappa`

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com> Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

O'sullivan, M. F., Henshall, J. K., Dickson, J. W., 1999. A simplified method for estimating soil compaction. *Soil and Tillage Research*, 49: 325-335.

**See Also**

[stressTraffic](#)

**Examples**

```
# EXAMPLE 1
water.content <- 15
compressive_properties5(water.content=water.content, soil="SandyLoam")
compressive_properties5(water.content=water.content, soil="ClayLoam")

# EXAMPLE 2 - SandyLoam

water.content <- seq(from=5,to=20,len=20)
out <- compressive_properties5(water.content=water.content, soil="SandyLoam")
plot(x=water.content ,y=out$N,
     ylab="N", xlab="Bulk density") # plot for N
plot(x=water.content ,y=out$lambda,
     ylab="lambda", xlab="Bulk density") # plot for lambda
plot(x=water.content ,y=out$kappa,
     ylab="kappa", xlab="Bulk density") # plot for kappa
```

```
# EXAMPLE 3 - ClayLoam

water.content <- seq(from=10,to=25,len=20)
out <- compressive_properties5(water.content=water.content, soil="ClayLoam")
plot(x=water.content ,y=out$N,
     ylab="N", xlab="Bulk density") # plot for N
plot(x=water.content ,y=out$lambda,
     ylab="lambda", xlab="Bulk density") # plot for lambda
plot(x=water.content ,y=out$kappa,
     ylab="kappa", xlab="Bulk density") # plot for rkappa

# End (not run)
```

---

criticalmoisture

*Critical Moisture and Maximum Bulk Density*


---

### Description

Function to determine the soil Critical Moisture and the Maximum Bulk Density based on the Proctor (1933) compaction test. It estimates compaction curve by fitting a quadratic regression model.

### Usage

```
criticalmoisture(theta, Bd, samples = NULL, graph = TRUE, ...)
```

```
maxbulkdensity(theta, Bd, samples = NULL, graph = TRUE, ...)
```

### Arguments

theta	a vector containing the soil moisture values.
Bd	a vector containing the the soil bulk density values.
samples	optional; a vector indicating the multiple samples. Default is NULL (one sample). See details.
graph	logical; if TRUE (default), the soil compaction curve is plotted.
...	further graphical arguments.

### Details

If samples is ispecified, then it must has the same length of theta and Bd.

### Value

An object of class 'criticalmoisture', i.e., a matrix containing the quadratic model coefficients (rows 1 to 3), the R-squared (row 4), the sample size (row 5), the critical soil moisture (row 6) and the maximum bulk density (row 7), per sample.

**Note**

maxbulkdensity is just an alias of criticalmoisture.

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Proctor, E. R. (1933). Design and construction of rolled earth dams. *Eng. News Record*, 3: 245-284, 286-289, 348-351, 372-376.

Silva, A. P. et al. (2010). Indicadores da qualidade física do solo. In: Jong Van Lier, Q. (Ed). *Física do solo*. Vicosa (MG): Sociedade Brasileira de Ciencia do Solo. p.541-281.

**See Also**

[maxcurv](#)

**Examples**

```
# example 1 (1 sample)
mois <- c(0.083, 0.092, 0.108, 0.126, 0.135)
bulk <- c(1.86, 1.92, 1.95, 1.90, 1.87)
criticalmoisture(theta = mois, Bd = bulk)

# example 2 (4 samples)
data(bulkDensity)
attach(bulkDensity)
criticalmoisture(theta = MOIS, Bd = BULK, samples = Id)

# End (not run)
```

---

fitbusscher

*Self-starting Nls Busscher's (1990) Model for Soil Penetration Resistance*

---

**Description**

Function to self start the nonlinear Busscher's (1990) model for penetration resistance, i.e.,  $Pr = b_0 * (\theta^{b_1}) * (Bd^{b_2})$ . It creates initial estimates (by log-linearization) of the parameters  $b_0$ ,  $b_1$  and  $b_2$  and uses them to provide its least-squares estimates through [nls](#).

**Usage**

```
fitbusscher(Pr, theta, Bd, ...)
```



**Arguments**

Pr            a numeric vector containing penetration resistance values.  
theta        a numeric vector containing soil moisture values at which to evaluate the model.  
Bd            a numeric vector containing bulk density values at which to evaluate the model.  
...           further arguments to [nls](#).

**Value**

A [nls](#) output (see `help(nls)`).

**Author(s)**

Anderson Rodrigo da Silva <[anderson.agro@hotmail.com](mailto:anderson.agro@hotmail.com)>

**References**

Busscher, W. J. (1990). Adjustment of flat-tipped penetrometer resistance data to common water content. *Transactions of the ASAE*, 3:519-524.

**See Also**

[fitlbc](#), [nls](#), [summary.nls](#), [predict.nls](#), [Rsq](#)

**Examples**

```
data(compaction)
attach(compaction)
out <- fitbusscher(Pr = PR, theta = Mois, Bd = BD)
summary(out)
Rsq(out)

# 3D plot
X <- seq(min(Mois), max(Mois), len = 30) # theta
Y <- seq(min(BD), max(BD), len = 30) # Bd
f <- function(x, y) coef(out)[1] * (x^coef(out)[2]) * (y^coef(out)[3])
Z <- outer(X, Y, f)
persp(X, Y, Z,
      xlab = "Soil moisture",
      ylab = "Soil bulk density",
      zlab = "Penetration resistance",
      ticktype = "detailed",
      phi = 20, theta = 30)

# End (not run)
```

fitlbc

*Parameter Estimation of the Load Bearing Capacity Model***Description**

This function creates initial parameter estimates of the nonlinear Load Bearing Capacity (Dias Jr., 1994) model, i.e.,  $\sigma_P = 10^{(b_0 + b_1 * \theta)}$ , by using two methods: a `getInitial` method or a log-linearization. Then, it uses them to provide its least-squares estimates via [nls](#).

**Usage**

```
fitlbc(theta, sigmaP, ...)
```

**Arguments**

<code>theta</code>	a numeric vector containing soil moisture values.
<code>sigmaP</code>	a numeric vector containing values of soil preconsolidation stress.
<code>...</code>	further arguments to <a href="#">nls</a> .

**Value**

A [nls](#) object.

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Dias Junior, M. S. (1994). *Compression of three soils under longterm tillage and wheel traffic*. 1994. 114p. Ph.D. Thesis - Michigan State University, East Lansing.

**See Also**

[sigmaP](#), [fitbusscher](#), [maxcurv](#), [Rsq](#)

**Examples**

```
data(compaction)
attach(compaction)
out <- fitlbc(theta = Mois, sigmaP = PS)
summary(out)
Rsq(out)
curve(10^(coef(out)[1] + coef(out)[2]*x))

# End (not run)
```

**Description**

An interactive graphical adjustment of the soil water retention curve via van Genuchten's (1980) formula. The nonlinear least-squares estimates can be achieved taking the graphical initial values.

**Usage**

```
fitsoilwater(theta, x, xlab = NULL, ylab = NULL, ...)
```

**Arguments**

theta	a numeric vector containing the values of soil water content.
x	a numeric vector containing the matric potential values.
xlab	a label for the x axis; if is NULL, the label "Matric potential" is used.
ylab	a label for the y axis; if is NULL, the label "Soil water content" is used.
...	further graphical arguments; see <a href="#">par</a> .

**Value**

A plot of theta versus x and the curve of the current fitted model according to the adjusted parameters in an external interactive panel. Pressing the button "NLS estimates" a [nls](#) summary of the fitted model is printed on console whether convergence is achieved, otherwise a warning box of "No convergence" is shown.

**Author(s)**

Anderson Rodrigo da Silva <[anderson.agro@hotmail.com](mailto:anderson.agro@hotmail.com)>

**References**

Genuchten, M. T. van. (1980). A closed form equation for predicting the hydraulic conductivity of unsaturated soils. *Soil Science Society of America Journal*, 44:892-898.

**See Also**

[nls](#), [soilwater](#)

**Examples**

```
# Liu et al. (2011)
h <- c(0.001, 50.65, 293.77, 790.14, 992.74, 5065, 10130, 15195)
w <- c(0.5650, 0.4013, 0.2502, 0.2324, 0.2307, 0.1926, 0.1812, 0.1730)
fitsoilwater(w, h)

# End (not run)
```

---

`fitsoilwater2`*Interactive Estimation of the Groenevelt and Grant (2004) Model Parameters*

---

**Description**

An interactive graphical adjustment of the soil water retention curve via Groenevelt and Grant (2004) formula. The nonlinear least-squares estimates can be achieved taking the graphical initial values.

**Usage**

```
fitsoilwater2(theta, x, x0 = 6.653, xlab = NULL, ylab = NULL, ...)
```

**Arguments**

<code>theta</code>	a numeric vector containing the values of soil water content.
<code>x</code>	a numeric vector containing pF (pore water suction) values. See <a href="#">soilwater2</a> .
<code>x0</code>	the value of pF at which the soil water content becomes zero. The default is 6.653.
<code>xlab</code>	a label for the x axis; if is NULL, the label "pF" is used.
<code>ylab</code>	a label for the y axis; if is NULL, the label "Soil water content" is used.
<code>...</code>	further graphical arguments; see <a href="#">par</a> .

**Value**

A plot of `theta` versus `x` and the curve of the current fitted model according to the adjusted parameters in an external interactive panel. Pressing the button "NLS estimates" a [nls](#) summary of the fitted model is printed on console whether convergence is achieved, otherwise a warning box of "No convergence" is shown.

**Author(s)**

Anderson Rodrigo da Silva <[anderson.agro@hotmail.com](mailto:anderson.agro@hotmail.com)>

**References**

Groenevelt & Grant (2004). A new model for the soil-water retention curve that solves the problem of residual water contents. *European Journal of Soil Science*, 55:479-485.

**See Also**

[nls](#), [soilwater2](#), [soilwater](#)

## Examples

```
w <- c(0.417, 0.354, 0.117, 0.048, 0.029, 0.017, 0.007, 0)
pF <- 0:7
fitsoilwater2(w, pF)

# End (not run)
```

---

fitsoilwater3

*Interactive Estimation of the Dexter's (2008) Model Parameters*

---

## Description

An interactive graphical adjustment of the soil water retention curve through the Dexter's (2008) formula. The nonlinear least-squares estimates can be achieved taking the graphical initial values.

## Usage

```
fitsoilwater3(theta, x, xlab = NULL, ylab = NULL, ...)
```

## Arguments

theta	a numeric vector containing the values of soil water content.
x	a numeric vector containing the values of applied air pressure.
xlab	a label for the x axis; if is NULL, the label "pF" is used.
ylab	a label for the y axis; if is NULL, the label "Soil water content" is used.
...	further graphical arguments; see <a href="#">par</a> .

## Value

A plot of theta versus x and the curve of the current fitted model according to the adjusted parameters in an external interactive panel. Pressing the button "NLS estimates" a [nls](#) summary of the fitted model is printed on console whether convergence is achieved, otherwise a warning box of "No convergence" is shown.

## Author(s)

Anderson Rodrigo da Silva <[anderson.agro@hotmail.com](mailto:anderson.agro@hotmail.com)>

## References

Dexter et al. (2008). A user-friendly water retention function that takes account of the textural and structural pore spaces in soil. *Geoderma*, 143:243–253.

## See Also

[soilwater3](#), [nls](#), [fitsoilwater2](#)

**Examples**

```
# data extracted from Liu et al. (2011)
h <- c(0.001, 50.65, 293.77, 790.14, 992.74, 5065, 10130, 15195)
w <- c(0.5650, 0.4013, 0.2502, 0.2324, 0.2307, 0.1926, 0.1812, 0.1730)
fitsoilwater3(w, h)

# End (not run)
```

---

fitsoilwater4

*Self-starting Nls Power Models for Soil Water Retention*


---

**Description**

Function to self start the following nonlinear power models for soil water retention:

$$\theta = \exp(a + b * Bd)\psi^c$$

(Silva et al., 1994)

$$\theta = a\psi^c$$

(Ross et al., 1991)

where  $\theta$  is the soil water content.

fitsoilwater() creates initial estimates (by log-linearization) of the parameters a, b and c and uses them to provide its least-squares estimates through [nls](#).

**Usage**

```
fitsoilwater4(theta, psi, Bd, model = c("Silva", "Ross"))
```

**Arguments**

theta	a numeric vector containing values of soil water content.
psi	a numeric vector containing values of water potential (Psi).
Bd	a numeric vector containing values of dry bulk density.
model	a character; the model to be used for calculating the soil water content. It must be one of the two: "Silva" (default) or "Ross".

**Value**

A "nls" object containing the fitted model.

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

## References

Ross et al. (1991). Equation for extending water-retention curves to dryness. *Soil Science Society of America Journal*, 55:923-927.

Silva et al. (1994). Characterization of the least limiting water range of soils. *Soil Science Society of America Journal*, 58:1775-1781.

## See Also

[fitsoilwater4](#), [soilwater](#), [soilwater2](#), [soilwater3](#)

## Examples

```
data(skip1994)
# Example 1
ex1 <- with(skip1994,
  fitsoilwater4(theta = W, psi = h, model = "Ross"))
ex1
summary(ex1)

# Example 2
ex2 <- with(skip1994,
  fitsoilwater4(theta = W, psi = h, Bd = BD, model = "Silva"))
ex2
summary(ex2)

# Not run
```

---

fitsoilwater5

*Interactive Estimation of the Modified van Genuchten's Model Parameters*

---

## Description

An interactive graphical adjustment of the soil water retention curve via the van Genuchten's formula, modified by Pierson and Mulla (1989). The nonlinear least-squares estimates can be achieved taking the graphical initial values. It may be useful to estimate the parameters needed in the high-energy-moisture-characteristics (HEMC) method, which is used to analyze the aggregate stability.

## Usage

```
fitsoilwater5(theta, x, theta_S, xlab = NULL, ylab = NULL, ...)
```

## Arguments

theta	a numeric vector containing the values of soil water content.
x	a numeric vector containing the matric potential values.
theta_S	an offset; a value for the parameter theta_S, the water content at saturation. See details.

xlab            a label for the x axis; if is NULL, the label "Matric potential" is used.  
 ylab            a label for the y axis; if is NULL, the label "Soil water content" is used.  
 ...             further graphical arguments; see [par](#).

### Details

The parameter `theta_S` must be passed as an argument. It is recommended to consider it as the highest water content value in the data set or the water content at saturation.

### Value

A plot of `theta` versus `x` and the curve of the current fitted model according to the adjusted parameters in an external interactive panel. Pressing the button "NLS estimates" a [nls](#) summary of the fitted model is printed on console whether convergence is achieved, otherwise a warning box of "No convergence" is shown.

### Author(s)

Anderson Rodrigo da Silva <[anderson.agro@hotmail.com](mailto:anderson.agro@hotmail.com)>

### References

Pierson, F.B.; Mulla, D.J. (1989) An Improved Method for Measuring Aggregate Stability of a Weakly Aggregated Loessial Soil. *Soil Sci. Soc. Am. J.*, 53:1825–1831.

### See Also

[nls](#), [soilwater5](#)

### Examples

```
h <- seq(0.1, 40, by = 2)
w <- c(0.735, 0.668, 0.635, 0.612, 0.559, 0.462, 0.369, 0.319, 0.296, 0.282,
0.269, 0.256, 0.249, 0.246, 0.239, 0.236, 0.229, 0.229, 0.226, 0.222)
plot(w ~ h)

# suggestions of starting values: thetaR = 0.35, alpha = 0.1, n = 10,
# b0 = 0.02, b1 = -0.0057, b2 = 0.00004 (Not run)

fitsoilwater5(theta = w, x = h, theta_S = 0.70)

# End (Not run)
```



**Description**

An accessorial function to convert an object of class 'function' to an object of class 'formula'.

**Usage**

```
fun2form(fun, y = NULL)
```

**Arguments**

fun	a object of class 'function'. It must be a one-line-written function, with no curly braces "{}".
y	optional; a character defining the left side of the formula, $y = \text{fun}()$ .

**Value**

An object of class [formula](#).

**Warning**

Numerical values into fun with three or more digits may cause miscalculation.

**Author(s)**

Anderson Rodrigo da Silva <[anderson.agro@hotmail.com](mailto:anderson.agro@hotmail.com)>

**See Also**

[function](#), [formula](#)

**Examples**

```
g <- function(x) Asym * exp(-b2 * b3 ^ x) # Gompertz Growth Model
fun2form(g, "y")

# f1 <- function(w) {exp(w)} # error
# fun2form(f1, "x")
f2 <- function(w) exp(w) # ok
fun2form(f2, "x")

# End (not run)
```

---

`getInitiallbc`*Get Initial Parameter Estimates for the Load Bearing Capacity Model*

---

**Description**

This is a `getInitial` function that evaluates initial parameter estimates for the Load Bearing Capacity model via `SSLbc`.

**Usage**

```
getInitiallbc(theta, sigmaP)
```

**Arguments**

<code>theta</code>	a numeric vector containing values of soil moisture.
<code>sigmaP</code>	a numeric vector containing values of preconsolidation stress.

**Value**

A numeric vector containing the estimates of the parameters `b0` and `b1`.

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Dias Junior, M. S. (1994). *Compression of three soils under longterm tillage and wheel traffic*. 1994. 114p. Ph.D. Thesis - Michigan State University, East Lansing.

**See Also**

`getInitial`, `SSLbc`, `nls`, `sigmaP`

**Examples**

```
data(compaction)
attach(compaction)
getInitiallbc(theta = Mois, sigmaP = PS)

# End (not run)
```

---

 hemc

*High-Energy-Moisture-Characteristics Aggregate Stability*


---

### Description

A function to determine the modal suction, volume of drainable pores, structural index and stability ratio using the high-energy-moisture-characteristics (HEMC) method by Pierson & Mulla (1989), which is used to analyze the aggregate stability. Before using `hemc()`, the user may estimate the parameters of the Modified van Genuchten's Model through the function `fitsoilwater5()`.

### Usage

```
hemc(x, theta_R, theta_S, alpha, n, b1, b2,
     graph = TRUE, from = 1, to = 30,
     xlab = expression(Psi ~ (J~kg^{-1})),
     ylab = expression(d ~ theta/d ~ Psi), ...)
```

### Arguments

<code>x</code>	a vector containing matric potential values.
<code>theta_R</code>	a numeric vector of length two containing the parameter values in the following orde: fast and slow.
<code>theta_S</code>	a numeric vector of length two containing the parameter values in the following orde: fast and slow.
<code>alpha</code>	a numeric vector of length two containing the parameter values in the following orde: fast and slow.
<code>n</code>	a numeric vector of length two containing the parameter values in the following orde: fast and slow.
<code>b1</code>	a numeric vector of length two containing the parameter values in the following orde: fast and slow.
<code>b2</code>	a numeric vector of length two containing the parameter values in the following orde: fast and slow.
<code>graph</code>	logical; if TRUE (default), a graphical solution is shown).
<code>from</code>	the lower limit for the x-axis
<code>to</code>	the lower limit for the x-axis
<code>xlab</code>	a label for the x-axis
<code>ylab</code>	a label for the y-axis
<code>...</code>	further graphical arguments

### Value

A list of a two objects: 1) a matrix containing the Modal Suction, the Volume od Drainable Pores (VDP) and the Structural Index for both, fast and slow wetting; and 2) the value of Stability Ratio.

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Pierson, F.B.; Mulla, D.J. (1989). An Improved Method for Measuring Aggregate Stability of a Weakly Aggregated Loessial Soil. *Soil Sci. Soc. Am. J.*, 53:1825–1831.

**See Also**

[fitsoilwater5](#)

**Examples**

```
hemc(x = seq(1, 30), theta_R = c(0.27, 0.4), theta_S = c(0.65, 0.47),
alpha = c(0.1393, 0.0954), n = c(6.37, 7.47),
b1 = c(-0.008421, -0.011970), b2 = c(0.0001322, 0.0001552))

# End (Not run)
```

---

hydraulicCutOff	<i>The matric potential at the point of hydraulic cut-off obtained from DE (Dexter et al., 2008) and GG (Groenevelt &amp; Grant, 2004) water retention curves.</i>
-----------------	--

---

**Description**

The pore water suction at the point of hydraulic cut-off occurs at the point where the residual water content, obtained from Dexter et al. (2008), intercepts with the Groenevelt & Grant (2004) retention curve.

**Usage**

```
hydraulicCutOff(theta_R, k0, k1, n, x0 = 6.653)
```

**Arguments**

theta_R	a parameter that represents the residual water content at the the Dexter's (2008) Water Retention Model.
k0	a parameter value, extracted from the water retention curve based on the Groenevelt & Grant (2004) formula.
k1	a parameter value, extracted from the water retention curve based on the Groenevelt & Grant (2004) formula.
n	a parameter value, extracted from the water retention curve based on the Groenevelt & Grant (2004) formula.
x0	the value of pF (pore water suction) at which the soil water content becomes zero. The default is 6.653.

**Value**

The water suction at the point of hydraulic cut-off.

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Dexter, A.R.; Czyz, E.A.; Richard, G.; Reszkowska, A. (2008). A user-friendly water retention function that takes account of the textural and structural pore spaces in soil. *Geoderma*, 143:243–253.

Groenevelt, P.H.; Grnat, C.D. (2004). A new model for the soil-water retention curve that solves the problem of residual water contents. *European Journal of Soil Science*, 55:479–485.

**See Also**

[fitsoilwater2](#), [fitsoilwater3](#)

**Examples**

```
# Dexter et al. (2012), Table 4A
hydraulicCutOff(0.1130, 6.877, 0.6508, 1.0453)
hydraulicCutOff(0.1122, 12.048, 0.4338, 2.0658)

# End (not run)
```

---

hydraulicCutOff2	<i>The matric potential at the point of hydraulic cut-off using the point of maximum curvature of DE (Dexter et al. 2008) water retention curve.</i>
------------------	--

---

**Description**

The pore water suction at the point of hydraulic cut-off occurs at the point where the residual water content, obtained from Dexter et al. (2008), intercepts with the Groenevelt & Grant (2004) retention curve. This function calculates the Hydraulic Cut-Off using the point of maximum curvature of the DE (Dexter et al. 2008) curve.

**Usage**

```
hydraulicCutOff2(theta_R, a1, a2, p1, p2, graph = FALSE, ...)
```

**Arguments**

theta_R	the residual water content from Dexter's (2008) water retention curve (g/g).
a1	a water content parameter from Dexter's (2008) water retention curve (g/g).
a2	a water content parameter from Dexter's (2008) water retention curve (g/g).
p1	a matric potential parameter from Dexter's (2008) water retention curve (hPa).
p2	a matric potential parameter from Dexter's (2008) water retention curve (hPa).
graph	logical; if TRUE a graphical solution with the maximum curvature point is displayed.
...	further graphical arguments. See <a href="#">par</a>

**Details**

The arguments are the fitting parameters from Dexter's (2008) water retention curve, which can be fitted using [fitsoilwater3](#). Further examples of how to use these parameters are given in Dexter et al. (2012).

**Value**

A `data.frame` containing the values of matric potential (hPa), pF and water content (w) at the hydraulic cut-off (hco) point.

**Author(s)**

Renato Paiva de Lima <[renato\\_agro@hotmail.com](mailto:renato_agro@hotmail.com)>

**References**

- Dexter, A.R.; Czyz, E.A.; Richard, G.; Reszkowska, A. (2008). A user-friendly water retention function that takes account of the textural and structural pore spaces in soil. *Geoderma*, 143:243–253.
- Dexter, A.R., Czyz, E.A., Richard, G. (2012). Equilibrium, non-equilibrium and residual water: consequences for soil water retention. *Geoderma*, 177:63–71.

**See Also**

[hydraulicCutOff](#), [fitsoilwater3](#)

**Examples**

```
# Example 1: soils from Dexter et al. (2012), Table 4

hydraulicCutOff2(theta_R=0.1130,a1=0.0808,a2=0.0576,p1=4043.2,p2=269.1,
                 graph = TRUE, ylim=c(-0.05,0.15)) # Soil 1

hydraulicCutOff2(theta_R=0.0998,a1=0.1456,a2=0.0162,p1=3156.0,p2=71.51,
                 graph = TRUE, ylim=c(-0.20,0.30)) # Soil 4
```

```

hydraulicCutOff2(theta_R=0.0709,a1=0.0195,a2=0.1794,p1=4467.5,p2=1395.5,
                 graph = TRUE, ylim=c(-0.20,0.30)) # Soil 7

hydraulicCutOff2(theta_R=0.0359,a1=0.1014,a2=0.0459,p1=1282.4,p2=56.93,
                 graph = TRUE, ylim=c(-0.10,0.20)) # Soil 10

hydraulicCutOff2(theta_R=0.0736,a1=0.0522,a2=0.0321,p1=3516.2,p2=90.54,
                 graph = TRUE, ylim=c(-0.05,0.15)) # Soil 14

# Example 2:

# Fitting the water retention curve through the Dexter's (2008) curve
h <- c(0.001, 50.65, 293.77, 790.14, 992.74, 5065, 10130, 15195)
w <- c(0.5650, 0.4013, 0.2502, 0.2324, 0.2307, 0.1926, 0.1812, 0.1730)
if (interactive()) {
  fitsoilwater3(theta=w, x=h)
}

# Using the fitted parameter
hydraulicCutOff2(theta_R=0.1738,a1=0.07505,a2=0.316,p1=3673,p2=70.38,
                 graph = TRUE, ylim=c(-0.40,0.60))

# End (not run)

```

---

iwc

*Integral Water Capacity (IWC)*


---

### Description

Quantifying the soil water availability for plants through the IWC approach. The theory was based on the work of Groenevelt et al. (2001), Groenevelt et al. (2004) and Asgarzadeh et al. (2014), using the van Genuchten-Mualem Model for estimation of the water retention curve and a simple power model for penetration resistance. The salinity effect on soil available water is also implemented here, according to Groenevelt et al. (2004).

### Usage

```

iwc(theta_R, theta_S, alpha, n, a, b, hos = 0,
    graph = TRUE,
    xlab = "Matric head (cm)",
    ylab = expression(cm^-1),
    xlim1 = NULL,
    xlim2 = NULL,
    xlim3 = NULL,
    ylim1 = NULL,
    ylim2 = NULL,
    ylim3 = NULL,
    col12 = c("black", "blue", "red"),

```

```
col3 = c("orange", "black"),
lty12 = c(1, 3, 2),
lty3 = c(2, 1), ...)
```

### Arguments

theta_R	the residual water content ( $m^3 m^{-3}$ ); a numeric parameter from van Genuchten's model; see details.
theta_S	the water content at saturation ( $m^3 m^{-3}$ ); a numeric parameter from van Genuchten's model; see details.
alpha	a scale parameter from van Genuchten's model; see details.
n	a shape parameter from van Genuchten's model; see details.
a	a parameter of the soil penetration resistance model; see details.
b	a parameter of the soil penetration resistance model; see details.
hos	optional; the value of osmotic head of the saturated soil extract (cm). Used only if one is concerned about the salinity effects on the water available for plants. Default is zero. See Groenevelt et al. (2004) for more details.
graph	logical; if TRUE (default), graphics for both dry and wet range are built.
xlab	a label for x-axis.
ylab	a label for y-axis.
xlim1, xlim2, xlim3	the x limits (x1, x2) of each plot. See <a href="#">plot.default</a> .
ylim1, ylim2, ylim3	the y limits (y1, y2) of each plot. See <a href="#">plot.default</a> .
col12	a vector of length 3 containing the color of each line of the first two plots. See <a href="#">par</a> .
col3	a vector of length 2 containing the color of each line of the third plot. See <a href="#">par</a> .
lty12	a vector of length 3 containing the line types for the first two plots. See <a href="#">par</a> .
lty3	a vector of length 2 containing the line types for the third plot. See <a href="#">par</a> .
...	further graphical parameters. See <a href="#">par</a> .

### Details

The parameters of the van Genuchten-Mualem Model can be estimated through the function `fitsoilwater()`. The soil penetration resistance model is given by:  $PR = a * h^b$ , where  $h$  is the soil water content and  $a$  and  $b$  are the fitting parameters.

### Value

A table containing each integration of IWC (integral water capacity, in m/m) and EI (integral energy calculation, in J/kg).

### Author(s)

Anderson Rodrigo da Silva <[anderson.agro@hotmail.com](mailto:anderson.agro@hotmail.com)>



## References

- Asgarzadeh, H.; Mosaddeghi, M.R.; Nikbakht, A.M. (2014) SAWCal: A user-friendly program for calculating soil available water quantities and physical quality indices. *Computers and Electronics in Agriculture*, 109:86–93.
- Groenevelt, P.H.; Grant, C.D.; Semetsa, S. (2001) A new procedure to determine soil water availability. *Australian Journal Soil Research*, 39:577–598.
- Groenevelt, P.H., Grant, C.D., Murray, R.S. (2004) On water availability in saline soils. *Australian Journal Soil Research*, 42:833–840.

## See Also

[soilwater](#), [fitsoilwater](#), [llwr](#)

## Examples

```
# example 1 (Fig 1b, Asgarzadeh et al., 2014)
iwc(theta_R = 0.0160, theta_S = 0.4828, alpha = 0.0471, n = 1.2982,
a = 0.2038, b = 0.2558, graph = TRUE)

# example 2 (Table 1, Asgarzadeh et al., 2014)
iwc(theta_R = 0.166, theta_S = 0.569, alpha = 0.029, n = 1.308,
a = 0.203, b = 0.256, graph = TRUE)

# example 3: evaluating the salinity effect
iwc(theta_R = 0.166, theta_S = 0.569, alpha = 0.029, n = 1.308,
a = 0.203, b = 0.256, hos = 200, graph = TRUE)

# End (Not run)
```

---

Kr\_h

*Unsaturated Hydraulic Conductivity*

---

## Description

A closed-form analytical expressions for calculating the relative unsaturated hydraulic conductivity as a function of soil water tension (h) based on van Genuchten's water retention curve.

## Usage

```
Kr_h(Ks, alpha, n, h, f=0.5)
```

## Arguments

Ks	Saturated hydraulic conductivity (e.g. cm/day).
alpha	The scale parameter of the van Genuchten's model (hPa <sup>-1</sup> ).
n	The shape parameter in van Genuchten's formula.
h	The water tension (hPa).
f	The pore-connectivity parameter. Default 0.5 [Mualem, 1976].

**Value**

numeric, the value of unsaturated hydraulic conductivity.

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com>

**References**

Guarracino, L. (2007). Estimation of saturated hydraulic conductivity Ks from the van Genuchten shape parameter alpha. *Water Resources Research*, 43(11).

Van Genuchten, M. T. (1980). A closed-form equation for predicting the hydraulic conductivity of unsaturated soils 1. *Soil Science Society of America Journal* 44(5):892-898.

**Examples**

```
# EXAMPLE 1
Kr_h(Ks = 1.06*10^2, alpha = 0.048, n = 1.5,h=100, f=0.5)

# EXAMPLE 2
x <- seq(log10(1), log10(1000),len=100)
h <- 10^x
y <- Kr_h(Ks = 1.06*10^2, alpha = 0.048, n = 1.5,h=h, f=0.5)
plot(x=h,y=y, log="yx", xlab="h (hPa)", yaxt='n',
      ylab="", ylim=c(0.001,100), xlim=c(1,10000))
mtext(expression(K[r] ~ (cm~d^-1)), 2, line=2)
ax <- c(0.001, 0.01, 0.1, 1, 10, 100)
axis(2,at=ax, labels=ax)

# End (not run)
```

---

Kr\_theta

*Unsaturated Hydraulic Conductivity as a function of water content*

---

**Description**

A closed-form analytical expressions for calculating the relative unsaturated hydraulic conductivity as a function of soil water content based on van Genuchten's water retention curve.

**Usage**

```
Kr_theta(theta, thetaS, thetaR, alpha, n, Ks, f=0.5)
```

**Arguments**

theta	The volumetric water content ( $m^3/m^3$ ).
thetaS	The volumetric water content at the saturation ( $m^3/m^3$ ).
thetaR	The volumetric residual water content ( $m^3/m^3$ ).
alpha	The scale parameter of the van Genuchten's model ( $hPa^{-1}$ ).
n	The shape parameter in van Genuchten's formula.
Ks	Saturated hydraulic conductivity (e.g. cm/day).
f	The pore-connectivity parameter. Default 0.5 [Mualem, 1976].

**Value**

numeric, the value of unsaturated hydraulic conductivity.

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com>

**References**

- Guarracino, L. (2007). Estimation of saturated hydraulic conductivity Ks from the van Genuchten shape parameter alpha. *Water Resources Research*, 43(11).
- Van Genuchten, M. T. (1980). A closed-form equation for predicting the hydraulic conductivity of unsaturated soils. *Soil Science Society of America Journal* 44(5):892-898.
- Mualem, Y. (1976). A new model for predicting the hydraulic conductivity of unsaturated porous media. *Water Resour. Res.* 43(11): 513-522,

**Examples**

```
# EXEMPLE 1
Kr_theta(theta=0.45,thetaS=0.5,thetaR=0.15,
         alpha = 0.02, n = 2, Ks = 1.06*10^2, f=0.5)

# EXEMPLE 2
thetaS <- 0.50
thetaR <- 0.15
theta <- seq(thetaS, thetaR, len=50)
y <- Kr_theta(theta=theta,thetaS=thetaS,thetaR=thetaR,
             alpha = 0.02, n = 2, Ks = 1.06*10^2, f=0.5)

# Just for this example, we are removing the "0" value
# for plotting the graph in log scale, since log10(0) results in "-Inf"
Kr <- y[-50]
w <- theta[-50]

plot(x=w,y=Kr,xlab=expression(theta~(m^3~m^-3)),
     ylim=c(0.001,100), log="y", yaxt='n',
     ylab="", xlim=c(0.15,0.50))
mtext(expression(K[r] ~ (cm~d^-1)), 2, line=2)
```

```
ax <- c(0.001, 0.01, 0.1, 1, 10, 100)
axis(2,at=ax, labels=ax)
# End (not run)
```

---

liquidlimit

*Soil Liquid Limit*


---

### Description

Function to determine the soil Liquid Limit by using the Sowers (1965) method.

$$LL = \theta * (n/25)^{0.12}$$

### Usage

```
liquidlimit(theta, n)
```

### Arguments

theta	the soil moisture value corresponding to n drops.
n	the number of drops.

### Value

The soil moisture value corresponding to the Liquid Limit.

### Author(s)

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

### References

- Sowers, G. F. (1965). Consistency. In: BLACK, C.A. (Ed.). *Methods of soil analysis*. Madison: American Society of Agronomy. Part 1, p.391-399. (Agronomy, 9).
- Sowers, G. F. (1965). Consistency. In: KLUTE, A. (Ed.). 2 ed. *Methods of soil analysis*. Madison: American Society of Agronomy. Part 1, p.545-566.

### See Also

[criticalmoisture](#)

**Examples**

```
liquidlimit(theta = 0.34, n = 22)

M <- c(0.34, 0.29, 0.27, 0.25, 0.20)
N <- c(22, 24, 25, 26, 28)
liquidlimit(theta = M, n = N)

# End (not run)
```

llwr

*Least Limiting Water Range (LLWR)***Description**

Graphical solution for the Least Limiting Water Range and parameter estimation of the related water retention and penetration resistance curves. A summary containing standard errors and statistical significance of the parameters is also given.

**Usage**

```
llwr(theta, h, Bd, Pr,
      particle.density, air,
      critical.PR, h.FC, h.WP,
      water.model = c("Silva", "Ross"),
      Pr.model = c("Busscher", "noBd"),
      pars.water = NULL, pars.Pr = NULL,
      graph = TRUE, graph2 = TRUE,
      xlab = expression(Bulk~Density~(Mg~m^{-3})),
      ylab = expression(theta~(m^3~m^{-3})),
      main = "Least Limiting Water Range", ...)
```

**Arguments**

theta	a numeric vector containing values of volumetric water content ( $m^3 m^{-3}$ ).
h	a numeric vector containing values of matric head (cm, Psi, MPa, kPa, ...).
Bd	a numeric vector containing values of dry bulk density ( $Mg m^{-3}$ ). Note that Bd can also be a vector of length 1. See details.
Pr	a numeric vector containing values of penetration resistance (MPa, kPa, ...).
particle.density	the value of the soil particle density ( $Mg m^{-3}$ ).
air	the value of the limiting volumetric air content ( $m^3 m^{-3}$ ).
critical.PR	the value of the critical soil penetration resistance.
h.FC	the value of matric head at the field capacity (cm, MPa, kPa, hPa, ...).
h.WP	the value of matric head at the wilting point (cm, MPa, kPa, hPa, ...).

water.model	a character; the model to be used for calculating the soil water content. It must be one of the two: "Silva" (default) or "Ross". See details.
Pr.model	a character; the model to be used to predict soil penetration resistance. It must be one of the two: "Busscher" (default) or "noBd". See details.
pars.water	optional; a numeric vector containing the estimates of the three parameters of the soil water retention model employed. If NULL (default), llwr() estimates them using a Newton-Raphson algorithm. See details.
pars.Pr	optional; a numeric vector containing estimates of the three parameters of the model proposed by Busscher (1990) for the functional relationship among Pr, theta and Bd. If NULL (default), llwr() estimates them using a Newton-Raphson algorithm. Moreover, if Pr.model = "noBd", then the third value is considered to be null.
graph	logical; if TRUE (default) a graphical solution for the Least Limiting Water Range is plotted.
graph2	logical; if TRUE (default) a line of the Least Limiting Water Range as a function of bulk density is plotted. If graph = FALSE, then llwr() will automatically consider graph2 = FALSE too.
xlab	a title for the x axis; the default is <i>Bulk Density</i> ( $Mg\ m^{-3}$ ).
ylab	a title for the y axis; the default is $\theta$ ( $m^3\ m^{-3}$ ).
main	a main title for the graphic; the default is "Least Limiting Water Range"
...	further graphical arguments.

### Details

The numeric vectors theta, h, Bd and Pr are supposed to have the same length, and their values should have appropriate unit of measurement. For fitting purposes, it is not advisable to use vectors with less than five values. It is possible to calculate the LLWR for a specific (unique) value of bulk density. In This case, Bd should be a vector of length 1 and, therefore, it is not possible to fit the models "Silva" and "Busscher", for water content and penetration resistance, respectively.

The model employed by Silva et al. (1994) for the soil water content ( $\theta$ ) as a function of the soil bulk density ( $\rho$ ) and the matric head (h) is:

$$\theta = \exp(a + b\rho)h^c$$

The model proposed by Ross et al. (1991) for the soil water content ( $\theta$ ) as a function of the matric head (h) is:

$$\theta = ah^c$$

The penetration resistance model, as presented by Busscher (1990), is given by

$$Pr = b0 * (\theta^{b1}) * (\rho^{b2})$$

If the argument Bd receives a single value of bulk density, then llwr() fits the following simplified model (option noBd):

$$Pr = b0 * \theta^{b1}$$

**Value**

A list of	
limiting.theta	a $n \times 4$ matrix containing the limiting values of water content for each input value of bulk density at the volumetric air content (thetaA), penetration resistance (thetaPR), field capacity (thetaFC) and wilting point (thetaWP).
pars.water	a "nls" object or a numeric vector containing estimates of the three parameters of the model employed by Silva et al. (1994) for the functional relationship among theta, Bd and h.
r.squared.water	a "Rsq" object containing the pseudo and the adjusted R-squared for the water model.
pars.Pr	a "nls" object or a numeric vector containing estimates of the three parameters of the penetration resistance model.
r.squared.Pr	a "Rsq" object containing the pseudo and the adjusted R-squared for the penetration resistance model.
area	numeric; the value of the shaded (LLWR) area. Calculated only when Bd is a vector of length > 1.
LLWR	numeric; the value of LLWR ( $m^3 m^{-3}$ ) corresponding to Bd. Calculated only when Bd is a single value.

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

- Busscher, W. J. (1990). Adjustment of flat-tipped penetrometer resistance data to common water content. *Transactions of the ASAE*, 3:519-524.
- Leao et al. (2005). An Algorithm for Calculating the Least Limiting Water Range of Soils. *Agronomy Journal*, 97:1210-1215.
- Leao et al. (2006). Least limiting water range: A potential indicator of changes in near-surface soil physical quality after the conversion of Brazilian Savanna into pasture. *Soil & Tillage Research*, 88:279-285.
- Ross et al. (1991). Equation for extending water-retention curves to dryness. *Soil Science Society of America Journal*, 55:923-927.
- Silva et al. (1994). Characterization of the least limiting water range of soils. *Soil Science Society of America Journal*, 58:1775-1781.

**See Also**

[fitbusscher](#)

## Examples

```
# Example 1 - part of the data set used by Leao et al. (2005)
data(skp1994)
ex1 <- with(skp1994,
  llwr(theta = W, h = h, Bd = BD, Pr = PR,
  particle.density = 2.65, air = 0.1,
  critical.PR = 2, h.FC = 100, h.WP = 15000))
ex1

# Example 2 - specifying the parameters (Leao et al., 2005)
a <- c(-0.9175, -0.3027, -0.0835) # Silva et al. model of water content
b <- c(0.0827, -1.6087, 3.0570) # Busscher's model
ex2 <- with(skp1994,
  llwr(theta = W, h = h, Bd = BD, Pr = PR,
  particle.density = 2.65, air = 0.1,
  critical.PR = 2, h.FC = 0.1, h.WP = 1.5,
  pars.water = a, pars.Pr = b))
ex2

# Example 3 - specifying a single value for Bd
ex3 <- with(skp1994,
  llwr(theta = W, h = h, Bd = 1.45, Pr = PR,
  particle.density = 2.65, air = 0.1,
  critical.PR = 2, h.FC = 100, h.WP = 15000))
ex3

# End (not run)
```

---

llwrPTF

*Least Limiting Water Range (LLWR) Using Pedo-Transfer Functions*


---

## Description

It calculates Least Limiting Water Range (LLWR) using pedo-transfer functions in according to Silva & Kay (1997) and Silva et al. (2008), for Canadian and Brazilian soils, respectively.

## Usage

```
llwrPTF(air, critical.PR, h.FC, h.WP, p.density, Bd, clay.content, org.carbon = NULL)
```

## Arguments

air	the value of the limiting volumetric air content, $m^3m^{-3}$
critical.PR	the value of the critical soil penetration resistance, MPa
h.FC	the value of matric suction at the field capacity, hPa
h.WP	the value of matric suction at the wilting point, hPa
p.density	the value of the soil particle density, $Mgm^{-3}$



Bd	a numeric vector containing values of dry bulk density, $Mgm^{-3}$ . Note that Bd can also be a vector of length 1.
clay.content	a numeric vector containing values of clay content to each bulk density, %
org.carbon	a numeric vector containing values of organic carbon to each bulk density, %, for Canadian soils. Default is 2%. See details.

### Details

Note that org.carbon is only required for Canadian soil. If it is not passed, LLWR for Canadian soil is calculated with 2% of organic carbon.

### Value

A list of

LLWR.B	LLWR for Brazilian soils
LLWR.C	LLWR for Canadian soils

### Author(s)

Renato Paiva de Lima <renato\_agro\_@hotmail.com>  
 Anderson Rodrigo da Silva <anderson.agro@hotmail.com>  
 Alvaro Pires da Silva <apisilva@usp.br>

### References

- Keller, T; Silva, A.P.; Tormena, C.A.; Giarola, N.B.F., Cavalieri, K.M.V., Stettler, M.; Arvidsson, J. 2015. SoilFlex-LLWR: linking a soil compaction model with the least limiting water range concept. *Soil Use and Management*, 31:321-329.
- Silva, A.P.; Kay, B.D. 1997. Estimating the least limiting water range of soil from properties and management. *Soil Science Society of America Journal*, 61:877-883.
- Silva, A.P., Kay, B.D.; Perfect, E. 1994. Characterization of the least limiting water range. *Soil Science Society of America Journal*, 61:877-883.
- Silva, A.P., Tormena, C.A., Jonez, F.; Imhoff, S. 2008. Pedotransfer functions for the soil water retention and soil resistance to penetration curves. *Revista Brasileira de Ciencia do Solo*, 32:1-10.

### Examples

```
# EXEMPLE 1 (for Brazilian Soils)
llwrPTF(air=0.1,critical.PR=2, h.FC=100, h.WP=15000,p.density=2.65,
        Bd=c(1.2,1.3,1.4,1.5,1.35),clay.content=c(30,30,35,38,40))

# EXEMPLE 2 (for Canadian Soils)
llwrPTF(air=0.1,critical.PR=2, h.FC=100, h.WP=15000,p.density=2.65,
        Bd=c(1.2,1.3,1.4),clay.content=c(30,30,30), org.carbon=c(1.3,1.5,2))

# EXEMPLE 3 (combining it with soil stress)
stress <- stressTraffic(inflation.pressure=200,
```

```

recommended.pressure=200,
tyre.diameter=1.8,
tyre.width=0.4,
wheel.load=4000,
conc.factor=c(4,5,5,5,5,5),
      layers=c(0.05,0.1,0.3,0.5,0.7,1),
plot.contact.area = FALSE)

stress.p <- stress$Stress$sigma_mean
layers <- stress$Stress$Layers
n <- length(layers)
def <- soilDeformation(stress = stress.p,
      p.density = rep(2.67,n),
      iBD = rep(1.55,n),
      N = rep(1.9392,n),
      CI = rep(0.06037,n),
      k = rep(0.00608,n),
      k2 = rep(0.01916,n),
      m = rep(1.3,n),graph=TRUE,ylim=c(1.4,1.8))

# Graph LLWR, considering Brazilian soils
plot(x = 1, y = 1,
      xlim=c(0,0.2),ylim=c(1,0),xaxt = "n",
      ylab = "Soil Depth",xlab = "", type="l", main="")
axis(3)
mtext("LLWR",side=3,line=2.5)

i.LLWR <- llwrPTF(air=0.1,critical.PR=2, h.FC=100,
      h.WP=15000,p.density=2.65,
      Bd=def$iBD,clay.content=rep(20,n))
f.LLWR <- llwrPTF(air=0.1,critical.PR=2, h.FC=100,
      h.WP=15000,p.density=2.65,
      Bd=def$fBD,clay.content=rep(20,n))

points(x=i.LLWR$LLWR.B, y=layers, type="l"); points(x=i.LLWR$LLWR.B, y=layers,pch=15)
points(x=f.LLWR$LLWR.B, y=layers, type="l", col=2); points(x=f.LLWR$LLWR.B, y=layers,pch=15, col=2)

# End (not run)

```

---

LLWR\_App

*A shiny for calculation of the usual least limiting water range*


---

## Description

A shiny for calculation of the usual least limiting water range

## Usage

```
LLWR_App()
```

**Arguments**

Available on the interactive web page

**Value**

shiny app

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com>

**See Also**

[stressTraffic](#)

---

llwr\_llmpr

*Least Limiting Water (LLWR) and Matric Potential Ranges (LLMPR)*


---

**Description**

A graphical solution and calculation of the least limiting water range and least limiting water matric potential range, including the corresponding the water content and water tensions limits.

**Usage**

```
llwr_llmpr(thetaR, thetaS, alpha, n, d, e, f = NULL, critical.PR, PD, Bd = NULL,
           h.FC, h.PWP, air.porosity,
           labels = c("AIR", "FC", "PWP", "PR"), ylab = "",
           graph1 = TRUE, graph2 = FALSE, ...)
```

**Arguments**

thetaR	the residual water content, $m^3m^{-3}$
thetaS	the water content at saturation, $m^3m^{-3}$
alpha	the scale parameter of the van Genuchten's model, $hPa^{-1}$
n	the shape parameter of the van Genuchten's model
d	a parameter of Busscher soil penetration resistance model. See details.
e	a parameter of Busscher soil penetration resistance model. See details.
f	a parameter of Busscher soil penetration resistance model. See details.
critical.PR	the limiting value of soil penetration resistance, MPa
PD	particle density, $Mgm^{-3}$
Bd	the bulk density to be displayed at bottom of the graph (optional), $Mgm^{-3}$
h.FC	the value of water tension at field capacity, hPa

h.PWP	the value of water tension at wilting point, hPa
air.porosity	the volumetric air-filled porosity
labels	the labels to h.FC, h.PWP, air.porosity and critical.PR
ylab	a title for the y-axis
graph1	logical; if TRUE (default) a graphical solution for the Least Limiting Water Range is displayed
graph2	logical; if TRUE (default) a graphical solution for the Least Limiting Matric Potential Range is displayed
...	Further graphical arguments

### Details

The penetration resistance model, as presented by Busscher (1990), is given by  $PR = d * \theta^e * BD^f$ . In this model, BD (bulk density) is calculated from thetaS (soil total porosity) and PD (particles density), i.e.,  $BD = PD * thetaS^{-1}$ . If the argument f is not passed, the model becomes  $PR = d * \theta^e$ .

### Value

A list of the LLWR and LLMPR, including the corresponding the water content and water tensions limits.

### Author(s)

Renato Paiva de Lima <renato\_agro\_@hotmail.com>

### References

- Leon, H. N., Almeida, B. G., Almeida, C. D. G. C., Freire, F. J., Souza, E. R., Oliveira, E. C. A., Silva, E. P. 2019. Medium-term influence of conventional tillage on the physical quality of a Typic Fragiudult with hardsetting behavior cultivated with sugarcane under rainfed conditions. *Catena*, 175: 37-46.
- Busscher, W. J. 1990. Adjustment of flat-tipped penetrometer resistance data to common water content. *Transactions of the ASAE*, 3: 519-524.
- van Genuchten, M. T. 1980. A closed-form equation for predicting the hydraulic conductivity of unsaturated soils 1. *Soil Science Society of America journal*, 44: 892-898.
- Silva et al. 1994. Characterization of the least limiting water range of soils. *Soil Science Society of America Journal*, 58: 1775-1781.
- Assouline, S., Or, D. 2014. The concept of field capacity revisited: Defining intrinsic static and dynamic criteria for soil internal drainage dynamics. *Water Resources Research*, 50: 4787-4802.
- Millington, R. J., Quirk, J. P. 1961. Permeability of porous solids. *Transactions of the Faraday Society*, 57: 1200-1207.
- Dexter, A. R., Czyz, E. A., Richard, G. 2012. Equilibrium, non-equilibrium and residual water: consequences for soil water retention. *Geoderma*, 177: 63-71.
- Moraes, M. T., Bengough, A. G., Debiasi, H., Franchini, J. C., Levien, R., Schnepf, A., Leitner, D., 2018. Mechanistic framework to link root growth models with weather and soil physical properties, including example applications to soybean growth in Brazil. *Plant and Soil*, 428: 67-92.

**Examples**

```

# Parameters from Leon et al. (2018), for usual physical restrictions threshold

llwr_llmpr(thetaR=0.1180, thetaS=0.36, alpha=0.133, n=1.30,
           d=0.005, e=-2.93, f=3.54, PD=2.65,
           critical.PR=4, h.FC=100, h.PWP=15000, air.porosity=0.1,
           labels=c("AFP", "FC", "PWP", "PR"),
           graph1=TRUE, graph2=FALSE, ylab=expression(psi~(hPa)), ylim=c(15000,1))
mtext(expression("Bulk density"~(Mg~m^-3)),1,line=2.2, cex=0.8)

llwr_llmpr(thetaR=0.1180, thetaS=0.36, alpha=0.133, n=1.30,
           d=0.005, e=-2.93, f=3.54, PD=2.65,
           critical.PR=4, h.FC=100, h.PWP=15000, air.porosity=0.1,
           graph1=FALSE, graph2=TRUE,
           labels=c("Air-filled porosity", "Field capacity",
                   "Permanent wilting point", "Penetration resistance"),
           ylim=c(0.1,0.30), ylab=expression(theta~(m^3~m^-3)))
mtext(expression("Bulk density"~(Mg~m^-3)),1,line=2.2, cex=0.8)

# Without bulk density effects in Busscher's model (i.e. f=NULL)

llwr_llmpr(thetaR=0.1180, thetaS=0.36, alpha=0.133, n=1.30,
           d=0.0165, e=-2.93, PD=2.65,
           critical.PR=3, h.FC=100, h.PWP=15000, air.porosity=0.1,
           graph1=TRUE, graph2=FALSE, ylim=c(15000,1),
           ylab=expression(psi~(hPa)))
mtext(expression("Bulk density"~(Mg~m^-3)),1,line=2.2, cex=0.8)

llwr_llmpr(thetaR=0.1180, thetaS=0.36, alpha=0.133, n=1.30,
           d=0.0165, e=-2.93, PD=2.65,
           critical.PR=3, h.FC=100, h.PWP=15000, air.porosity=0.1,
           graph1=FALSE, graph2=TRUE,
           ylim=c(0.1,0.30), ylab=expression(theta~(m^3~m^-3)))
mtext(expression("Bulk density"~(Mg~m^-3)),1,line=2.2, cex=0.8)

# Parameters from Leon et al. (2018), calculated physical restrictions threshold

thetaR <- 0.1180
thetaS <- 0.36
alpha <- 0.133
n <- 1.30
clay.content <- 15 # clay content 15 %
mim.gas.difusion <- 0.005
root.elongation.rate <- 0.3 # root elongation rate 30%

FC <- (1/alpha)*((n-1)/n)^((1-2*n)/n) # Assouline and Or (2014)
PWP <- 10^(3.514 + 0.0250*clay.content) # Dexter et al. (2012)
AIR.critical <- (mim.gas.difusion*(thetaS^2)^(1/(10/3))) # Millington and Quirk (1961)
PR.critical <- log(root.elongation.rate)/-0.4325 # Moraes et al. (2018)

```

```

llwr_llmpr(thetaR=thetaR, thetaS=thetaS, alpha=alpha, n=n,
           d=0.005, e=-2.93, f=3.54, PD=2.65, ylim=c(15000,1),
           critical.PR=PR.critical, h.FC=FC, h.PWP=PWP, air.porosity=AIR.critical,
           graph1=TRUE, graph2=FALSE, ylab=expression(psi~(hPa)))
mtext(expression("Bulk density"~(Mg~m^-3)),1,line=2.2, cex=0.8)

llwr_llmpr(thetaR=thetaR, thetaS=thetaS, alpha=alpha, n=n,
           d=0.005, e=-2.93, f=3.54, PD=2.65,
           critical.PR=PR.critical, h.FC=FC, h.PWP=PWP, air.porosity=AIR.critical,
           graph1=FALSE, graph2=TRUE,
           ylim=c(0.1,0.30), ylab=expression(theta~(m^3~m^-3)))
mtext(expression("Bulk density"~(Mg~m^-3)),1,line=2.2, cex=0.8)

# End (not run)

```

---

LLWR\_LLMPR\_App

*A shiny for calculation of least limiting water and matric potential ranges of agricultural soils with calculated physical restriction thresholds*

---

## Description

A shiny for calculation of least limiting water and matric potential ranges of agricultural soils with calculated physical restriction thresholds

## Usage

```
LLWR_LLMPR_App()
```

## Arguments

Available on the interactive web page

## Value

shiny app

## Author(s)

Renato Paiva de Lima <renato\_agro\_@hotmail.com>

## See Also

[stressTraffic](#)

---

maxcurv

*Maximum Curvature Point*


---

## Description

Function to determine the maximum curvature point of an univariate nonlinear function of  $x$ .

## Usage

```
maxcurv(x.range, fun,
method = c("general", "pd", "LRP", "spline"),
x0ini = NULL,
graph = TRUE, ...)
```

## Arguments

<code>x.range</code>	a numeric vector of length two, the range of $x$ .
<code>fun</code>	a function of $x$ ; it must be a one-line-written function, with no curly braces <code>{}</code> .
<code>method</code>	a character indicating one of the following: "general" - for evaluating the general curvature function ( $k$ ), "pd" - for evaluating perpendicular distances from a secant line, "LRP" - a NLS estimate of the maximum curvature point as the breaking point of Linear Response Plateau model, "spline" - a NLS estimate of the maximum curvature point as the breaking point of a piecewise linear spline. See details.
<code>x0ini</code>	an initial $x$ -value for the maximum curvature point. Required only when "LRP" or "spline" are used.
<code>graph</code>	logical; if TRUE (default) a curve of <code>fun</code> is plotted.
<code>...</code>	further graphical arguments.

## Details

The method "LRP" can be understood as an especial case of "spline". And both models are fitted via [nls](#). The method "pd" is an adaptation of the method proposed by Lorentz et al. (2012). The "general" method should be preferred for finding *global* points. On the other hand, "pd", "LRP" and "spline" are suitable for finding *local* points of maximum curvature.

## Value

A list of

<code>fun</code>	the function of $x$ .
<code>x0</code>	the $x$ critical value.
<code>y0</code>	the $y$ critical value.
<code>method</code>	the method of determination (input).

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Lorentz, L.H.; Erichsen, R.; Lucio, A.D. (2012). Proposal method for plot size estimation in crops. *Revista Ceres*, 59:772–780.

**See Also**

[function](#), [curve](#)

**Examples**

```
# Example 1: an exponential model
f <- function(x) exp(-x)
maxcurv(x.range = c(-2, 5), fun = f)

# Example 2: Gompertz Growth Model
Asym <- 8.5
b2 <- 2.3
b3 <- 0.6
g <- function(x) Asym * exp(-b2 * b3 ^ x)
maxcurv(x.range = c(-5, 20), fun = g)

# using "pd" method
maxcurv(x.range = c(-5, 20), fun = g, method = "pd")

# using "LRP" method
maxcurv(x.range = c(-5, 20), fun = g, method = "LRP", x0ini = 6.5)

# Example 3: Lessman & Atkins (1963) model for optimum plot size
a = 40.1
b = 0.72
cv <- function(x) a * x^-b
maxcurv(x.range = c(1, 50), fun = cv)

# using "spline" method
maxcurv(x.range = c(1, 50), fun = cv, method = "spline", x0ini = 6)

# End (not run)
```



**Description**

It calculates the sedimentation time of soil particle in aqueous media using Stokes equation, i.e., the time needed for the particles of soil larger than the size attributed as input to sediment in aqueous media, usually water.

**Usage**

```
particle.sedimentation(d, h=0.2, g=9.81, v=0.001, Pd=2650, Wd=1000)
```

**Arguments**

d	the lower limit of soil particle diameter (micrometers) to sediment withing the calculated time.
h	the vertical distance (meters) from which the particles fall. Default is 0.2 m.
g	the acceleration of gravity, in m/s <sup>2</sup> . Default is 9.81 m/s <sup>2</sup> .
v	the viscosity of the fluid, in N/s/m <sup>2</sup> . Default is 0.001 N/s/m <sup>2</sup> , for water at 20 degrees Celsius.
Pd	the particle density, in kg/m <sup>3</sup> . Default is 2650 kg/m <sup>3</sup> .
Wd	the density of the fluid, in kg/m <sup>3</sup> . Default is 1000 kg/m <sup>3</sup> .

**Value**

A data.frame containing the estimated time for the sedimentation of particles.

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com>

**References**

Hillel, D. (2003). *Introduction to environmental soil physics*. Elsevier. p.39-51. Doi:10.1016/B978-012348655-4/50004-6

**Examples**

```
# Example 1
particle.sedimentation(d=2, h=0.2, g=9.81, v=1.002*10^-3, Pd=2650, Wd=1000)

# Example 2
d <- c(2000, 200, 50, 10, 2, 1)
time <- particle.sedimentation(d=d, h=0.2, g=9.81, v=1.002*10^-3, Pd=2650, Wd=1000)

plot(x=d, y=time$hours, log = "x", xaxt = "n",
      ylab = "time of sedimentation (hours)", xlab = "particle diameter (micrometer)")
axis(1,at=d, labels=d)

# End (not run)
```

---

particle.sedimentation\_App

*A shiny for time of particle sedimentation*

---

**Description**

A shiny for time of particle sedimentation

**Usage**

```
particle.sedimentation_App()
```

**Arguments**

Available on the interactive web page

**Value**

shiny app

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com>

**See Also**

[stressTraffic](#)

---

plotCIsigmaP

*Percentile Confidence Intervals for Simulated Preconsolidation Stress*

---

**Description**

Build and plot percentile confidence intervals for preconsolidation stress simulated from [simSigmaP](#).

**Usage**

```
plotCIsigmaP(msim, conf.level = 0.95, shade.col = "orange",  
ordered = TRUE, xlim = NULL, xlab = expression(sigma[P]),  
las = 1, mar = c(4.5, 6.5, 2, 1), ...)
```

**Arguments**

<code>msim</code>	an object of class "simSigmaP".
<code>conf.level</code>	the confidence level for the intervals.
<code>shade.col</code>	a character or number indicating the color of the shaded area delimiting each CI. See <a href="#">colors</a> .
<code>ordered</code>	logical; should the intervals be displayed according to the value of the simulated mean?
<code>xlim</code>	optional; a numeric vector of length two containing the limits of the <i>x</i> -axis.
<code>xlab</code>	optional; a character indicating the <i>x</i> -axis label.
<code>las</code>	optional; see <a href="#">par</a> .
<code>mar</code>	optional; see <a href="#">par</a> .
<code>...</code>	further graphical parameters; see <a href="#">par</a> .

**Value**

A numeric matrix containing the simulated mean, coefficient of variation, lower and upper CI limits and the name of the method used to calculate the preconsolidation stress.

**Author(s)**

Anderson Rodrigo da Silva <[anderson.agro@hotmail.com](mailto:anderson.agro@hotmail.com)>

**See Also**

[simSigmaP](#), [sigmaP](#)

**Examples**

```
# input data: stress and void ratio
pres <- c(1, 12.5, 25, 50, 100, 200, 400, 800, 1600)
VR <- c(1.43, 1.41, 1.40, 1.39, 1.35, 1.31, 1.25, 1.18, 1.12)

# simulation (may take a few seconds)
simres <- simSigmaP(VR, pres, nsim = 30)
head(simres)

# percentile confidence intervals
ci <- plotCIsigmaP(simres, conf.level = 0.95,
  shade.col = "blue", ordered = TRUE)
print(ci)

# End (Not run)
```

---

PredComp

*A shiny for simulation of soil compaction*

---

**Description**

A shiny for for simulation of soil compaction

**Usage**

PredComp()

**Arguments**

Available on the interactive web page

**Value**

shiny app

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com>

**See Also**

[stressTraffic](#)

---

psd

*Soil Pore Size Distribution*

---

**Description**

The unimodal soil pore size distribution based on van Genuchten's model.

**Usage**

psd(thetaS, thetaR, alpha, n, h)

**Arguments**

thetaS	the water content at saturation.
thetaR	the residual water content.
alpha	the scale parameter of the van Genuchten's model (hPa <sup>-1</sup> ).
n	the shape parameter in van Genuchten's formula.
h	a vector of water tension (hPa) on the range of water retention curve.

**Value**

A numeric vector containing the soil pore size distribution as a function of soil water tension.

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com>

**References**

Ghiberto, P. J., Imhoff, S., Libardi, P. L., Silva, A. P. D., Tormena, C. A., Pilatti, M. A. (2015). Soil physical quality of Mollisols quantified by a global index. *Scientia Agricola*, 72(2):167-174.

**Examples**

```
# EXAMPLE 1
x <- seq(log10(1),log10(15000),len=100)
h <- 10^x
y <- psd(thetaR = 0.15,thetaS = 0.55, alpha = 0.048, n = 1.5, h=h)
plot(x=h,y=y, log="x", xlab="h (hPa)", ylab=expression(delta*theta/delta*h), ylim=c(0,0.005))

# EXAMPLE 2
x <- seq(log10(1),log10(15000),len=100)
h <- 10^x
y <- psd(thetaR = 0.20,thetaS = 0.61, alpha = 0.1232, n = 1.3380,h=h)
plot(x=h,y=y, log="x", xlab="h (hPa)", ylab=expression(delta*theta/delta*h), ylim=c(0,0.01))

# EXAMPLE 3
x <- seq(log10(1),log10(15000),len=100)
h <- 10^x
y <- psd(thetaR = 0.154,thetaS = 0.600, alpha = 0.103, n = 2.365,h=h)
plot(x=h,y=y, log="x", xlab="h (hPa)", ylab=expression(delta*theta/delta*h), ylim=c(0,0.03))
ax <- c(1,10,100,1000,10000)
radius <- r(h=ax)
axis(3,at=ax, labels=round(radius,2))
mtext("Equivalent pore radius"~(mu*m),3,line=2.5, cex=0.9)

# End (not run)
```

r

*Equation of capillary***Description**

The equivalent pore radius as a function of soil water tension.

**Usage**

```
r(h, surface.tension.water=0.072, water.density=1000, water.pore.contact.angle=0)
```

**Arguments**

h                    The water tension (hPa).  
 surface.tension.water                    Surface tension of water (N/m).  
 water.density      Density of water (kg/m<sup>3</sup>).  
 water.pore.contact.angle                Water pore contact angle (degrees).

**Value**

The equivalent pore radius, in micrometer..

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com>

**References**

Ghiberto, P. J., Imhoff, S., Libardi, P. L., Silva, A. P. D., Tormena, C. A., Pilatti, M. A. (2015). Soil physical quality of Mollisols quantified by a global index. *Scientia Agricola*, 72(2):167-174.

**Examples**

```
x <- seq(log10(1), log10(15000), len=50)
h <- 10^x
y <- r(h=h)
plot(x=h, y=y, log="yx", xlab="h (hPa)", yaxt='n', ylab="", ylim=c(0.1, 1500))
ax <- c(0.1, 1, 10, 100, 1000, 1500)
axis(2,at=ax, labels=ax)
mtext("Pore radius"~ (mu*m), 2, line=2.5)

# End (not run)
```

---

 Rsq

*Multiple R-squared*


---

**Description**

Function to calculate the *multiple R-squared* and the *adjusted R-squared* from a fitted model via [lm](#) or [aov](#), i.e., linear models. For a model fitted via [nls](#), nonlinear models, the *pseudo R-squared* is returned.

**Usage**

```
Rsq(model)
```

**Arguments**

model                    a model fitted via [lm](#), [aov](#) or [nls](#).

**Value**

A list of

R.squared                the multiple R-squared (for linear models) or the Pseudo R-squared (for nonlinear models).

adj.R.squared          the adjusted R-squared.

**Author(s)**

Anderson Rodrigo da Silva <[anderson.agro@hotmail.com](mailto:anderson.agro@hotmail.com)>

**See Also**

[lm](#), [summary.lm](#), [aov](#), [nls](#)

**Examples**

```
# example 1 [linear model]
y <- rnorm(10)
x <- 1:10
fit <- lm(y ~ x)
summary(fit)
Rsq(fit)

# example 2 [nonlinear model for Load Bearing Capacity]
data(compaction)
attach(compaction)
out <- fitlbc(theta = Mois, sigmaP = PS)
summary(out)
Rsq(out)

# End (not run)
```

---

r\_App

*A shiny for equation of capillarity*

---

**Description**

A shiny for equation of capillarity

**Usage**

r\_App()

**Arguments**

Available on the interactive web page

**Value**

shiny app

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com>

**See Also**

[stressTraffic](#)

---

sigmaP

*Preconsolidation Stress*

---

**Description**

A function to determine the preconsolidation stress ( $\sigma_P$ ). It is a parameter obtained from the soil compression curve and has been used as an indicator of soil load-bearing capacity as well as to characterize the impacts suffered by the use of machines. The function `sigmaP()` contains implementations of the main methods for determining the pre-consolidation stress, such as the Casagrande method, the method of Pacheco Silva, the regression methods and the method of the virgin compression line intercept.

**Usage**

```
sigmaP(voidratio, stress, n4VCL = 3,
       method = c("casagrande", "VCLzero", "reg1", "reg2", "reg3", "reg4", "pacheco"),
       mcp = NULL, graph = TRUE, ...)
```

**Arguments**


<code>voidratio</code>	a numeric vector containing void ratio (or bulk density) values.
<code>stress</code>	a numeric vector containing the applied stress sequence.
<code>n4VCL</code>	the number of points for calculating the slope of the soil Virgin Compression Line (VCL), which is obtained by linear regression.
<code>method</code>	a character indicating which method is to be computed; one of the following: casagrande (default), VCLzero, reg1, reg2, reg3, reg4 or pacheco; see Details.
<code>mcp</code>	the maximum curvature point in log10 scale of stress; required only if the method casagrande is used.
<code>graph</code>	logical; if TRUE (default) the compression curve is plotted.
<code>...</code>	further graphical arguments.



**Details**

casagrande is the method proposed by Casagrande (1936). The preconsolidation stress obtained via VCLzero corresponds to the intersection of the soil *Virgin Compression Line* (VCL) with the x-axis at zero applied stress, as described by Arvidsson & Keller (2004). reg1, reg2, reg3 and reg4 are regression methods that obtain the preconsolidation stress value as the intercept of the VCL and a regression line fitted with the first two, three, four and five points of the curve, respectively, as described by Dias Junior & Pierce (1995). pacheco is the method of Pacheco Silva (ABNT, 1990).

You may follow the flowchart below to understand the determination of the preconsolidation stress through  $\sigma_p()$ .



*ws* - the weight of wet soil; *ds* - the weight of dry soil; *dc* - the diameter of the cylinder; *hc* - the height of the cylinder; *pd* - the particle density; *df* - a numeric vector containing soil deformation values;  $\sigma_p$  - preconsolidation pressure; *VCL* - Virgin Compression Line.

**Value**

A list of

sigmaP	the preconsolidation stress.
method	the method used as argument.
mcp	the maximum curvature point in log10 scale of stress; stored only if the method casagrande is used.
CI	the compression index.
SI	the swelling index.

### Author(s)

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

### References

- ABNT - Associacao Brasileira de Normas Tecnicas. (1990). *Ensaio de adensamento unidimensional*: NBR 12007. Rio de Janeiro. 13p.
- Arvidsson, J.; Keller, T. (2004). Soil precompression stress I. A survey of Swedish arable soils. *Soil & Tillage Research*, 77:85-95.
- Bowles, J. A. (1986). *Engineering Properties of Soils and their Measurements*, 3rd edition. McGraw-Hill Book Company, Inc. NY, 218pp.
- Casagrande, A. (1936). *The determination of the pre-consolidation load and its practical significance*. In: Proceedings of the International Conference on Soil Mech. and Found. Eng. (ICSMFE), Cambridge, MA, 22-26 June 1936, vol. 3. Harvard University, Cambridge, MA, USA, pp. 60-64.
- Dias Junior, M. S.; Pierce, F. J. (1995). A simple procedure for estimating preconsolidation pressure from soil compression curves. *Soil Technology*, 8:139-151.

### See Also

[voidratio](#), [maxcurv](#), [fitlbc](#)

### Examples

```
pres <- c(1, 12.5, 25, 50, 100, 200, 400, 800, 1600)
VR <- c(0.846, 0.829, 0.820, 0.802, 0.767, 0.717, 0.660, 0.595, 0.532)

plot(VR ~ log10(pres), type = "b") # find the 'mcp'
sigmaP(VR, pres, method = "casagrande", mcp = 1.6, n4VCL = 2)

# fitting the VCL
sigmaP(VR, pres, method = "casagrande", mcp = 1.6, n4VCL = 3)

# self-calculation of "mcp" argument for Casagrande method
sigmaP(VR, pres, method = "casagrande", n4VCL = 3)

# Pacheco method
sigmaP(VR, pres, method = "pacheco")

# Regression method
sigmaP(VR, pres, method = "reg3")
```

```
# End (not run)
```

---

```
simSigmaP
```

```
Simulating Preconsolidation Stress
```

---

## Description

Simulating preconsolidation pressure, compression and swelling indices, based on a multivariate Gaussian distribution for the parameters of the compression curve.

## Usage

```
simSigmaP(voidratio, stress,
  what.out = c("sigmaP", "CI", "SI"),
  method = c("casagrande", "VCLzero", "reg1", "reg2", "reg3", "reg4", "pacheco"),
  n4VCL = 3, nsim = 100)
```

## Arguments

<code>voidratio</code>	a numeric vector containing void ratio (or bulk density) values.
<code>stress</code>	a numeric vector containing the applied stress sequence.
<code>what.out</code>	a character indicating which <code>sigmaP()</code> output should be simulated. It must be on of "sigmaP" (default), "CI" or "SI".
<code>method</code>	a character vector indicating which methods should be used.
<code>n4VCL</code>	the number of points for calculating the slope of the soil Virgin Compression Line (VCL), which is obtained by linear regression. Default is 3.
<code>nsim</code>	the number of simulations. Default is 100. Warning: it may cause time demanding.

## Value

A numeric matrix containing the simulated values for each method selected as input.

## Author(s)

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

## See Also

[sigmaP](#), [plotCIsigmaP](#)

**Examples**

```
# input data: stress and void ratio
pres <- c(1, 12.5, 25, 50, 100, 200, 400, 800, 1600)
VR <- c(1.43, 1.41, 1.40, 1.39, 1.35, 1.31, 1.25, 1.18, 1.12)

# simulation (may take a few seconds)
simres <- simSigmaP(VR, pres, nsim = 30)
head(simres)

# plot percentile confidence intervals
ci <- plotCIsigmaP(simres, conf.level = 0.95,
  shade.col = "blue", ordered = TRUE)
print(ci)

# End (Not run)
```

Sindex

*The S Index***Description**

Function to calculate the S index (Dexter, 2004) for evaluating the soil physical quality based on the *Water Retention Curve* (van Genuchten, 1980).

$$S = -n * (\theta_S - \theta_R) * (1 + 1/m)^{-(1+m)}$$

**Usage**

```
Sindex(theta_R, theta_S, alpha, n, m = 1 - 1/n, vcov = NULL,
  nsim = 999, conf.level = 0.95, graph = TRUE, ...)
```

**Arguments**

theta_R	the residual water content.
theta_S	the water content at saturation.
alpha	a scale parameter of the van Genuchten's formula.
n	a shape parameter in van Genuchten's formula.
m	a shape parameter in van Genuchten's Formula. Default is $1 - 1/n$ (Mualem, 1976).
vcov	optional (default is NULL); a variance-covariance matrix of the estimates which is used to perform Monte Carlo simulations of the parameters theta_R, theta_S, alpha and n for building a simulated confidence interval of the S index (in modulus).
nsim	the number of Monte Carlo simulations; default is 999. It is used only if vcov is specified.
conf.level	the confidence level; default is 0.95. It is used only if vcov is specified.
graph	logical; if TRUE (default), the soil water retention curve is plotted.
...	further graphical arguments.

**Value**

A list of

h_i	the modulus of the water potential at the inflection point.
theta_i	the water content at the inflection point.
S.index	the modulus of the S index.
PhysicalQuality	A character indicating the soil physical quality, as proposed by Dexter (2004).
simCI	the simulated confidence interval. It is stored only if vcov is specified.
conf.level	the confidence level for the simulated confidence interval. It is stored only if vcov is specified.

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

- Dexter, A. R. (2004). Soil physical quality Part I. Theory, effects of soil texture, density, and organic matter, and effects on root growth. *Geoderma*, 120:201-214.
- Genuchten, M. T. van. (1980). A closed form equation for predicting the hydraulic conductivity of unsaturated soils. *Soil Science Society of America Journal*, 44:892-898.
- Mualem, Y. (1976). A new model for predicting the hydraulic conductivity of unsaturated porous media, *Water Resource Research*, 12:513-522.

**See Also**

[soilwater](#), [fitsoilwater](#)

**Examples**

```
# Dexter (2004, Table 1)
Sindex(0, 0.395, 0.0217, 1.103, xlim = c(0, 1000))
Sindex(0, 0.335, 0.0616, 1.139, xlim = c(0, 1000))
# ...
Sindex(0, 0.226, 0.0671, 1.581, xlim = c(0, 1000))

# End (not run)
```

---

skp1994

*LLWR Data Set*

---

**Description**

Data set presented by Leao et al. (2005), for determining the Least Limiting Water Range.

**Usage**

```
data(skp1994)
```

**Format**

A data frame with 64 observations on the following 4 variables:

BD a numeric vector containing soil bulk density values, in Mg/m<sup>3</sup>.

W a numeric vector containing volumetric water content values, in m<sup>3</sup>/m<sup>3</sup>.

PR a numeric vector containing penetration resistance values, in MPa.

h a numeric vector containing matric head values, in cm.

**Source**

Leao et al. (2005). An Algorithm for Calculating the Least Limiting Water Range of Soils. *Agronomy Journal*, 97:1210-1215.

**Examples**

```
data(skp1994)
summary(skp1994)
```

---

SoilAggregate

*Soil Aggregate Size Data Set*

---

**Description**

Data set for determining soil aggregate size distribution.

**Usage**

```
data(SoilAggregate)
```

**Format**

A data frame with 12 observations on 7 variables.

ID a factor with the names of the soil samples.

D3

D1.5

D0.75

D0.375

D0.178

D0.053

**Examples**

```
data(SoilAggregate)
summary(SoilAggregate)
```

---

soilDeformation	<i>Soil deformation by O'Sullivan and Robertson (1996)</i>
-----------------	--

---

**Description**

It calculates bulk density variation as a function of the applied mean normal stress using critical state theory, by O'Sullivan and Robertson (1996).

**Usage**


```
soilDeformation(stress, p.density, iBD, N, CI, k, k2, m, graph = FALSE, ...)
```

**Arguments**

stress	a numeric vector containing the values of mean normal stress, kPa; Note that stress can also be a vector of length 1.
p.density	a numeric vector containing the values of particle density to each stress, $Mgm^{-3}$ .
iBD	a numeric vector containing the values of initial bulk density to each stress, $Mgm^{-3}$ .
N	the specific volume at $p = 1$ kPa, to each stress
CI	the compression index, to each stress; check details
k	the recompression index, to each stress; check details
k2	the slope of the steeper recompression line to each stress (similar to the $k'$ in O'Sullivan and Robertson (1996) model); check details
m	the value that separates yield line and VCL to each stress; check details
graph	logical; shall soilDeformation plot the graph model (only the first parameters set is plotted)?
...	further graphical arguments. See <a href="#">par</a> .

### Details

The specific volume ( $v$ ) is given as  $v = PD/BD$ , where PD is particle density and BD is the bulk density. Please, check each parameter from O'Sullivan and Robertson (1996) model in the figure below.



### Value

A list of

iBD	initial bulk density, $Mgm^{-3}$
fBD	final bulk density, $Mgm^{-3}$
vi	initial specific volume
vf	final specific volume
I%	variation of soil bulk density (%) after the applied stress

### Author(s)

Renato Paiva de Lima <renato\_agro\_@hotmail.com>

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

Alvaro Pires da Silva <apisilva@usp.br>



## References

O'Sullivan, M.F.; Robertson, E.A.G. 1996. Critical state parameters from intact samples of two agricultural soils. *Soil and Tillage Research*, 39:161-173.

Keller, T.; Defosse, P.; Weisskopf, P.; Arvidsson, J.; Richard, G. 2007. SoilFlex: a model for prediction of soil stresses and soil compaction due to agricultural field traffic including a synthesis of analytical approaches. *Soil and Tillage Research*, 93:391-411.

## Examples

```
# EXAMPLE 1
soilDeformation(stress = 300,
                p.density = 2.67,
                iBD = 1.55,
                N = 1.9392,
                CI = 0.06037,
                k = 0.00608,
                k2 = 0.01916,
                m = 1.3, graph=TRUE, ylim=c(1.4, 1.8))

# EXAMPLE 2 (combining it with soil stress)
stress <- stressTraffic(inflation.pressure=200,
                      recommended.pressure=200,
                      tyre.diameter=1.8,
                      tyre.width=0.4,
                      wheel.load=4000,
                      conc.factor=c(4,5,5,5,5,5),
                      layers=c(0.05,0.1,0.3,0.5,0.7,1),
                      plot.contact.area = FALSE)

stress.mean <- stress$Stress$sigma_mean
layers <- stress$Stress$Layers
n <- length(layers)

def <- soilDeformation(stress = stress.mean,
                      p.density = rep(2.67, n),
                      iBD = rep(1.55, n),
                      N = rep(1.9392, n),
                      CI = rep(0.06037, n),
                      k = rep(0.00608, n),
                      k2 = rep(0.01916, n),
                      m = rep(1.3, n), graph=TRUE, ylim=c(1.4, 1.8))

# Graph
plot(x = 1, y = 1,
     xlim=c(1.4, 1.7), ylim=c(1, 0), xaxt = "n",
     ylab = "Soil Depth", xlab = "", type="l", main="")
axis(3)
mtext("Bulk Density", side=3, line=2.5)

initial.BD <- def$iBD
final.BD <- def$fBD
```

```

points(x=initial.BD, y=layers, type="l")
points(x=initial.BD, y=layers,pch=15)
points(x=final.BD, y=layers, type="l", col=2)
points(x=final.BD, y=layers,pch=15, col=2)

# End (not run)

```

---

soilStrength

*Estimation of precompression stress by Severiano et al. (2013)*


---

### Description

It calculates the precompression stress using the pedo-transfer function from Severiano et al. (2013)

### Usage

```
soilStrength(clay.content, matric.suction = NULL, water.content = NULL)
```

### Arguments

`clay.content` a numeric vector containing the values of clay for each soil layer, %. Note that it can also be a unique value.

`matric.suction` a numeric vector containing the values of matric suction for each clay content, kPa.

`water.content` a numeric vector containing the values of water content for each clay content,  $m^3 m^{-3}$ . Note that `water.content` must be passed if `matric.suction` is not. See details.

### Details

Intervals of soil water content/matric suction to be used as input for estimating soil strength according to Severiano et al. (2013).

Clay content (%)	Matric suction (kPa)	Water content ( $m^3 m^{-3}$ )
< 20	1-10,000	0.41-0.05
21-31	1-10,000	0.44-0.09
32-37	1-10,000	0.45-0.11
38-52	1-10,000	0.49-0.13
< 52	1-10,000	0.50-0.15

### Value

A two-columns data frame:

`Pc` the precompression stress (Severiano et al. 2013)

`LL.Pc` the lower limit of precompression stress in according to the Terranimo model criteria (see Stettler et al. 2014). Note:  $LL.Pc = Pc * 0.5$

`UL.Pc` the upper limit of precompression stress in according to the Terranimo model criteria (see Stettler et al. 2014). Note:  $UL.Pc = Pc * 1.1$

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com>  
 Anderson Rodrigo da Silva <anderson.agro@hotmail.com>  
 Alvaro Pires da Silva <apisilva@usp.br>

**References**

Severiano, E.C.; Oliveira, G.C.; Dias Junior, M.S.; Curi, N.C.; Costa, K. A.P.; Carducci, C.E. 2013. Preconsolidation pressure, soil water retention characteristics, and texture of Latosols in the Brazilian Cerrado. *Soil Research*, 51:193-202.

Stettler, M., Keller, T., Weisskopf, P., Lamande, M., Lassen, P., Schjonning, P., 2014. Terranimo - a web-based tool for evaluating soil compaction. *Landtechnik*, 69:132-137.

**See Also**

[stressTraffic](#)

**Examples**

```
# EXEMPLE 1 (using water content)
soilStrength(clay.content=c(25,28,30,30,30),
             water.content = c(0.26,0.27,0.29,0.32,0.32))

# EXEMPLE 2 (using matric suction)
soilStrength(clay.content=c(25,28,30,30,30),
             matric.suction = c(100,330,1000,3000,5000))

# EXAMPLE 3 (combining it with soil stress)
stress <- stressTraffic(inflation.pressure=200,
                      recommended.pressure=200,
                      tyre.diameter=1.8,
                      tyre.width=0.4,
                      wheel.load=4000,
                      conc.factor=c(4,5,5,5,5),
                      layers=c(0.05,0.1,0.3,0.5,0.7,1),
                      plot.contact.area = FALSE)

strength <- soilStrength(clay.content=c(25,28,30,30,30,30),
                       matric.suction = c(30,100,100,100,200,200))

# Graph
plot(x = 1, y = 1,
     xlim=c(0,300),ylim=c(1,0),xaxt = "n",
     ylab = "Soil Depth",xlab = "", type="l", main="")
axis(3)
mtext("Vertical Stress",side=3,line=2.5)
```

```

stressz <- stress$Stress$sigma_vertical
layers <- stress$Stress$Layers
points(x=stressz, y=layers, type="l")

# Green zone
x0 <- strength$LL.Pc
x1 <- rep(0,length(layers))
y0 <- layers
y1 <- rev(layers)
polygon(x=c(x0,x1), y = c(y0,y1),density = NA,
        col=rgb(red=0, green=1, blue=0, alpha=0.3))

# Yellow zone
x0 <- strength$UL.Pc
x1 <- rev(strength$LL.Pc)
y0 <- layers
y1 <- rev(layers)
polygon(x=c(x0,x1), y = c(y0,y1),density = NA,
        col=rgb(red=1, green=1, blue=0, alpha=0.3))

# Red zone
x0 <- rep(300,length(layers))
x1 <- rev(strength$UL.Pc)
y0 <- layers
y1 <- rev(layers)
polygon(x=c(x0,x1), y = c(y0,y1),density = NA,
        col=rgb(red=1, green=0, blue=0, alpha=0.3))

# End (not run)

```

---

soilStrength2	<i>Estimation of precompression stress by Schjonning and Lamande (2018)</i>
---------------	---

---

### Description

It calculates the precompression stress using the pedo-transfer function from Schjonning and Lamande (2018)

### Usage

```
soilStrength2(bulk.density, matric.suction, clay.content)
```

### Arguments

clay.content    a numeric vector containing the values of clay content, %  
matric.suction    a numeric vector containing the values of matric suction, hPa  
bulk.density    a numeric vector containing the values of soil bulk density,  $Mgm^{-3}$

**Details**

The function returns 0 for soil properties values beyond the range for which the function was built.

**Value**

PC                    the precompression stress

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com> Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Schjonning, P.; Lamande, M., 2018. Models for prediction of soil precompression stress from readily available soil properties. *Geoderma*, 320:115-125.

**See Also**

[stressTraffic](#)

**Examples**

```
# EXAMPLE 1

soilStrength2(bulk.density=1.46, matric.suction=100, clay.content=0.3)

# EXAMPLE 2

matric.suction <- seq(from=60,to=1000,len=50) # range of matric suction from 60 to 1200 hPa
out <- soilStrength2(bulk.density=1.46, matric.suction=matric.suction, clay.content=0.3)
plot(x=matric.suction,y=out,
     ylab="Precompression stress (kPa)", xlab="Matric suction (hPa)")

# End (not run)
```

---

soilStrength3

*Estimation of precompression stress by Saffih-Hdadi et al. (2009)*

---

**Description**

It calculates the precompression stress using the pedo-transfer function from Saffih-Hdadi et al. (2009)

**Usage**

```
soilStrength3(bulk.density, water.content,
             texture=c("VeryFine","Fine","MediumFine","Medium","Coarse"))
```

**Arguments**

bulk.density a numeric vector containing the values of soil bulk density,  $Mgm^{-3}$   
water.content a numeric vector containing the values of gravimetric water content, %  
texture the soil texture group. See examples

**Details**

The function returns 0 for soil properties values beyond the range for which the function was built.

**Value**

PC the precompression stress

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com> Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Saffih-Hdadi, K.; Defosse, P.; Richard, G.; Cui, Y. J.; Tang, A. M.; Chaplain, V, 2009. A method for predicting soil susceptibility to the compaction of surface layers as a function of water content and bulk density *Geoderma*, 115: 96-103.

**See Also**

[stressTraffic](#)

**Examples**

```
# EXAMPLE 1

soilStrength3(bulk.density=1.1, water.content=40, texture="VeryFine")
soilStrength3(bulk.density=1.2, water.content=20, texture="Fine")
soilStrength3(bulk.density=1.3, water.content=15, texture="MediumFine")
soilStrength3(bulk.density=1.4, water.content=15, texture="Medium")
soilStrength3(bulk.density=1.5, water.content=10, texture="Coarse")

# End (not run)
```

soilStrength4

*Estimation of precompression stress by Lebert and Horn (1991)***Description**

It calculates the soil strength through precompression stress using the pedo-transfer function from Lebert and Horn (1991)

**Usage**

```
soilStrength4(BD=1.55,AC=10,AWC=15,PWP=26,Ks=0.29,
              OM=1.5,C=30,phi=36,texture="Clay>35", pF=1.8)
```

**Arguments**

BD	a numeric vector containing the values of soil bulk density, $Mgm^{-3}$
AC	a numeric vector containing the values of volumetric air capacity at the specified pF, %
AWC	a numeric vector containing the values of volumetric available water at the specified pF, %
PWP	a numeric vector containing the values of volumetric non available water capacity (pF > 4.2), %
Ks	a numeric vector containing the values of saturated hydraulic conductivity, $10^3cms^{-1}$
OM	a numeric vector containing the values of organic matter, %
C	a numeric vector containing the values of cohesion at the specified pF, kPa
phi	a numeric vector containing the values of angle of internal friction at the specified pF, degree
texture	the soil texture classification. See details
pF	the '1.8' or '2.5' value pF

**Details**

The function returns '0' for soil properties values beyond the range for which the function was built. The default for this function is the values given in the application example by Horn and Fleige (2003). In the 'texture' argument, the user must pass the textural classification 'Sand', 'SandLoam', 'Silt', 'Clay<35' or 'Clay>35'. See examples.

**Value**

PC the precompression stress

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com> Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Lebert, M., Horn, R. (1991) A method to predict the mechanical strength of agricultural soils. *Soil and Tillage Research*, 19: 275-256.

Horn, R., Fleige, H. (2003) A method for assessing the impact of load on mechanical stability and on physical properties of soils. *Soil and Tillage Research*, 73: 89-99.

**See Also**

[stressTraffic](#)

**Examples**

```
soilStrength4(BD=1.55,AC=10,AWC=15,PWP=26,Ks=0.29,OM=1.5,
             C=30,phi=36,texture="Clay>35", pF=1.8) # Exemple from Horn and Fleige (2003), Table 7

# End (not run)
```

---

soilStrength5	<i>Estimation of precompression stress by Imhoff et al. (2004)</i>
---------------	--

---

**Description**

It calculates the soil strength using precompression stress using the pedo-transfer function from Imhoff et al. (2004)

**Usage**

```
soilStrength5(bulk.density, water.content, clay.content)
```

**Arguments**

bulk.density	a numeric vector containing the values of soil bulk density, $Mgm^{-3}$
water.content	a numeric vector containing the values of water content, (g/g)
clay.content	a numeric vector containing the values of clay content, %

**Details**

The function returns 0 for soil properties values beyond the range for which the function was built.

**Value**

PC	the precompression stress
----	---------------------------

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com> Anderson Rodrigo da Silva <anderson.agro@hotmail.com>



## References

Imhoff, S., Da Silva, A. P., Fallow, D. (2004) Susceptibility to Compaction, Load Support Capacity, and Soil Compressibility of Hapludox. *Soil Science Society of America Journal*, 68: 17-24.

## See Also

[stressTraffic](#)

## Examples

```
# EXAMPLE 1

soilStrength5(clay.content=60, water.content=0.30, bulk.density=1.25)
soilStrength5(clay.content=35, water.content=0.23, bulk.density=1.40)
soilStrength5(clay.content=20, water.content=0.10, bulk.density=1.60)

# EXAMPLE 2
water.content <- seq(0.1,0.30,len=20) # range of water content from 0.1 to 0.30 (g g-1)
out <- soilStrength5(clay.content=20, water.content=water.content , bulk.density=1.60)
plot(x=water.content,y=out,
     ylab="Precompression stress (kPa)", xlab="Water content")

# End (not run)
```

---

soilwater

*Soil Water Retention, based on the van Genuchten's (1980) formula*

---

## Description

Function to calculate the soil water content based on the van Genuchten's (1980) formula:

$$\theta = \theta_R + (\theta_S - \theta_R)(1 + (\alpha x)^n)^{-m}$$

## Usage

```
soilwater(x, theta_R, theta_S, alpha, n, m = 1 - 1/n,
          saturation.index = FALSE)
```

## Arguments

x	the matric potential.
theta_R	the residual water content.
theta_S	the water content at saturation.
alpha	a scale parameter of the van Genuchten's formula.
n	a shape parameter in van Genuchten's formula.

`m` a shape parameter in van Genuchten's Formula. Default is  $1 - 1/n$  (Mualem, 1976).

`saturation.index` logical; if FALSE (default) the outcome is the soil water content, otherwise the saturation index is returned.

### Value

The the soil water content or the saturation index (a value between 0 and 1).

### Author(s)

Anderson Rodrigo da Silva <anderson.agro@hotmail.com> (code adapted from the function `swc()`, package *soilwater* (Cordano *et al.*, 2012).)

### References

Genuchten, M. T. van. (1980). A closed form equation for predicting the hydraulic conductivity of unsaturated soils. *Soil Science Society of America Journal*, 44:892-898.

Mualem, Y. (1976). A new model for predicting the hydraulic conductivity of unsaturated porous media. *Water Resources Research*, 12:513-522.

### See Also

[fitsoilwater](#)

### Examples

```
# example 1
soilwater(x = 0.1, theta_R = 0.06, theta_S = 0.25, alpha = 21, n = 2.08)
curve(soilwater(x, theta_R = 0.06, theta_S = 0.25, alpha = 21, n = 2.08))

# example 2 (punctual predictions)
p <- seq(0, 1, length.out = 10)
m <- soilwater(x = p, theta_R = 0.06, theta_S = 0.25,
alpha = 21, n = 2.08)
points(m ~ p, type = "b", col = "red")

# End (not run)
```

**Description**

Function to calculate the soil water content based on the Groenevelt & Grant (2004) model. It is based on thermodynamic principles. Therefore, it is appropriate for the case in which thermodynamic equilibrium has been attained by diffusion of water. In this case, the water retention curve is given by:

$$\theta = k_1 \exp(-k_0/x_0^n) - k_1 \exp(-k_0/x^n)$$

where  $x = \log h$  (pore water suction), and  $h$  is in units of hPa

**Usage**

```
soilwater2(x, x0 = 6.653, k0, k1, n)
```

**Arguments**

x	a numeric vector containing pF values.
x0	the value of pF (pore water suction) at which the soil water content becomes zero. The default is 6.653.
k0	a parameter value.
k1	a parameter value.
n	a parameter value.

**Value**

The the soil water content.

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Groenevelt & Grant (2004). A newmodel for the soil-water retention curve that solves the problem of residualwater contents. *European Journal of Soil Science*, 55:479-485.

**See Also**

[fitsoilwater2](#), [soilwater](#)

**Examples**

```
pF <- 0:7
soilwater2(pF, k0 = 1.867, k1 = 0.426, n = 2.358)

# End (not run)
```

soilwater3

*Soil Water Retention, based on the Dexter's (2008) formula***Description**

Function to calculate the soil water content based on the Dexter's (2008) formula. It is based on the segregation of pore space in two categories what are called bi-modal pore size distributions. In this model, the pore space is divided into two parts: the textural porosity which occurs between the primary mineral particles, and the structural porosity which occurs between micro aggregates and/or any other compound particles. This is called the double-exponential (DE) water retention equation, given by:

$$\theta = \theta_R + a_1 \exp(-x/p_1) + a_2 \exp(-x/p_2)$$

where  $\theta$  is the gravimetric water content.

**Usage**

```
soilwater3(x, theta_R, a1, p1, a2, p2)
```

**Arguments**

x	a numeric vector containing the values of applied air pressure.
theta_R	a parameter that represents the residual water content.
a1	a parameter that represents the drainable part of the textural pore space in units of gravimetric water content at saturation.
p1	a parameter that represents the applied air pressures characteristic for displacement of water from the textural pore space.
a2	a parameter that represents the total structural pore space in units of gravimetric water content at saturation.
p2	a parameter that represents the applied air pressure that is characteristic for displacing water from the structural pores.

**Value**

The the soil water content.

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Dexter et al. (2008). A user-friendly water retention function that takes account of the textural and structural pore spaces in soil. *Geoderma*, 143:243-253.

**See Also**

[fitsoilwater3](#), [soilwater](#), [soilwater2](#)

**Examples**

```
soilwater3(x = 0, theta_R = 0.058, a1 = 0.233, p1 = 3.274, a2 = 0.070, p2 = 1.78)
soilwater3(x = 3, theta_R = 0.058, a1 = 0.233, p1 = 3.274, a2 = 0.070, p2 = 1.78)
soilwater3(x = 4, theta_R = 0.058, a1 = 0.233, p1 = 3.274, a2 = 0.070, p2 = 1.78)

# End (not run)
```

---

soilwater4

*Soil Water Retention, based on Power Models*


---

**Description**

Function to calculate the soil water content based on the following formulas:

$$\theta = \exp(a + b * Bd)\psi^c$$

(Silva et al., 1994)

$$\theta = a\psi^c$$

(Ross et al., 1991)

where  $\theta$  is the soil water content.

**Usage**

```
soilwater4(psi, Bd, a, b, c, model = c("Silva", "Ross"))
```

**Arguments**

psi	a numeric vector containing values of water potential (Psi).
Bd	a numeric vector containing values of dry bulk density.
a	a model-fitting parameter. See details.
b	a model-fitting parameter. See details.
c	a model-fitting parameter. See details.
model	a character; the model to be used for calculating the soil water content. It must be one of the two: "Silva" (default) or "Ross".

**Details**

The parameters "a" and "c" have the same meaning in both models, but be aware that the parameter "a" of the model employed by Silva et al. (1994) is parameter "a" of the Ross et al. (1991) in a log10 scale.

**Value**

The the soil water content.

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Ross et al. (1991). Equation for extending water-retention curves to dryness. *Soil Science Society of America Journal*, 55:923-927.

Silva et al. (1994). Characterization of the least limiting water range of soils. *Soil Science Society of America Journal*, 58:1775-1781.

**See Also**

[fitsoilwater4](#), [soilwater](#), [soilwater2](#), [soilwater3](#)

**Examples**

```
# End (not run)
```

---

soilwater5

*Soil Water Retention, based on the modified van Genuchten's formula*

---

**Description**

Function to calculate the soil water content based on the modified van Genuchten's formula, as suggested by Pierson and Mulla (1989):

$$\theta = \theta_R + (\theta_S - \theta_R)(1 + (\alpha x)^n)^{-m} + b_0 + b_1 x + b_2 * x^2$$

**Usage**

```
soilwater5(x, theta_R, theta_S, alpha, n, m = 1 - 1/n, b0, b1, b2)
```

**Arguments**

x	the matric potential.
theta_R	the residual water content.
theta_S	the water content at saturation.
alpha	a scale parameter of the van Genuchten's formula.
n	a shape parameter in van Genuchten's formula.
m	a shape parameter in van Genuchten's Formula. Default is $1 - 1/n$ (Mualem, 1976).

b0	a parameter added to the van Ganuchten's formula.
b1	a parameter added to the van Ganuchten's formula.
b2	a parameter (of quadratic term) added to the van Ganuchten's formula.

**Value**

The the soil water content or the saturation index (a value between 0 and 1).

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Pierson, F.B.; Mulla, D.J. (1989) An Improved Method for Measuring Aggregate Stability of a Weakly Aggregated Loessial Soil. *Soil Sci. Soc. Am. J.*, 53:1825–1831.

**See Also**

[fitsoilwater5](#)

**Examples**

```
soilwater5(x = 20, theta_R = 0.2735, theta_S = 0.478, alpha = 0.1029,
n = 9.45, b0 = 0.2278, b1 = -0.0165, b2 = 0.000248)
curve(soilwater5(x, theta_R = 0.2735, theta_S = 0.478, alpha = 0.1029,
n = 9.45, b0 = 0.2278, b1 = -0.0165, b2 = 0.000248),
from = 0, to = 40,
ylab = "Water content",
xlab = "Matric potential")

# End (Not run)
```

---

SSlbc

*Self-Starting Nls Load Bearing Capacity Model*

---

**Description**

A [selfStart](#) model that evaluates the Load Bearing Capacity (Dias Jr., 1994) function and its gradient. It has an initial attribute that creates initial estimates of the parameters b0 and b1.

**Usage**

```
SSlbc(theta, b0, b1)
```

**Arguments**

theta	a numeric vector of soil moisture values at which to evaluate the model.
b0	a numeric parameter.
b1	a numeric parameter.

**Value**

a numeric vector with the same length of theta. It is the value of the expression  $10^{(b_0+b_1*\theta)}$ . Also, the gradient matrix with respect to the parameters is attached as an attribute named *gradient*.

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**References**

Dias Junior, M. S. (1994). *Compression of three soils under longterm tillage and wheel traffic*. 1994. 114p. Ph.D. Thesis - Michigan State University, East Lansing.

**See Also**

[getInitiallbc](#), [fitlbc](#), [selfStart](#), [nls](#), [sigmaP](#)

**Examples**

```
data(compaction)
attach(compaction)
ss <- SS1bc(Mois, 2.79, -2.33)
ss[1:50] # prediction
PS # original data of preconsolidation stress
ss # prediction and gradient

# End (not run)
```

---

stressTraffic

*Predicting Soil Stress Due to Agricultural Trafficability*

---

**Description**

Contact area, stress distribution and stress propagation based on the SoilFlex model (Keller 2005; Keller et al. 2007) are calculated.

**Usage**

```
stressTraffic(inflation.pressure, recommended.pressure, tyre.diameter,
             tyre.width, wheel.load, conc.factor, layers, plot.contact.area = FALSE, ...)
```



**Arguments**

inflation.pressure	tyre inflation pressure, kPa
recommended.pressure	recommended tyre inflation pressure at given load, kPa
tyre.diameter	overall diameter of the unloaded tyre, m
tyre.width	tyre width, m
wheel.load	wheel load, kg
conc.factor	concentration factor; a numeric vector ranging from 3 (wet soil) to 6 (dry soil), depending on water content.
layers	a numeric vector containing values of depth (in meters) for the soil layers. Note that layers can also be a unique value
plot.contact.area	logical; shall soilTraffic plot the distribution of stress over the contact area?
...	further graphical arguments. See <a href="#">par</a> .

**Value**

A list of

Area	Contact area parameters.
Loads	Estimated wheel loads.
Stress	Stress propagation into soil; sigma_vertical: vertical stress; sigma_mean: mean normal stress
stress.matrix	The matrix of applied stress at a specific depth and radial distance from the tyre centre.
fZStress	The function of stress propagation in z direction (vertical stress).
fmeanStress	The function of mean normal stress propagation.
fStress	The function of stress propagation.
fXStress	The function of stress propagation in x (footprint length or driving) direction.
fYStress	The function of stress propagation in y (tire width) direction.

**Author(s)**

Renato Paiva de Lima <renato\_agro\_@hotmail.com>  
 Anderson Rodrigo da Silva <anderson.agro@hotmail.com>  
 Alvaro Pires da Silva <apisilva@usp.br>

**References**

Keller, T. 2005. A model to predict the contact area and the distribution of vertical stress below agricultural tyres from readily-available tyre parameters. *Biosyst. Eng.* 92, 85-96.  
 Keller, T.; Defosse, P.; Weisskopf, P.; Arvidsson, J.; Richard, G. 2007. SoilFlex: a model for prediction of soil stresses and soil compaction due to agricultural field traffic including a synthesis of analytical approaches. *Soil and Tillage Research* 93, 391-411.

**Examples**

```

stress <- stressTraffic(inflation.pressure=200,
  recommended.pressure=200,
  tyre.diameter=1.8,
  tyre.width=0.4,
  wheel.load=4000,
  conc.factor=c(4,5,5,5,5,5),
  layers=c(0.05,0.1,0.3,0.5,0.7,1),
  plot.contact.area = TRUE)

stress

# Building a fancier plot for the contact area
# library(fields)
# image.plot(x = as.numeric(rownames(stress$stress.matrix)),
#           y = as.numeric(colnames(stress$stress.matrix)),
#           z = stress$stress.matrix,
#           xlab="Tyre footprint length (m)", ylab="Tyre width (m)")
# End (not run)

# Stress Propagation
# Vertical Stress
stress.v <- stress$Stress$sigma_vertical
layers <- stress$Stress$Layers
plot(x = 1, y = 1, xlim=c(0,300),ylim=c(1,0),xaxt = "n",
     ylab = "Soil Depth",xlab = "", type="l", main="")
axis(3)
mtext("Stress (kPa)",side=3,line=2.5)
lines(x=stress.v, y=layers)

# Mean normal stress
stress.p <- stress$Stress$sigma_mean
lines(x=stress.p, y=layers, lty=2)
legend("bottomright", c("Vertical stress", "Normal mean stress"), lty = 1:2)

# End (not run)

```

---

voidratio

*Void Ratio*


---

**Description**

A function to calculate the soil void ratio.

**Usage**

```

voidratio(wetsoil, drysoil, diam.cylinder, height.cylinder,
  dens.particle, deformation)

```

**Arguments**

wetsoil            the weight of wet soil.  
drysoil            the weight of dry soil.  
diam.cylinder    the diameter of the cylinder.  
height.cylinder            the height of the cylinder.  
dens.particle    the particle density.  
deformation      a numeric vector containing soil deformation values.

**Value**

A numeric vector with same length of deformation containig void ratio values.

**Author(s)**

Anderson Rodrigo da Silva <anderson.agro@hotmail.com>

**See Also**

[sigmaP](#)

**Examples**

```
def <- c(0, 0.0230, 0.0352, 0.0605, 0.1070, 0.1750, 0.2525, 0.3395, 0.4250)
pres <- c(1, 12.5, 25, 50, 100, 200, 400, 800, 1600)
VR <- voidratio(wetsoil = 170.62, drysoil = 134.08, diam.cylinder = 6.95,
               height.cylinder = 2.5, dens.particle = 2.61, def)
VR

plot(VR ~ pres, type = "b",
     ylab = "Void ratio",
     xlab = "Applied stress (kPa)",
     main = "Compression curve",
     log = "x")

# End (not run)
```

# Index

## \* datasets

- bulkDensity, 6
  - compaction, 7
  - skp1994, 62
  - SoilAggregate, 62
- aggreg.stability, 4
- aggreg.stability\_App, 5
- aov, 54, 55
- bulkDensity, 6
- colors, 51
- compaction, 7
- compressive\_properties, 7
- compressive\_properties2, 9
- compressive\_properties3, 10
- compressive\_properties4, 12
- compressive\_properties5, 13
- criticalmoisture, 15, 36
- curve, 48
- fitbusscher, 16, 18, 39
- fitlbc, 17, 18, 58, 80
- fitsoilwater, 19, 33, 61, 74
- fitsoilwater2, 20, 21, 29, 75
- fitsoilwater3, 21, 29, 30, 77
- fitsoilwater4, 22, 23, 78
- fitsoilwater5, 23, 28, 79
- formula, 25
- fun2form, 25
- function, 25, 48
- getInitial, 26
- getInitiallbc, 26, 80
- hemc, 27
- hydraulicCutOff, 28, 30
- hydraulicCutOff2, 29
- iwc, 31
- Kr\_h, 33
- Kr\_theta, 34
- liquidlimit, 36
- llwr, 33, 37
- LLWR\_App, 42
- llwr\_llmpr, 43
- LLWR\_LLMPR\_App, 46
- llwrPTF, 40
- lm, 54, 55
- maxbulkdensity (criticalmoisture), 15
- maxcurv, 16, 18, 47, 58
- nls, 16–22, 24, 26, 47, 54, 55, 80
- par, 19–21, 24, 30, 32, 51, 63, 81
- particle.sedimentation, 48
- particle.sedimentation\_App, 50
- plot.default, 32
- plotCIsigmaP, 50, 59
- PredComp, 52
- predict.nls, 17
- psd, 52
- r, 53
- r\_App, 55
- Rsqr, 17, 18, 54
- selfStart, 79, 80
- sigmaP, 18, 26, 51, 56, 59, 80, 83
- simSigmaP, 50, 51, 59
- Sindex, 60
- skp1994, 62
- SoilAggregate, 62
- soilDeformation, 63
- soilphysics (soilphysics-package), 3
- soilphysics-package, 3
- soilStrength, 66
- soilStrength2, 68
- soilStrength3, 69

soilStrength4, 71  
soilStrength5, 72  
soilwater, 19, 20, 23, 33, 61, 73, 75, 77, 78  
soilwater2, 20, 23, 74, 77, 78  
soilwater3, 21, 23, 76, 78  
soilwater4, 77  
soilwater5, 24, 78  
SSlbc, 26, 79  
stressTraffic, 6, 8, 9, 11, 12, 14, 43, 46, 50,  
52, 56, 67, 69, 70, 72, 73, 80  
summary.lm, 55  
summary.nls, 17  
voidratio, 58, 82